APPLICATIONS OF APPROXIMATION SCHEMES TO EVOLUTION EQUATIONS

ADRIAN DUMA AND CONSTANTIN P. NICULESCU

Abstract. We discuss here various properties of the nonlinear $A-$proper operators as well as a generalized Leray-Schauder Principle. Also, a method of approximating arbitrary continuous operators by $A-$proper mappings is described. We construct, via apropriate Browder-Petryshyn approximation schemes, approximative solutions for linear evolution equations in Banach spaces.

The purpose of this paper is to describe some approximation structures for nonlinear operators in Banach spaces as well as a number of applications to evolution equations. In sections 1 and 2 we prove results concerning the nonlinear $A-$proper operators, such as the positive decomposition, the $A-$properness of the Dirac mass operator, and a method of approximating arbitrary continuous operators by $A-$proper mappings. We also prove a generalized Leray-Schauder Principle via the $A-$proper mapping theory.

The third section is devoted to the existence of approximative solutions for linear evolution equations in Banach spaces. That is done via the same approximation schemes as in the case of $A-$proper operators. Our results complements somehow the classical results on C_0-semigroups and show what happens beyond the standard hypotheses.

Given a separable Banach space E with Schauder basis, we construct (via an approximation scheme), a linear operator A on E such that the dicerential equation

$$\begin{aligned} \frac{du}{dt} &= Au; \\ u(0) &= u_0 \end{aligned}$$

admits a sequence $(u_n)_n$ of C^∞ solutions for each $u_0 \in E$, $u_0 \neq 0$, such that $(u_n)_n$ converges uniformly to a C^∞ function u and $\left(\frac{du_n}{dt} - Au_n\right)_n$ converges uniformly to 0, but u is not a solution.

A complementary phenomenon is described in Theorem 3.2 below. For $E = C([0,1] \times [0,1])$, one shows the existence of a wildly discontinuous linear operator $A : E \to E$, for which the initial value problem

$$\begin{aligned} \frac{du}{dt} &= Au + f, \quad t \in [0,T] \\ u(0) &= u_0 \end{aligned}$$

1991 Mathematics Subject Classiøcation. 35F10, 47H09, 47H11.
Key words and phrases. $A-$proper mapping, evolution equation, Browder-Petryshyn approximation scheme.
Partially supported by CNCSIS Grant 303/1999.

has a generalized solution whatever are $f \in C^1([0,T], E)$, $u_0 \in E$ and $T > 0$.

1. $A-$properness via approximation schemes

Let X and Y be two separable Banach spaces and let

$$\Gamma = (\{X_n\}, \{Y_n\}, \{P_n\}, \{Q_n\})$$

be an approximation scheme, where $X_n \subset X$, $Y_n \subset Y$ are linear subspaces with $\dim X_n = \dim Y_n < \infty$, $P_n : X_n \to X$ are the canonical isometries and $Q_n : Y \to Y_n$ are continuous operators such that $Q_n y \to y \quad (y \in Y)$.

An operator $T : X \to Y$ is named $A-$proper with respect to Γ provided that the operators $T_n = Q_n T P_n$ are continuous and any bounded sequence $\{x_k; x_k \in X_{n_k}\}$ such that $T_{n_k} x_k \to f$, where $f \in Y$ has a subsequence $\{x_{k_j}\}$ so that $x_{k_j} \to x_0$ and $Tx_0 = f$.

Definition 1.1. The scheme Γ is called of type (C) if it satisfies the following conditions:

a) $X_n \subsetneq X_{n+1}$; $Y_n \subsetneq Y_{n+1}$ for any n;
b) $Q_n \in L(Y)$ are projectors such that $R(Q_n) = Y_n$;
c) $m \leq n$ implies $Q_m Q_n = Q_m$.

Theorem 1.1. Suppose that Γ is of type (C). Then there exists a continuous noncompact operator $S : X \to Y$, with bounded $R(S)$, such that for any $A-$proper operator $T : X \to Y$ which is uniformly continuous on bounded subsets, the sum-operator $T + S$ is $A-$ proper, too .

Proof. Let $\delta \in (0,1)$. We may construct the sequences $\{x_i\} \subset X$ and $\{z_i\} \subset Y$ such that

$$\|x_i\| = \|z_i\| = 1,$$
$$x_i \in X_i \ ; \quad z_i \in Y_i$$

and

$$d(x_i, X_j) \geq \delta, \quad d(z_i, Y_j) \geq \delta \quad (\text{for } j < i).$$

We define $y_i \in Y$ by

$$y_i = z_i - Q_{i-1} z_i.$$

Then

$$\begin{aligned} \sup \|y_i\| &\leq 1 + \sup \|Q_i\| < +\infty \\ y_i &\in Y_i \ , \ d(y_i, Y_j) \geq \delta \quad (j < i) \end{aligned}$$

and

$$Q_m y_k = \begin{cases} y_k, & \text{if} \quad m \geq k \\ 0, & \text{if} \quad m < k \end{cases}$$

.

Choose now $\mu \in (0, \delta/2)$ and the sequence $\{\varepsilon_n\} \subset (0, \mu]$ such that $\varepsilon_n \to 0$. We define the functions $\{\varphi_i\} \subset C(X, \mathbb{R})$ by

$$\varphi_i(x) = \max\left(0, 1 - \varepsilon_i^{-1} \|x - x_i\|\right) \qquad (x \in X).$$

Clearly,

$$i \neq j \ \Rightarrow \ supp\, \varphi_i \cap supp\, \varphi_j = \emptyset$$

and $0 \leq \varphi_i(x) \leq 1$ for all $i \in \mathbb{N}^\star$, and all $x \in X$.

We introduce the operator $S : X \to Y$, deøned by

$$S(x) = \sum_{i=1}^{\infty} \varphi_i(x)\, y_{i+1}.$$

Then

$$\sup_{x\in X} \|S(x)\| \le \sup_{n} \|y_n\| < +\infty.$$

We claim that S is continuous. Indeed, let $u_0 \in X$. We have the following two alternatives:

a) $\|u_0 - x_i\| > \varepsilon_i$ for any $i \in \mathbb{N}^\star$. We claim that there exists $\lambda > 0$ such that $u \in B(u_0, \lambda)$ implies

$$\|u - x_i\| > \varepsilon_i \qquad (i \in \mathbb{N}^\star).$$

Indeed, if this is not true, then we can ønd $n \in \mathbb{N}^\star$ and $u_1 \in B(u_0, \alpha)$ such that $\|u_1 - x_n\| \le \varepsilon_n$, where $\alpha \in (0, \delta/2 - \mu)$. Let us choose now a $\beta > 0$ with $\beta < \min\{\alpha, \|u_0 - x_n\| - \varepsilon_n\}$. Then there exist $u_2 \in B(u_0, \beta)$ and $m \in \mathbb{N}^\star$ such that $\|u_2 - x_m\| \le \varepsilon_m$.

We obtain

$$\|u_0 - x_n\| \le \alpha + \varepsilon_n, \quad \|u_0 - x_m\| \le \beta + \varepsilon_m,$$

hence

$$\|x_m - x_n\| \le \alpha + \beta + \varepsilon_m + \varepsilon_n \le 2\mu + 2\alpha < \delta$$

and thus $m = n$. Then we have

$$\|u_0 - x_n\| \le \beta + \varepsilon_n < \|u_0 - x_n\|,$$

a contradiction. Consequently, there exists $\lambda > 0$ such that

$$\varphi_i(u) = 0 \quad (\forall)\ i \in \mathbb{N}^\star,\, u \in B(u_0, \lambda)$$

so that $S\big|_{B(u_0,\lambda)} = 0$ and S is continuous at u_0.

b) There exists a (unique) $j \in \mathbb{N}^\star$ such that $\|u_0 - x_j\| \le \varepsilon_j$.

Let $\gamma \in (0, \delta - 2\mu)$. Then

$$S(u) = \varphi_j(u)\, y_{j+1} \quad \text{for } u \in \overline{B}(u_0, \gamma)$$

so that S is continuous at u_0.

Now, let $T : X \to Y$ be an $A-$proper operator which is uniformly continuous on bounded subsets. We claim that the operator $T+S$ is $A-$proper too. Let $\{n_k\} \subset \mathbb{N}^\star$ be such that $n_k \to \infty$ and let $\{w_{n_k}\,;\, w_{n_k} \in X_{n_k}\}$ be a bounded sequence such that

$$T_{n_k} w_{n_k} + S_{n_k} w_{n_k} \to f,$$

for some $f \in Y$.

Notice that

$$\varphi_m | X_{n_k} = 0 \quad (m > n_k)$$

so that

$$S(x) = \sum_{i=1}^{n_k} \varphi_i(x)\, y_{i+1} \qquad \text{on } X_{n_k}$$

which yields

$$S_{n_k}(x) = \sum_{i=1}^{k-1} \varphi_i(x)\, y_{i+1},$$

because $Q_{n_k} y_{n_k+1} = 0$.

Then for each $k \in \mathbb{N}^\star$ we may choose an $i_k \in \mathbb{N}^\star$, $1 \le i_k \le n_k - 1$, such that

$$S_{n_k}(w_{n_k}) = \varphi_{i_k}(w_{n_k})\, y_{i_k+1}.$$

The following two cases are possible:

1) There exist $\{k_l\} \subset \mathbb{N}^\star$, $k_l \underset{l}{\longrightarrow} \infty$ and $m \in \mathbb{N}^\star$ such that $i_{k_l} < m \quad (l \ge 1)$. Then we obtain

$$S_{n_{k(l)}}\left(w_{n_{k(l)}}\right) \in \bar{B}\left(0, \sup \|y_i\|\right) \cap sp\left\{y_j; j = \overline{2, m}\right\}.$$

Now, T being $A-$proper, it follows that $\left\{w_{n_{k(l)}}\right\}_l$ has no convergent subsequence and, since $T+S$ is continuous, it will be $A-$proper too.

2) $i_k \to \infty \quad (k \to \infty)$. Now, if there exists a sequence $\{k_j\} \subset N^\star$ such that $k_j \to +\infty \quad (j \to +\infty)$ and $\varphi_{i_{k(j)}}\left(w_{n_{k(j)}}\right) > 0 \qquad (j \ge 1)$ then we obtain

$$\left\| w_{n_{k(j)}} - x_{i_{k(j)}} \right\| < \varepsilon_{i_{k(j)}} \to 0 \qquad (j \to \infty).$$

We also have

$$S_{n_{k(j)}}\left(w_{n_{k(j)}}\right) = \varphi_{i_{k(j)}}\left(w_{n_{k(j)}}\right) y_{i_{k(j)}+1},$$

hence

$$(*) \qquad S_{i_{k(j)}}\left(w_{n_{k(j)}}\right) = \varphi_{i_{k(j)}}\left(w_{n_{k(j)}}\right) Q_{i_{k(j)}} y_{i_{k(j)}+1} = 0$$

Now, since T is uniformly continuous on bounded subsets, we have

$$(**) \qquad \left\| T w_{n_{k(j)}} - T x_{i_{k(j)}} \right\| \to 0.$$

We also have

$$Q_{i_{k(j)}}\left[T_{n_{k(j)}} w_{n_{k(j)}} + S_{n_{k(j)}} w_{n_{k(j)}}\right] \to f$$

and, from $(\star)$, $T_{i_{k(j)}} w_{n_{k(j)}} \to f$ and, from $(\star\star)$, $T_{i_{k(j)}} x_{i_{k(j)}} \to f$, which is impossible, because T is $A-$proper and $\{x_n\}$ has no convergent subsequence.

Then there exists $k_0 \in \mathbb{N}^\star$ such that $k \ge k_0$ implies $\varphi_{i_k}(w_{n_k}) = 0$, consequently, $S_{n_k}(w_{n_k}) = 0$, so that

$$T_{n_k}(w_{n_k}) \to f.$$

Thus, $\{w_{n_k}\}$ has convergent subsequences and, since $T+S$ is continuous, it will be $A-$proper, too.

Finally, let us remark that $Sx_i = y_{i+1} \quad (i \in \mathbb{N}^\star)$, whence $S\left(\bar{B}(0,1)\right)$ is not relatively compact. ■

Theorem 1.2. Let E be a separable Banach space such that $E^\star$ is endowed with a sequence of ønite-dimensional subspaces $\{X_n\}_{n \ge 1}$, with the following properties:

a) for every n there is a projection $\Pi_n \in L(E^\star)$ with $R(\Pi_n) = X_n$;

b) $\Pi_n x \to x \quad \forall x \in E^\star$;

c) there exists an (algebraic-topological) isomorphism into $U : E^\star \to C(S_{E^\star})^\star$, where $S_{E^\star}$ denotes the closed unit ball of $E^\star$, endowed with the weak-star topology.

Then the operator $S_{E^\star} \to C(S_{E^\star})^\star$, $x \to \delta_x$ (the Dirac mass) is $A-$proper from $S_{E^\star} \subset E^\star$ to $C(S_{E^\star})^\star$.

Proof. For every $n \in \mathbb{N}^\star$ we choose $\mu_n \in U(X_n)$ with $\|\mu_n\| = n$. Let us define the finite dimensional subspace $Y_n \subset C(S_{E^\star})^\star$ by $Y_n = U(X_n)$. Then clearly $\dim Y_n = \dim X_n < +\infty$. We also introduce the sequence of continuous operators $Q_n : C(S_{E^\star})^\star \to Y_n$, given by

$$Q_n \nu = U \Pi_n j^\star \nu + dist\,(\nu, \{\varepsilon_x; x \in S_{E^\star}\})\ \mu_n,$$

where $j : E \to C(S_{E^\star})$ is the canonical isometrical embedding.

We claim that the operator $T : S_{E^\star} \subset E^\star \to C(S_{E^\star})^\star$; $Tx = \varepsilon_x$, is $A-$proper with respect to the approximation scheme

$$\Gamma = (\{X_n\}, \{P_n\}, \{Y_n\}, \{Q_n\}),$$

where $P_n : X_n \to E^\star$ are the canonical isometries.

Clearly, if $x \in S_{E^\star} \cap X_n$, then we have $Q_n T x = Ux$, hence the operator $Q_n T\,|_{S_{E^\star} \cap X_n}$ is continuous.

Now, let $\{x_m; x_m \in S_{E^\star} \cap X_m\}$ be a sequence such that

$$\|Q_m T x_m - Q_m \mu\| \to 0$$

for some $\mu \in C(S_{E^\star})^\star$. Then we obtain

$$\|Ux_m - U\Pi_m j^\star \mu + dist\,(\mu, \{\varepsilon_x; x \in S_{E^\star}\})\,\mu_m\| \to 0;$$

$$dist\,(\mu, \{\varepsilon_x; x \in S_{E^\star}\}) \cdot m \le \|U\| \left(1 + \sup_m \|\Pi_m\| \cdot \|\mu\|\right),$$

therefore there is an $x \in S_{E^\star}$ such that $\mu = \delta_x$. It follows that

$$\|Ux_m - U\Pi_m x\| \to 0,$$

thus

$$x_m \to x$$

and clearly $Tx = \mu$, which completes the proof. ■

2. The generalized Leray-Schauder Principle

Let X be a Banach space and let D be an open bounded subset of X with $0 \in D$. We say that $\Gamma \subset D$ is a pseudo-boundary of D if there exist a Banach space Y, a open bounded subset $D_\star \subset Y$ and a continuous $A-$proper operator $A : \overline{D_\star} \to \overline{D}$ with $Deg\,(A, D_\star, 0) \neq \{0\}$ such that $\Gamma = A(\partial D_\star)$.

Theorem 2.1. Let $K : \overline{D} \to X$ be a compact operator. Suppose that there exists a pseudo-boundary Γ of D such that

$$Kx \neq \lambda\, x \qquad (\lambda > 1;\ x \in \Gamma).$$

Then K has at least one fixed point.

Proof. Let us consider the $A-$proper homotopy $H : [0,1] \times \overline{D_\star} \to X$, given by

$$H(t,z) = Az - tKAz.$$

Then $H(t,z) = 0$ with $t \in (0,1)$ and $z \in \partial D_\star$ implies $Kx = \lambda\, x$ with $\lambda = t^{-1}$ and $x = Az \in \Gamma$, a contradiction. Clearly, $0 \notin H(\{0\} \times \partial D_\star)$. If $H(1,z) = 0$ with

$z \in \partial D_\star$, we obtain that Az is a fixed point of K. Finally, if $0 \notin H([0,1] \times \partial D_\star)$ then

$$\begin{aligned} Deg\,(A - KA,\, D_\star,\, 0) &= Deg\,(H(1,\cdot)\,,\, D_\star,\, 0) = Deg\,(H(0,\cdot)\,,\, D_\star,\, 0) \\ &= Deg\,(A,\, D_\star,\, 0) \neq \{0\}\,, \end{aligned}$$

which assures the existence of a $z \in D_\star$ such that $(A - KA)(z) = 0$. Consequently, $Az \in \overline{D}$ is a fixed point of K and the theorem is proved. ■

In what follows, X will be a Banach space and $\Delta \subset X$ will be an open bounded subset satisfying the following property:

($\mathcal{P}$) There is a finite-dimensional subspace $V \subset X$, a point $p \in \overline{\Delta_V} = \overline{\Delta} \cap V$ and a sequence of linear subspaces $X_n \subset V$ such that:

a) the sequence $\overline{\Delta_n} = \overline{\Delta} \cap X_n$ converges to $\{p\}$ in the hyperspace $2^{\overline{\Delta_V}}$ of $\overline{\Delta_V}$ (e.g., see [11]);

b) $\overline{\Delta_m} \cap \overline{\Delta_n} = \emptyset \qquad (m \neq n)\,;$

c) $p \notin \overline{\Delta_n} \qquad (n \in \mathbb{N}^\star)$.

We have the following approximation result:

Theorem 2.2. For each continuous operator $T : \overline{\Delta} \subset X \to \ell_2$ there exists a sequence of operators $\left\{T_n\,; T_n : \overline{\Delta} \to \ell_2\right\}$ that are $A-$proper with respect to various approximation schemes for the pair $(X,\, \ell_2)$ such that

$$\sup\, \left\{\|T_n x - Tx\|\ ;\ x \in \overline{\Delta}\,\right\} \underset{n\to\infty}{\longrightarrow} 0.$$

Proof. Let $Q := T\left(\overline{\Delta_V}\right)$. Since$T$ is continuous and $\overline{\Delta_V}$ is compact, so it is Q. Let us take $\varepsilon\ > 0$. We denote by $\{e_n\}_{n\geq 1}$ the standard orthonormal basis of ℓ_2 and by $\{\Pi_n\}_{n\geq 1}$ the associated sequence of ortho-projectors. Then there exists $k \in \mathbb{N}^\star$ such that

$$\sup_{x\in Q}\ \|\Pi_k x - x\|\ < \frac{\varepsilon}{2}.$$

Now we define the operator $\tilde{T} : \overline{\Delta} \to \ell_2$, given by the formula

$$\tilde{T}x = \begin{cases} Tx, & \text{if } x \in \overline{\Delta} \setminus \left(\bigcup\limits_{n\geq 1} \overline{\Delta_n}\right) \\ \Pi_k Tx + \frac{\varepsilon}{2} e_{k+m}, & \text{if } x \in \overline{\Delta_m} \quad (m \in \mathbb{N}^\star)\,. \end{cases}$$

Then it is clear that

$$\sup_{x\in\overline{\Delta}}\ \left\|\tilde{T}x - Tx\right\|\ < \varepsilon.$$

Let us observe that, if $x \in \overline{\Delta_m}$ and $y \in \overline{\Delta_n}$ $(m \neq n)$, then

$$\left\|\tilde{T}x - \tilde{T}y\right\|\ \geq \frac{\varepsilon}{\sqrt{2}}.$$

Let us choose now $f \in \ell_2 \backslash \{0\}$ and a sequence of subspaces $\{Y_n\}_{n\geq 1}$, $Y_n \subset \ell_2$, such that

$$\dim\ Y_n = \dim\ X_n$$

and $f \in Y_n$ $(n \in \mathbb{N}^\star)$. We then define a sequence of continuous nonlinear operators

$$\{Q_n\,;\, Q_n : \ell_2 \to Y_n\}_{n\geq 1}$$

where

$$Q_n x = dist\left(x, \tilde{T}\left(\overline{\Delta_n}\right)\right) \cdot \|x - Tp\| \cdot f.$$

We claim that the operator $\tilde{T}$ is $A-$proper with respect to the approximation scheme

$$\Gamma = (\{X_n\}, \{P_n\}, \{Y_n\}, \{Q_n\}),$$

where $P_n : X_n \to X$ are the canonical isometries. Indeed, let us observe that, if $x \in \overline{\Delta_n}$, then $Q_n\tilde{T}x = 0$. On the other hand, let $\left\{x_m \,;\, x \in \overline{\Delta_m}\right\}$ be a sequence such that

$$\left\|Q_m\tilde{T}x_m - Q_m y\right\| \to 0,$$

for some $y \in \ell_2$. It follows that necessarily $y = Tp\ \left(= \tilde{T}p\right)$. Moreover, the property $(\mathcal{P})$ implies that $x_m \to p$. The proof is complete. ■

Remark. The above argument can be easily modiøed in order to conclude that all the operators T_n are $A-$proper with respect to the same approximation scheme.

In what follows, BCA denotes the class of all bounded continuous $A-$proper operators with respect to a given approximation scheme.

Proposition 2.3. Let X and Y be two Banach spaces, Y being ordered by the cone Y^+, such that int $Y^+ \neq \emptyset$. Let $D \subset X$ be a open bounded non-empty subset. Then for each operator $A \in BCA\left(\overline{D}, Y\right)$ there exist two operators $A_\pm \in BCA\left(\overline{D}, Y^+\right)$ such that

$$A = A_+ - A_-$$

and

$$A_\pm^{-1}(\{0\}) = A^{-1}(\{0\}).$$

Proof. Since int $Y^+ \neq \emptyset$, there exist $y_0 \in Y$ and $r > 0$ such that

$$\overline{B}(y_0, r) \subset Y^+.$$

Let $r^\star = \min\left\{r, \frac{1}{2}\|y_0\|\right\}$. Since clearly $y_0 \neq 0$, it follows that $r^\star > 0$. Then we deøne the operators $A_\pm$ by

$$A_\pm x = \frac{1}{2}\left(\pm Ax + \frac{\|Ax\|}{r^\star}y_0\right) \qquad \left(x \in \overline{D}\right).$$

It is easy to see that $A_\pm \in BCA\left(\overline{D}, Y\right)$, because $\lambda A + K \in BCA\left(\overline{D}, Y\right)$ for each $\lambda \in \mathbb{R} \setminus \{0\}$ and $K \in K\left(\overline{D}, Y\right)$ (= the space of all compact maps). Moreover, $A_\pm x \geq 0\ \left(x \in \overline{D}\right)$, because $\overline{B}(y_0, r^\star) \subset Y^+$. Clearly, $A = A_+ - A_-$. Because

$$\frac{1}{2}\left(\frac{\|y_0\|}{r^\star} - 1\right)\|Ax\| \ \leq\ \|A_\pm x\| \ \leq\ \frac{1}{2}\left(\frac{\|y_0\|}{r^\star} + 1\right)\|Ax\| \ \left(x \in \overline{D}\right),$$

we conclude that $A^{-1}(\{0\}) = A_\pm^{-1}(\{0\})$. The proof is done. ■

3. Approximative solutions for evolution equations

Our ørst goal is to show that for every separable Banach space E with Schauder basis there exists a linear operator $A : E \to E$ with the following two properties:

A1) The problem

$$(P): \quad \frac{du}{dt} = Au \qquad (t \geq 0)$$

has a solution u of class C^{∞} for each choice of the initial datum $u(0) = u_0$ in E;

A2) For each $u_0 \in E,\ u_0 \neq 0$, there exist u and $\{u_n\} \subset C^{\infty}(\mathbb{R}^+, E)$ such that

$$\sup_{t \geq 0} \| u_n(t) - u(t) \| \to 0, \quad u(0) = u_0\,;$$

$$\sup_{t \geq 0} \left\| \frac{du_n}{dt}(t) - A(u_n(t)) \right\| \to 0\,;$$

$$\frac{du}{dt}(t) \neq A(u(t)) \qquad (t \geq 0)\,.$$

In order to prove this result we shall need the following construction: Let $\{e_n\}$ be a Schauder basis for E, let $E_0 = span\,\{e_n\}$ and let $\{e_n\} \cup \{b_x\,;\ x \in E_0\}$ be a Hamel basis for E. We deøne the linear operator $A : E \to E$ by $Ae_n = 0 \quad (n \in \mathbb{N}^\star)\,; \quad Ab_x = x \quad (x \in E_0)$. It is easy to see that $A^2 = 0$.

Proof of A1). For each $u_0 \in E$ the function $u(t) = u_0 + t\,Au_0 \quad (t \geq 0)$ is a $C^{\infty}-$solution of (P), with $u(0) = u_0$.

Proof of A2). Let us consider the sequence of linear continuous projections

$$P_n x = \sum_{k=1}^{n} e_k^\star(x)\, e_k \qquad (x \in E;\ n \in \mathbb{N}^\star)$$

where $\{e_n^\star\} \subset E^\star$ is the associated sequence of coeŒcient functionals to $\{e_n\}$.

Let us choose a sequence of integers $\{k_n\} \subset \mathbb{N}^\star,\ k_n \to \infty$, such that

$$\| P_{k_n} b_{-P_n u_0} - b_{-P_n u_0} \| < n^{-1} \qquad (n \in \mathbb{N}^\star)$$

and consider the functions

$$u_n(t) = e^{-t}\left[b_{-P_n u_0} - P_{k_n}\left(b_{-P_n u_0} - u_0\right)\right] \quad (t \geq 0)$$

and

$$u(t) = e^{-t} u_0 \qquad (t \geq 0)\,.$$

Then, letting $c_n = \| b_{-P_n u_0} - P_{k_n} b_{-P_n u_0} + P_{k_n} u_0 - u_0 \| \to 0$, we have

$$\sup_{t \geq 0} \| u_n(t) - u(t) \| \leq e^{-t} c_n;$$

$$\frac{du_n}{dt}(t) = -u_n(t)\,;$$

$$A(u_n(t)) = -e^{-t} P_n u_0;$$

$$\begin{aligned}\left\|\frac{du_n}{dt}(t)-A(u_n(t))\right\| &= \\ &= e^{-t}\left\|-b_{-P_n u_0}+P_{k_n}(b_{-P_n u_0}-u_0)+P_n u_0\right\| \\ &\le \left\|-b_{-P_n u_0}+P_{k_n} b_{-P_n u_0}\right\| + \left\|P_n u_0 - P_{k_n} u_0\right\| \to 0\end{aligned}$$

for each $t \geq 0$.

Finally, let us suppose that

$$\frac{du}{dt}(t_0)=A(u(t_0))$$

for some $t_0 \in \mathbb{R}^+$. Then

$$\begin{aligned}-e^{-t_0}u_0 &= e^{-t_0}Au \\ -u_0 &= Au_0 \\ -Au_0 &= 0 \\ u_0 &= 0\end{aligned}$$

a contradiction. ■

In the remainder of this section, we put $I=[0,1]$, $E=C\left(I^2\right)$, and keep øxed a number $T>0$. It is known that $E \approx C(I)\,\overline{\otimes}_{\varepsilon} C(I)$ (the completion of the injective tensor product, see [8]).

We shall need the following classical result:

Theorem 3.1. (A. A. Miliutin [11]; see also [9]). Let Q be an uncountable compact metric space. Then $C(Q)$ is linearly isomorphic to $C(I)$.

Let us consider the initial value problem

$$u(t)=Au(t)+f(t), \quad t\in[0,T]$$

$$u(0)=u_0,$$

where $A: D(A)=E \to E$ is a linear operator and $f:[0,T]\to E$, $u_0 \in E$ are given .

Theorem 3.2. There exists a discontinuous linear operator $A: D(A)=E\to E$, with dense kernel, such that for each $f \in C^1([0,T], E)$ and each $u_0 \in E$ there exist u , $u_1, u_2, \ldots \in C^1([0,T], E)$ such that $A\circ u_k \in C([0,T], E)$ and

$$\begin{aligned}u_k &\to u \quad \text{uniformly,} \\ \frac{du_k}{dt}-Au_k &\to f \quad \text{uniformly}\end{aligned}$$

and

$$u_k(0)\to u_0 \quad \text{in } E.$$

Proof. Using Miliutin's Theorem, we obtain an isomorphism $U: C(I)\to E$. Let $\{b_n\}$ be a normalized Schauder base of $C(I)$. Then there is an uncountable subset Γ of $C(I)$ such that $B=\{b_n\}\cup\Gamma$ is a Hamel base of $C(I)$. Let $\{c_n\}\subset\Gamma$ be such that $c_m \neq c_n \quad (m\neq n)$. We put $e(t)=1 \quad (t\in I)$. Then we consider the linear operator $T: C(I)\to C(I)$, deøned by

$$Tc_k = ke \quad (k \geq 1);$$

$$Tb = 0 \quad (b \in B \setminus \{c_k\}).$$

We also consider the following linear operators:

$$S : C(I) \otimes C(I) \to C(I) \otimes C(I), \quad S = T \otimes id_{C(I)}$$

$$B : C(I) \otimes C(I) \to C(I), \quad B\left(\sum_i f_i \otimes g_i\right)(t) = \sum_i f_i(t)\, g_i(t)$$

and $A : E \to E$, $A = UBSP$, where $P : E \to C(I) \otimes_\varepsilon C(I)$ is an algebraic projection.

Now let $u_0 \in E$ and $f \in C^1([0,T], E)$. Since the linear subspace $F = span\,\{b_n\} \otimes_\varepsilon C(I)$ is dense in E (see [8], p. 280), there is a sequence $\{q_n\} \subset F$ such that $q_n \to u_0$. Let us remark that $F \subset Ker\ S$.

We deøne $\{u_k\} \subset C^1([0,T], E)$ by

$$u_k(t) = q_k - k^{-1} c_k \otimes U^{-1}(f(t)).$$

Then, deøning $u(t) = u_0 \quad (t \in [0,T])$, we have $u_k \to u$ uniformly, because

$$\left\| k^{-1} c_k \otimes U^{-1}(f(t)) \right\| \leq k^{-1} \left\| U^{-1} \right\| \, \|f\|_\infty \to 0.$$

We also have

$$\frac{du_k}{dt} = -k^{-1} c_k \otimes U^{-1}(f(t)),$$

so that $\frac{du_k}{dt} \to 0$ uniformly. On the other hand,

$$\begin{aligned} A(u_k(t)) &= UBSP(u_k(t)) = UBS(u_k(t)) \\ &= -k^{-1} UBS\left(c_k \otimes U^{-1}(f(t))\right) \\ &= -UB\left(e \otimes U^{-1}(f(t))\right) \\ &= -UU^{-1}(f(t)) = -f(t). \end{aligned}$$

Finally, let us remark that A has a dense kernel (since $F \subset Ker\ A$), and $A \neq 0$, because

$$A(c_1 \otimes c_1) = UBSP(c_1 \otimes c_1) = UB(e \otimes c_1) = Uc_1 \neq 0).$$

Hence A is discontinuous and the proof of our theorem is done. ■

References

[1] H.AMANN and S.A.WEISS, On the uniqueness of the topological degree, Math.Z. 130 (1973), 39-54.

[2] F.E. BROWDER, Existence of periodic solutions for nonlinear equations of evolution, Proc.N.A.S., 53 (1965) , 5, 1100-1103 .

[3] F.E. BROWDER, Fixed point theory and nonlinear problems, Bull. Amer. Math. Soc. (New Series), vol. 9, No.1, July 1983, 1-39.

[4] F.E. BROWDER and W.V.PETRYSHYN, Noncompact operators in Banach spaces, Bull. Amer. Math. Soc. 74 (1968), 641-646.

[5] F.E. BROWDER and W.V.PETRYSHYN, Approximation methods and the generalized topological degree for nonlinear mappings in Banach spaces, J. Funct.Anal. 3 (1969), 217-245.
[6] G. DA PRATO and E. SINESTRARI, On the Phillips and Tanabe Regularity Theorems, Semesterbericht Funktionalanalysis Tbingen, Band 8, Sommersemester 1985, 117-124.
[7] K. DEIMLING, Nonlinear Functional Analysis, Springer Verlag, Berlin-Heidelberg-New York, 1985.
[8] G. KTHE, Topological Vector Spaces II, Springer-Verlag, New York-Heidelberg-Berlin,1979.
[9] H.E. LACEY, The isometric theory of classical Banach spaces, Springer-Verlag, Berlin-Heidelberg-New York, 1974.
[10] N.G. LLOYD, Degree Theory, Cambridge University Press, 1978.
[11] A. A. Miliutin, Isomorphisms of the spaces of contiunous functions over compact sets of cardinality of the continuum, Teor. Funkcii Funkcional. Anal. i Priloen, 2 (1966), 150-156.
[12] J. van MILL, Inønite dimensional topology, North-Holland, Amsterdam-New York-Oxford, 1989.
[13] I. SINGER, Bases in Banach Spaces II, Springer-Verlag, Berlin-Heidelberg- New York,1981.

Department of Mathematics, University of Craiova, Str. A.I. Cuza 13, Craiova 1100, Romania

E-mail address: tempus@oltenia.ro