


Section 5.2 Logarithmic Functions

Objective 1: Understanding the Definition of a Logarithmic Function

Every exponential function of the form $f(x) = b^x$, where $b > 0$ and $b \neq 1$, is one-to-one and thus has an inverse function. The graph of $f(x) = b^x$, $b > 1$ and its inverse, f^{-1} , are shown below. Recall from Section 5.1, the graph of $f(x) = b^x$, $b > 1$ contains the points $\left(-1, \frac{1}{b}\right)$, $(0, 1)$ and $(1, b)$, and since $b^x \rightarrow 0$ as $x \rightarrow -\infty$, the x -axis is a horizontal asymptote for the graph. Recall from Section 3.6 that the graph of f^{-1} is obtained by reflecting the graph of f about the line $y = x$. Therefore, the graph of f^{-1} will contain the points $\left(\frac{1}{b}, -1\right)$, $(1, 0)$ and $(b, 1)$, and the y -axis will be a vertical asymptote for the graph.



To find the equation of f^{-1} , we begin with the process from Section 3.6:

Step 1: Change $f(x)$ to y : $y = b^x$

Step 2: Interchange x and y : $x = b^y$

Step 3: Solve for y : ??

Before we can solve for y , we must introduce the following definition:

Definition: For $x > 0$, $b > 0$ and $b \neq 1$, the **logarithmic function with base b** is defined by $y = \log_b x$ if and only if $x = b^y$.

Step 3. **Solve for y :** $x = b^y$ can be written as $y = \log_b x$

Step 4. **Change y to $f^{-1}(x)$:** $f^{-1}(x) = \log_b x$

Review of Rules of Exponents and Rewriting an Expression in the Form b^u

See Section 5.1a.

Review of Evaluating Expressions with Negative and Rational Exponents

LSU Videos “Negative Exponents” and “Rational Exponents” (0:00 – 16:30) are found on the course website.

Objective 2: Evaluating Logarithmic Expressions

The expression $\log_b x$ is the exponent to which b must be raised to in order to get x .

Objective 3: Understanding the Properties of Logarithms

General Properties of Logarithms

For $b > 0$ and $b \neq 1$,

(1) $\log_b b = 1$ and

(2) $\log_b 1 = 0$.

Cancellation Properties of Exponentials and Logarithms

For $b > 0$ and $b \neq 1$,

(1) $b^{\log_b x} = x$ and

(2) $\log_b b^x = x$.

Objective 4: Using the Common and Natural Logarithms

Definition: For $x > 0$, the **common logarithmic function** is defined by $y = \log x$ if and only if $x = 10^y$.



Definition: For $x > 0$, the **natural logarithmic function** is defined by $y = \ln x$ if and only if $x = e^y$.

Objective 5: Understanding the Characteristics of Logarithmic Functions

Characteristics of Logarithmic Functions

For $b > 0$ and $b \neq 1$, the logarithmic function with base b is defined by $y = \log_b x$.

The domain of $f(x) = \log_b x$ is $(0, \infty)$ and the range is $(-\infty, \infty)$. The graph of $f(x) = \log_b x$ has one of the following two shapes depending on the value of b :



The graph of $y = \log_b x$, $b > 0$ and $b \neq 1$ has the following properties:



1. The graph intersects the x -axis at $(1, 0)$.
2. The graph contains the points $(b, 1)$ and $(\frac{1}{b}, -1)$.
3. If $b > 1$, the graph is increasing on the interval $(0, \infty)$.
If $0 < b < 1$, the graph is decreasing on the interval $(0, \infty)$.
4. The y -axis ($x = 0$) is a vertical asymptote.
5. The function is one-to-one.

Review of Using Transformations to Graph Functions

See Section 3.4.

Objective 6: Sketching the Graphs of Logarithmic Functions Using Transformations

The graph of $f(x) = \log_2(x-1)$ can be obtained by horizontally shifting the graph of $f(x) = \log_2 x$ to the right one unit. The graph of the function $f(x) = \log_2 x$ is shown below on the left. It contains the points $\left(\frac{1}{2}, -1\right)$, $(1, 0)$ and $(2, 1)$ and has vertical asymptote $x = 0$. To shift the graph of this function right one unit, add 1 to each of the x -coordinates of the points on the graph. The resulting graph of $f(x) = \log_2(x-1)$, shown below on the right, contains the points $\left(\frac{3}{2}, -1\right)$, $(2, 0)$ and $(3, 1)$ and has vertical asymptote $x = 1$. The domain of $f(x) = \log_2(x-1)$ is $(1, \infty)$ and the range is $(-\infty, \infty)$.



Review of Solving Linear Inequalities of the form $ax + b \geq 0$

See Section 1.7.

Objective 7: Finding the Domain of Logarithmic Functions

If $f(x) = \log_b [g(x)]$, then the domain of f can be found by solving the inequality $g(x) > 0$.