

Instructions. Answer each of the questions on your own paper. Put your name on each page of your paper. Be sure to show your work so that partial credit can be adequately assessed. *Credit will not be given for answers (even correct ones) without supporting work.* A table of Laplace transforms, a table of convolutions, and the statement of the main partial fraction decomposition theorem have been appended to the exam.

1. [20 Points] Compute the inverse Laplace transform of each of the following rational functions.

(a) $F(s) = \frac{10s - 2}{(s - 1)^2(s + 3)}$

(b) $G(s) = \frac{2s + 5}{s^2 + 4s + 13}$

2. [9 Points] Compute the Laplace transform of $f(t) = te^{3t}$, $g(t) = \sin 2t$ and the convolution product $(f * g)(t)$. (Note that you are **not** asked to compute $(f * g)(t)$, only the Laplace transform of $(f * g)(t)$.)

3. [12 Points] Find the Laplace transform $Y(s)$ of the solution $y(t)$ of the initial value problem

$$4y'' + 8y' + 3y = 2te^{-5t}, \quad y(0) = 2, \quad y'(0) = -1.$$

Note that you are only asked to find the Laplace transform $Y(s)$ of $y(t)$, not $y(t)$ itself.

4. [8 Points] Find the standard basis \mathcal{B}_q of \mathcal{E}_q for $q(s) = (s - 3)^3(s^2 + 4)$.

5. [12 Points] Solve the initial value problem: $y'' + 7y' + 10y = 0$, $y(0) = 1$, $y'(0) = 1$.

6. [12 Points] Find the general solution of $4y'' - 4y' + y = 0$.

7. [12 Points] Find the general solution of $y'' + 4y' + 29y = 0$.

8. [15 Points] Find the general solution of the following differential equation using the method of undetermined coefficients:

$$y'' + y' - 2y = 6e^t.$$

Laplace Transform Table

	$f(t)$	\rightarrow	$F(s) = \mathcal{L}\{f(t)\}(s)$
1.	1	\rightarrow	$\frac{1}{s}$
2.	t^n	\rightarrow	$\frac{n!}{s^{n+1}}$
3.	e^{at}	\rightarrow	$\frac{1}{s-a}$
4.	$t^n e^{at}$	\rightarrow	$\frac{n!}{(s-a)^{n+1}}$
5.	$\cos bt$	\rightarrow	$\frac{s}{s^2+b^2}$
6.	$\sin bt$	\rightarrow	$\frac{b}{s^2+b^2}$
7.	$e^{at} \cos bt$	\rightarrow	$\frac{s-a}{(s-a)^2+b^2}$
8.	$e^{at} \sin bt$	\rightarrow	$\frac{b}{(s-a)^2+b^2}$

Laplace Transform Principles

Linearity	$\mathcal{L}\{af(t) + bg(t)\} = a\mathcal{L}\{f\} + b\mathcal{L}\{g\}$
Input Derivative Principles	$\mathcal{L}\{f'(t)\}(s) = s\mathcal{L}\{f(t)\} - f(0)$
	$\mathcal{L}\{f''(t)\}(s) = s^2\mathcal{L}\{f(t)\} - sf(0) - f'(0)$
First Translation Principle	$\mathcal{L}\{e^{at}f(t)\} = F(s-a)$
Transform Derivative Principle	$\mathcal{L}\{-tf(t)\}(s) = \frac{d}{ds}F(s)$
The Dilation Principle	$\mathcal{L}\{f(bt)\}(s) = \frac{1}{b}\mathcal{L}\{f(t)\}(s/b)$
The Convolution Principle	$\mathcal{L}\{(f * g)(t)\}(s) = F(s)G(s).$

Table of Convolutions

$f(t)$	$g(t)$	$f * g(t)$
t	t^n	$\frac{t^{n+2}}{(n+1)(n+2)}$
t	$\sin at$	$\frac{at - \sin at}{a^2}$
t^2	$\sin at$	$\frac{2}{a^3}(\cos at - (1 - \frac{a^2 t^2}{2}))$
t	$\cos at$	$\frac{1 - \cos at}{a^2}$
t^2	$\cos at$	$\frac{2}{a^3}(at - \sin at)$
t	e^{at}	$\frac{e^{at} - (1 + at)}{a^2}$
t^2	e^{at}	$\frac{2}{a^3}(e^{at} - (a + at + \frac{a^2 t^2}{2}))$
e^{at}	e^{bt}	$\frac{1}{b-a}(e^{bt} - e^{at}) \quad a \neq b$
e^{at}	e^{at}	te^{at}
e^{at}	$\sin bt$	$\frac{1}{a^2 + b^2}(be^{at} - b \cos bt - a \sin bt)$
e^{at}	$\cos bt$	$\frac{1}{a^2 + b^2}(ae^{at} - a \cos bt + b \sin bt)$
$\sin at$	$\sin bt$	$\frac{1}{b^2 - a^2}(b \sin at - a \sin bt) \quad a \neq b$
$\sin at$	$\sin at$	$\frac{1}{2a}(\sin at - at \cos at)$
$\sin at$	$\cos bt$	$\frac{1}{b^2 - a^2}(a \cos at - a \cos bt) \quad a \neq b$
$\sin at$	$\cos at$	$\frac{1}{2}t \sin at$
$\cos at$	$\cos bt$	$\frac{1}{a^2 - b^2}(a \sin at - b \sin bt) \quad a \neq b$
$\cos at$	$\cos at$	$\frac{1}{2a}(at \cos at + \sin at)$

Partial Fraction Expansion Theorems

The following two theorems are the main partial fractions expansion theorems, as presented in the text.

Theorem 1 (Linear Case). *Suppose a proper rational function can be written in the form*

$$\frac{p_0(s)}{(s - \lambda)^n q(s)}$$

and $q(\lambda) \neq 0$. Then there is a unique number A_1 and a unique polynomial $p_1(s)$ such that

$$\frac{p_0(s)}{(s - \lambda)^n q(s)} = \frac{A_1}{(s - \lambda)^n} + \frac{p_1(s)}{(s - \lambda)^{n-1} q(s)}. \quad (1)$$

The number A_1 and the polynomial $p_1(s)$ are given by

$$A_1 = \frac{p_0(\lambda)}{q(\lambda)} \quad \text{and} \quad p_1(s) = \frac{p_0(s) - A_1 q(s)}{s - \lambda}. \quad (2)$$

Theorem 2 (Irreducible Quadratic Case). *Suppose a real proper rational function can be written in the form*

$$\frac{p_0(s)}{(s^2 + cs + d)^n q(s)},$$

where $s^2 + cs + d$ is an irreducible quadratic that is factored completely out of $q(s)$. Then there is a unique linear term $B_1s + C_1$ and a unique polynomial $p_1(s)$ such that

$$\frac{p_0(s)}{(s^2 + cs + d)^n q(s)} = \frac{B_1s + C_1}{(s^2 + cs + d)^n} + \frac{p_1(s)}{(s^2 + cs + d)^{n-1} q(s)}. \quad (3)$$

If $a + ib$ is a complex root of $s^2 + cs + d$ then $B_1s + C_1$ and the polynomial $p_1(s)$ are given by

$$B_1(a + ib) + C_1 = \frac{p_0(a + ib)}{q(a + ib)} \quad \text{and} \quad p_1(s) = \frac{p_0(s) - (B_1s + C_1)q(s)}{s^2 + cs + d}. \quad (4)$$