

1152-55-67

Natalia Cadavid-Aguilar* (ncadavia@gmail.com), Instituto Politécnico Nacional 2508, Gustavo A. Madero, San Pedro Zacatenco, 07360 Ciudad de México, Mexico, and **Jesús González Espino Barros** (jesus@math.cinvestav.mx), Instituto Politécnico Nacional 2508, Gustavo A. Madero, San Pedro Zacatenco, 07360 Ciudad de México, Mexico. *On the effective topological complexity of Σ_g , $g \geq 2$.*

Let Σ_g be embedded in \mathbb{R}^3 so that Σ_g is invariant under reflections in the xy -, yz -, and xz -planes. Let σ stand for the “antipodal” (orientation-reversing) involution on Σ_g given by $\sigma(x, y, z) = (-x, -y, -z)$. If we consider Σ_g with a left action of $\mathbb{Z}_2 = \langle \sigma \rangle$, then our main result says that there is a monotonic sequence of effective topological complexities for orientable surfaces respect to \mathbb{Z}_2 -action as follows

$$3 \leq \text{TC}^\sigma(\Sigma_2) \leq \text{TC}^\sigma(\Sigma_3) \leq \text{TC}^\sigma(\Sigma_4) \leq \dots \leq 4.$$

(Received August 19, 2019)