

Some 2002–2008
William Lowell Putnam Mathematical Competition
Problems

2A2 Given any five points on a sphere, show that some four of them must lie on a closed hemisphere.

2B2 Consider a polyhedron with at least five faces such that exactly three edges emerge from each of its vertices. Two players play the following game:

Each player, in turn, signs his or her name on a previously unsigned face. The winner is the player who first succeeds in signing three faces that share a common vertex.

Show that the player who signs first will always win by playing as well as possible.

3A1 Let n be a fixed positive integer. How many ways are there to write n as a sum of positive integers, $n = a_1 + a_2 + \cdots + a_k$, with k an arbitrary positive integer and $a_1 \leq a_2 \leq \cdots \leq a_k \leq a_1 + 1$? For example, with $n = 4$ there are four ways: 4, 2+2, 1+1+2, 1+1+1+1.

4A1 Basketball star Shanille O'Keal's team statistician keeps track of the number, $S(N)$, of successful free throws she has made in her first N attempts of the season. Early in the season, $S(N)$ was less than 80% of N , but by the end of the season, $S(N)$ was more than 80% of N . Was there necessarily a moment in between when $S(N)$ was exactly 80% of N ?

4A2 For $i = 1, 2$ let T_i be a triangle with side lengths a_i, b_i, c_i , and area A_i . Suppose that $a_1 \leq a_2, b_1 \leq b_2, c_1 \leq c_2$, and that T_2 is an acute triangle. Does it follow that $A_1 \leq A_2$?

4B2 Let m and n be positive integers. Show that

$$\frac{(m+n)!}{(m+n)^{m+n}} < \frac{m!}{m^m} \frac{n!}{n^n}.$$

7B1 Let f be a polynomial with positive integer coefficients. Prove that if n is a positive integer, then $f(n)$ divides $f(f(n) + 1)$ if and only if $n = 1$. [Editor's note: one must assume f is nonconstant.]

8A2 Alan and Barbara play a game in which they take turns filling entries of an initially empty 2008×2008 array. Alan plays first. At each turn, a player chooses a real number and places it in a vacant entry. The game ends when all the entries are filled. Alan wins if the determinant of the resulting matrix is nonzero; Barbara wins if it is zero. Which player has a winning strategy?

8B1 What is the maximum number of rational points that can lie on a circle in \mathbb{R}^2 whose center is not a rational point? (A *rational point* is a point both of whose coordinates are rational numbers.)