

A NOTE ON THE CONNECTIVITY OF 2-POLYMATROID MINORS

ZACHARY GERSHKOFF AND JAMES OXLEY

ABSTRACT. Brylawski and Seymour independently proved that if M is a connected matroid with a connected minor N , and $e \in E(M) - E(N)$, then $M \setminus e$ or M/e is connected having N as a minor. This paper proves an analogous result for 2-polymatroids. Specifically, if M is a connected 2-polymatroid with a proper connected minor N , then there is an element e of $E(M) - E(N)$ such that $M \setminus e$ or M/e is connected having N as a minor. We also consider what can be said about the uniqueness of the way in which the elements of $E(M) - E(N)$ can be removed so that connectedness is always maintained.

1. INTRODUCTION

Tutte [9] proved that, whenever e is an element of a connected matroid M , at least one of $M \setminus e$ and M/e is connected. Brylawski [1] and Seymour [8] independently extended this theorem by showing that if N is a connected minor of M and e is in $E(M) - E(N)$, then $M \setminus e$ or M/e is connected having N as a minor. In this paper, we prove a similar result for 2-polymatroids.

For a positive integer k , a k -*polymatroid* M is a pair (E, r) consisting of a finite *ground set* E and a *rank function* r , from the power set of E into the integers, satisfying the following conditions:

- (i) $r(\emptyset) = 0$;
- (ii) if $X \subseteq Y \subseteq E$, then $r(X) \leq r(Y)$;
- (iii) if X and Y are subsets of E , then $r(X) + r(Y) \geq r(X \cup Y) + r(X \cap Y)$; and
- (iv) $r(\{e\}) \leq k$ for all $e \in E$.

A matroid is just a 1-polymatroid, so every matroid is a 2-polymatroid. We call M a *polymatroid* if M is a k -polymatroid for some k . Our focus here will be mainly on 2-polymatroids. Elements of a polymatroid of ranks 0, 1, and 2 are called *loops*, *points*, and *lines*, respectively. Non-loop elements p and q are *parallel* if $r(\{p, q\}) = r(\{p\}) = r(\{q\})$.

Many matroid concepts that are stated in terms of the rank function can be extended to polymatroids. In particular, for a polymatroid $M = (E, r)$ and a subset T of E , the *deletion* $M \setminus T$ and the *contraction* M/T of T from M are the polymatroids with ground set $E - T$ and rank functions $r_{M \setminus T}$ and $r_{M/T}$ where $r_{M \setminus T}(X) = r(X)$ and $r_{M/T}(X) = r(X \cup T) - r(T)$ for all subsets X of $E - T$. A *minor* of M is any polymatroid that can be obtained from M by a sequence of deletions and contractions. A polymatroid M is *connected*, or equivalently *2-connected*, if there is no non-empty proper subset X of its ground set E such that

Date: December 3, 2018.

1991 Mathematics Subject Classification. 05B35.

$r(X) + r(E - X) = r(E)$. We sometimes use $E(M)$ and r_M to denote the ground set and rank function of M .

The following is the main result of the paper.

Theorem 1.1. *Let M be a connected 2-polymatroid and let N be a connected minor of M . When $N \neq M$, there is an element e of $E(M) - E(N)$ such that $M \setminus e$ or M/e is connected having N as a minor.*

Unlike in the matroid case, it is not true that, for every element e of $E(M) - E(N)$, at least one of $M \setminus e$ and M/e is connected having N as a minor. For example, let $E(M) = \{x, y, z\}$ where x , y , and z are lines, $r(\{x, y\}) = r(\{y, z\}) = 3$, and $r(\{x, z\}) = 4$. Let N be the 2-polymatroid consisting of a single line z . Then both $M \setminus y$ and M/y are disconnected, as the former consists of two lines in rank 4, and the latter is isomorphic to the matroid $U_{2,2}$.

Theorem 1.1 will be proved in Section 3. The next section includes a number of preliminaries needed for this proof. In Section 4, we consider what can be said about the uniqueness of the element e .

2. PRELIMINARIES

Our matroid terminology follows Oxley [4]. Indeed, much of the notation from matroid theory carries over to polymatroids. For instance, when M is the polymatroid (E, r) and $T \subseteq E$, the deletion $M \setminus (E - T)$ is also denoted by $M|T$. Moreover, we frequently write $r(M)$ for $r(E)$. A subset S of E spans a subset T if $r(S \cup T) = r(S)$. A *component* of M is a maximal non-empty subset X of E such that $M|X$ is connected. As for matroids, the *connectivity function* λ_M or λ of M is defined for all subsets X of $E(M)$ by $\lambda_M(X) = r(X) + r(E - X) - r(E)$. For a positive integer j and a subset Z of $E(M)$, we call Z and $(Z, E(M) - Z)$ *j -separating* if $\lambda_M(Z) < j$.

The *local connectivity* $\square(X, Y)$ between subsets X and Y of E is given by $\square(X, Y) = r(X) + r(Y) - r(X \cup Y)$. Thus $\square(X, E - X) = \lambda(X)$. The following useful results for local connectivity and connectivity are proved for matroids in [4, Lemma 8.2.3]; the proofs there extend to polymatroids.

Lemma 2.1. *Let (E, r) be a polymatroid and let X_1, X_2, Y_1 , and Y_2 be subsets of E with $Y_1 \subseteq X_1$ and $Y_2 \subseteq X_2$. Then*

$$\square(Y_1, Y_2) \leq \square(X_1, X_2).$$

Lemma 2.2. *Let (E, r) be a polymatroid M and let X, C , and D be disjoint subsets of E . Then*

$$\lambda_{M \setminus D/C}(X) \leq \lambda_M(X).$$

Moreover, equality holds if and only if

$$r(X \cup C) = r(X) + r(C)$$

and

$$r(E - X) + r(E - D) = r(E) + r(E - (X \cup D)).$$

Next we note a useful consequence of Lemma 2.1.

Corollary 2.3. *Let X and Y be sets in a polymatroid M such that $X \cap Y \neq \emptyset$ and both $M|X$ and $M|Y$ are connected. Then $M|(X \cup Y)$ is connected.*

Proof. Suppose that $M|(X \cup Y)$ is disconnected letting Z be a component of it. Let $W = (X \cup Y) - Z$. By Lemma 2.1, $\cap(Z \cap X, W \cap X) \leq \cap(Z, W) = 0$. As $M|X$ is connected, $Z \cap X$ or $W \cap X$ is empty. By symmetry, $Z \cap Y$ or $W \cap Y$ is empty. As neither Z nor W is empty, we may assume that both $Z \cap X$ and $W \cap Y$ are empty. It follows that $X \cap Y$ is empty; a contradiction. \square

The following generalization of a matroid result was noted in [7, Lemma 3.12(ii)].

Lemma 2.4. *Let A, B , and C be subsets of the ground set of a polymatroid. Then*

$$\cap(A \cup B, C) + \cap(A, B) = \cap(A \cup C, B) + \cap(A, C).$$

We omit the proof of the next result, which follows easily from Lemma 2.2.

Lemma 2.5. *If Z is a component of a minor of a polymatroid M , then Z is contained in a component of M , and $M \setminus Z = M/Z$.*

As noted in [4, p.409], with every 2-polymatroid M , we can associate a matroid as follows. Let L be the set of lines of M . For each ℓ in L , freely add two points to ℓ letting M^+ be the resulting 2-polymatroid. Then M' , the *natural matroid derived from M* , is $M^+ \setminus L$. Oxley, Semple, and Whittle [7, Lemma 3.3] noted the following straightforward result.

Lemma 2.6. *Let M be a 2-polymatroid with $|E(M)| \geq 2$ and let M' be the natural matroid derived from M . Then M is connected if and only if M' is connected.*

The proof of our main theorem will use the operations of parallel connection and 2-sum of polymatroids as introduced by Matúš [3] and Hall [2]. For a positive integer k , let M_1 and M_2 be k -polymatroids (E_1, r_1) and (E_2, r_2) . Suppose first that $E_1 \cap E_2 = \emptyset$. The *direct sum* $M_1 \oplus M_2$ of M_1 and M_2 is the k -polymatroid $(E_1 \cup E_2, r)$ where, for all subsets A of $E_1 \cup E_2$, we have $r(A) = r(A \cap E_1) + r(A \cap E_2)$. Clearly a 2-polymatroid is connected if and only if it cannot be written as the direct sum of two non-empty 2-polymatroids. Now suppose that $E_1 \cap E_2 = \{p\}$ and $r_1(\{p\}) = r_2(\{p\})$. Let $P(M_1, M_2)$ be $(E_1 \cup E_2, r)$ where r is defined for all subsets A of $E_1 \cup E_2$ by

$$r(A) = \min\{r_1(A \cap E_1) + r_2(A \cap E_2), r_1((A \cap E_1) \cup p) + r_2((A \cap E_2) \cup p) - r_1(\{p\})\}.$$

Hall [2] notes that it is routine to check that $P(M_1, M_2)$ is a k -polymatroid. We call it the *parallel connection* of M_1 and M_2 with respect to the *basepoint* p . When M_1 and M_2 are both matroids, this definition coincides with the usual definition of the parallel connection of matroids.

Now let M_1 and M_2 be 2-polymatroids having at least two elements. Suppose that $E(M_1) \cap E(M_2) = \{p\}$, that neither $\lambda_{M_1}(\{p\})$ nor $\lambda_{M_2}(\{p\})$ is 0, and that $r_1(\{p\}) = r_2(\{p\}) = 1$. We define the *2-sum*, $M_1 \oplus_2 M_2$, of M_1 and M_2 to be $P(M_1, M_2) \setminus p$. This definition [7] extends Hall's definition since the latter requires each of M_1 and M_2 to have at least three elements. Weakening that requirement does not alter the validity of Hall's proof of the following result [2, Proposition 3.6].

Proposition 2.7. *Let M be a 2-polymatroid (E, r) having a partition (X_1, X_2) of E such that $r(X_1) + r(X_2) = r(E) + 1$. Then there are 2-polymatroids M_1 and M_2 with ground sets $X_1 \cup p$ and $X_2 \cup p$, where p is a new element not in E , such that $M = P(M_1, M_2) \setminus p$. In particular, for all $A \subseteq X_1 \cup p$,*

$$r_1(A) = \begin{cases} r(A), & \text{if } p \notin A; \\ r((A - p) \cup X_2) - r(X_2) + 1, & \text{if } p \in A. \end{cases}$$

The following was shown by Hall [2, Corollary 3.5].

Proposition 2.8. *Let M_1 and M_2 be 2-polymatroids (E_1, r_1) and (E_2, r_2) where $E_1 \cap E_2 = \{p\}$. Suppose $r_1(\{p\}) = r_2(\{p\}) = 1$ and each of M_1 and M_2 has at least two elements. Then the following are equivalent.*

- (i) M_1 and M_2 are both 2-connected;
- (ii) $M_1 \oplus_2 M_2$ is 2-connected; and
- (iii) $P(M_1, M_2)$ is 2-connected.

The next theorem, a special case of a result of Hall [2, Theorem 4.3], will play a crucial role in the proof of our main theorem.

Theorem 2.9. *Every connected 2-polymatroid M having at least two elements has distinct elements x and y such that each of $\{M \setminus x, M/x\}$ and $\{M \setminus y, M/y\}$ contains a connected 2-polymatroid.*

The following lemma holds for polymatroids in general and will be useful in Section 4.

Lemma 2.10. *Let M be a connected polymatroid and let M/e be disconnected. If Z is a component of M/e , then $\square(Z, \{e\}) > 0$.*

Proof. Let $Y = E(M) - (Z \cup e)$. Then

$$(2.1) \quad r_{M/e}(Z) + r_{M/e}(Y) = r(M/e) = r(M) - r_M(\{e\}).$$

Moreover, since M is connected,

$$(2.2) \quad r_M(Z) + r_M(Y \cup e) > r(M).$$

By the definition of local connectivity, $r_M(Y) + r_M(\{e\}) - \square(Y, \{e\}) = r(Y \cup e)$, so we can rewrite (2.2) as

$$(2.3) \quad r_M(Z) + r_M(Y) > r(M) - r_M(\{e\}) + \square(Y, \{e\}).$$

By subtracting (2.1) from (2.3), we obtain

$$(2.4) \quad (r_M(Z) - r_{M/e}(Z)) + (r_M(Y) - r_{M/e}(Y)) > \square(Y, \{e\}).$$

The differences on the left-hand side can be rewritten as local connectivities. Thus $\square(Z, \{e\}) + \square(Y, \{e\}) > \square(Y, \{e\})$, so $\square(Z, \{e\}) > 0$. \square

3. A SPLITTER THEOREM FOR CONNECTED 2-POLYMATROIDS

This section is devoted to proving the main result of the paper.

Proof of Theorem 1.1. Assume that the theorem fails. Then it follows from Theorem 2.9 that N is non-empty. Hence, as M is connected, it has no loops. Next we note the following.

3.1.1. *If x is an element of M such that both $M \setminus x$ and M/x have N as a minor, then x is a line of M .*

Clearly neither $M \setminus x$ nor M/x is connected. It follows, by a result of Oxley and Whittle [5, Theorem 3.1], that $r_M(\{x\}) \neq 1$. Hence x is a line of M .

Take e in $E(M) - E(N)$. Then, for some M_0 in $\{M \setminus e, M/e\}$, the 2-polymatroid M_0 has N as a minor. By assumption, M_0 is not connected. Take an element f of a component of M_0 that avoids $E(N)$. Then both $M \setminus f$ and M/f have N as a minor. Thus, by 3.1.1, f is a line of M . Moreover,

3.1.2. $r(M \setminus f) = r(M)$.

To see this, suppose $r(E - f) < r(E)$. Then, as M is connected, $r(E - f) = r(E) - 1$. Let M' be the natural matroid derived from M and let f_1 and f_2 be the points of M' corresponding to f . Then $M' \setminus f_1$ has f_2 as a coloop, so M' / f_1 is connected. Now M / f is disconnected, so, by Lemma 2.6, $M' / f_1, f_2$ is disconnected. Therefore, $M' / f_1 \setminus f_2$ is connected. But $M' \setminus f_2$ has f_1 as a coloop, so $M' \setminus f_2 / f_1 = M' \setminus f_2 \setminus f_1$. As this matroid is connected, by Lemma 2.6, $M \setminus f$ is too; a contradiction.

3.1.3. *Let K be a component of M / f . Then $M|(K \cup f)$ is connected.*

Suppose $M|(K \cup f)$ is disconnected. Then K is the disjoint union of sets X and Y such that $r(X \cup f) + r(Y) = r(K \cup f)$. As Y is 1-separating in $M|(K \cup f)$, it is 1-separating in $(M|(K \cup f)) / f$, that is, in $(M / f)|K$. But K is a component of the last matroid, so $K = Y$. Thus $X = \emptyset$, so $r(K \cup f) = r(K) + r(\{f\})$. It follows that K is 1-separating in M ; a contradiction. Hence 3.1.3 holds.

Now let F be a component of M / f that avoids $E(N)$. By 3.1.1, every element of F is a line in M . Let $G = E(M) - f - F$. By 3.1.3, $M|(F \cup f)$ is connected. Next we show the following.

3.1.4. *There is a line g in F such that $(M|(F \cup f)) \setminus g$ or $(M|(F \cup f)) / g$ is connected. Moreover, $\Pi(\{f\}, \{g\}) < 2$.*

By Theorem 2.9, F contains an element g such that $(M|(F \cup f)) \setminus g$ or $(M|(F \cup f)) / g$ is connected. As g is in F , we see that g is a line. Since the theorem fails, $M \setminus g$ is not connected, so $\Pi(\{f\}, \{g\}) < 2$. Thus 3.1.4 holds.

3.1.5. *$M|(G \cup f)$ is connected; $(M|(F \cup f)) / g$ is connected; and $\Pi(\{f\}, \{g\}) = 1$.*

To see this, first note that, by 3.1.3, $M|(K \cup f)$ is connected for each component K of M / f . Then, by Corollary 2.3, $M|(G \cup f)$ is connected. The same argument shows that $(M|(F \cup f)) \setminus g$ is disconnected for if it is connected, then so is $M \setminus g$; a contradiction. Thus, by 3.1.4, $(M|(F \cup f)) / g$ is connected.

Now suppose that $\Pi_M(\{f\}, \{g\}) = 0$. Then, as $\Pi_{M/f}(G, \{g\}) = 0$, one easily checks that $\Pi_M(G \cup f, \{g\}) = 0$. Hence $M|(G \cup f) = (M|(G \cup f \cup g)) / g = (M/g)|(G \cup f)$. Thus, as $(M|(F \cup f)) / g$ and $M|(G \cup f)$ are connected and both contain f , Corollary 2.3 implies that M/g is connected; a contradiction. Hence 3.1.5 holds.

Recall that f is a line of M such that $M \setminus f$ and M / f are disconnected. Moreover, F is a component of M / f and $E(N) \subseteq G = E(M) - f - F$. Let A be a component of $M \setminus f$ avoiding $E(N)$ and let $B = E(M) - f - A$. The next two observations follow because M is connected.

3.1.6. *Neither A nor B spans f .*

3.1.7. *$r(G \cup f) < r(G) + 2$ and $r(F \cup f) < r(F) + 2$.*

Next we show the following.

3.1.8. *At least one of $A \cap G$, $A \cap F$, $B \cap F$, and $B \cap G$ is empty.*

Suppose that all four intersections are non-empty. By 3.1.2, $r(E - f) = r(E)$. Thus $r(A) + r(B) = r(E)$ and $r(F \cup f) + r(G \cup f) = r(E) + 2$. Adding these two equations and applying submodularity to the left-hand side gives

$$r(A \cup F \cup f) + r(A \cap F) + r(B \cup G \cup f) + r(B \cap G) \leq 2r(E) + 2,$$

so

$$(3.1) \quad [r(A \cup F \cup f) + r(B \cap G)] + [r(B \cup G \cup f) + r(A \cap F)] \leq 2r(E) + 2.$$

As $(A \cup F \cup f, B \cap G)$ and $(B \cup G \cup f, A \cap F)$ are partitions of $E(M)$, we deduce, since M is connected, that equality holds in (3.1). Hence the two specified partitions are 2-separating in M . By symmetry, so are $(A \cup G \cup f, B \cap F)$ and $(B \cup F \cup f, A \cap G)$. By Propositions 2.7 and 2.8, M can be written as the 2-sum with basepoint p_{AF} of two connected 2-polymatroids, one with ground set $(A \cap F) \cup p_{AF}$ and the other, Q_0 , with ground set $(E(M) - (A \cap F)) \cup p_{AF}$. By arguing in terms of the natural matroid derived from M , it is straightforward to check that, in Q_0 , each of $A \cap G$, $B \cap F$, and $B \cap G$ is 2-separating. Hence we can decompose Q_0 as a 2-sum of two connected 2-polymatroids one with ground set $(A \cap G) \cup p_{AG}$. Repeating this process twice more, we obtain a connected 2-polymatroid Q with ground set $\{f, p_{AF}, p_{AG}, p_{BF}, p_{BG}\}$ where M is obtained from Q by attaching, via 2-sums, connected 2-polymatroids with ground sets $(A \cap F) \cup p_{AF}$, $(A \cap G) \cup p_{AG}$, $(B \cap F) \cup p_{BF}$, and $(B \cap G) \cup p_{BG}$.

As $M|A$ is connected, Proposition 2.8 implies that p_{AG} and p_{AF} are parallel in Q . Since $(M/f)|F$ is connected, p_{BF} and p_{AF} are parallel in Q/f . But p_{AG} and p_{AF} are also parallel in Q/f unless they are loops. In the exceptional case, $A \cap F$ contains a component of M/f ; a contradiction. We deduce that the component of M/f containing F also contains $A \cap G$; a contradiction. Thus 3.1.8 holds.

By 3.1.8, A or B is contained in F or G , and F or G is contained in A or B . We know that $B \cap G$ is non-empty because it contains $E(N)$.

Suppose both F and G span f . Then A or B spans f ; a contradiction to 3.1.6. By 3.1.7, there are two remaining cases to consider:

- (i) $r(F \cup f) = r(F) + 1$; and
- (ii) $r(F \cup f) = r(F)$ and $r(G \cup f) = r(G) + 1$.

By 3.1.5, $(M|(F \cup f))/g$ is connected and $\square(\{f\}, \{g\}) = 1$. Thus $r(\{f, g\}) = 3$. Assume (i) holds. Then $\square_{M/g}(F - g, \{f\}) = r(F) + r(\{f, g\}) - r(F \cup f) - r(\{g\}) = 0$. Thus $\{f\}$ is a component of $(M|(F \cup f))/g$. As the last matroid is connected, we deduce that $F = \{g\}$. Thus $M \setminus g = M|(G \cup f)$ so, by 3.1.5, $M \setminus g$ is connected; a contradiction.

We now know that (ii) holds. As neither A nor B spans f , neither has F as a subset. Thus both $A \cap F$ and $B \cap F$ are non-empty. As $B \cap G$ is non-empty, 3.1.8 implies that $A \cap G$ is empty. Then $G \subseteq B$. But $r(G \cup f) = r(G) + 1$. Therefore $r(B \cup f) \leq r(B) + 1$. Since $r(B \cup f) \neq r(B)$, it follows that $r(B \cup f) = r(B) + 1$.

We have $r(A) + r(B) = r(M \setminus f)$, and, by 3.1.2, $r(M \setminus f) = r(M)$. As $r(B \cup f) = r(B) + 1$, we deduce that $r(A) + r(B \cup f) = r(M) + 1$, so M can be written as a 2-sum with basepoint p of two connected 2-polymatroids with ground sets $A \cup p$ and $B \cup f \cup p$. Let the former be M_1 .

Suppose M_1 has at least three elements. Then, by Theorem 2.9, M_1 has an element q such that $q \neq p$ and $M_1 \setminus q$ or M_1/q is connected. Observe that if M_1/q is connected, then $\square(\{q\}, \{p\}) = 0$ otherwise p is a loop in the contraction; a contradiction. It follows, by [7, Lemma 4.3], that $M \setminus q$ or M/q is the 2-sum of two connected 2-polymatroids each with at least two elements. Thus, by Proposition 2.8, $M \setminus q$ or M/q is connected. As q is in A and hence in F , both $M \setminus q$ and M/q have N as a minor and so we obtain a contradiction.

We may now assume that M_1 consists of a single line a through p .

3.1.9. M/a is connected.

Assume M/a is disconnected. Then its ground set has a partition (V, W) such that $r_{M/a}(V) + r_{M/a}(W) = r(M/a)$. Now we may assume that f is in V . Thus $W \subseteq B$ since $A = \{a\}$. As $\sqcap_M(A, B) = 0$, it follows that $r_{M/a}(W) = r_M(W)$. Hence $r_M(V \cup a) + r_M(W) = r(M)$; a contradiction. We conclude that 3.1.9 holds.

As $a \in F$, we know that M/a has N as a minor. Thus we have a contradiction that completes the proof of the theorem. \square

The argument above relies heavily on the fact that we have a 2-polymatroid. However, we believe that the main theorem also holds for k -polymatroids for all $k > 2$.

Conjecture 3.2. *Let M be a connected k -polymatroid and let N be a connected minor of M . When $N \neq M$, there is an element e of $E(M) - E(N)$ such that $M \setminus e$ or M/e is connected having N as a minor.*

4. UNIQUENESS

By Theorem 1.1, for every connected 2-polymatroid M and every connected proper minor N of M , we can remove the elements of $E(M) - E(N)$ one at a time so that we stay connected and maintain N as a minor. In this section, we consider what can be said about the uniqueness of this sequence of element removals

Now let M be a connected polymatroid and N be a connected proper minor of M . An *admissible ordering* of $E(M) - E(N)$ is an ordering (a_1, a_2, \dots, a_n) of the set $E(M) - E(N)$ such that, for each k in $\{1, 2, \dots, n\}$, there is a connected minor M_k of M with ground set $E(M) - \{a_1, a_2, \dots, a_k\}$ such that M_k is a minor of M_{k-1} , where $(M_0, M_n) = (M, N)$. We give an example below to show that an admissible ordering may be unique. We shall show, however, that we always retain some flexibility with respect to the way in which the elements are removed unless $|E(M) - E(N)| = 1$. Formally, a *constrained admissible ordering* is an ordering $((\alpha_1, a_1), (\alpha_2, a_2), \dots, (\alpha_n, a_n))$ such that $E(M) - E(N) = \{a_1, a_2, \dots, a_n\}$ where each α_i is a deletion or contraction operation, and, for each k in $\{1, 2, \dots, n\}$, there is a connected minor M_k of M with ground set $E(M) - \{a_1, a_2, \dots, a_k\}$ where M_k is obtained from M_{k-1} by removing a_k by the operation designated by α_k , and $(M_0, M_n) = (M, N)$.

To construct a 2-polymatroid with a unique admissible ordering, let N be a simple non-empty connected matroid. Take $N \oplus U_{n,n}$ where the ground set of $U_{n,n}$ is $\{b_0, b_1, \dots, b_{n-1}\}$. Take $b_n \in E(N)$ and consider the 2-polymatroid M whose ground set is $E(N) \cup \{f_i : 1 \leq i \leq n\}$ where $f_i = \{b_{i-1}, b_i\}$ for all i , and the rank function of M is induced by that of $N \oplus U_{n,n}$. Then M is connected, $M \setminus f_1, f_2, \dots, f_k$ is connected for all k in $\{1, 2, \dots, n\}$, and $M \setminus f_1, f_2, \dots, f_n = N$. Thus (f_1, f_2, \dots, f_n) is an admissible ordering of $E(M) - E(N)$. It is not difficult to check that the admissible ordering is unique. Note, however, that $M \setminus f_1 \setminus f_2 = M/f_1 \setminus f_2$, so this example does not give us a unique constrained admissible ordering. Indeed, as the next result shows, except in the trivial case, there can never be such a unique ordering.

Theorem 4.1. *Let M be a connected 2-polymatroid and N be a connected proper minor of M . Then there is a unique constrained admissible ordering of $E(M) - E(N)$ if and only if $|E(M) - E(N)| = 1$.*

The next two lemmas contain the core of the proof of this theorem.

Lemma 4.2. *Let each of \dagger and \ddagger denote a deletion or contraction operation. Suppose both $M \dagger e$ and $M \dagger e \ddagger f$ are connected, but $M \ddagger f$ is not. Then $\{e\}$ and Z are the components of $M \ddagger f$. Moreover,*

$$M \ddagger f \setminus e = M \ddagger f / e.$$

Proof. Let (X, Y) be a 1-separating partition of $E(M \ddagger f)$ with Y minimal and non-empty avoiding e . Then $\lambda_{M \ddagger f}(Y) = 0$. Thus, by Lemma 2.2, $\lambda_{M \ddagger f \dagger e}(Y) = 0$. As $M \dagger e \ddagger f$ is connected, $X - e = \emptyset$. Thus $\{e\}$ and Z are components of $M \ddagger f$. Hence, by Lemma 2.5, $M \ddagger f \setminus e = M \ddagger f / e$. \square

Lemma 4.3. *Let M be a connected polymatroid and N be a connected minor of M . Let e and f be distinct elements of $E(M) - \{e, f\}$ and let $Z = E(M) - \{e, f\}$. Suppose that $\{\alpha, \beta\} = \{\setminus, /\} = \{\gamma, \delta\}$. Assume that $M\alpha e$ and $M\alpha e \gamma f$ are connected. Then*

- (i) $M\gamma f$ is connected and $M\alpha e \gamma f = M\gamma f \alpha e$; or
- (ii) $M\gamma f$ is disconnected, $M\alpha e \gamma f = M\beta e \gamma f$, and
 - (a) $M\beta e$ is connected; or
 - (b) $M\beta e$ is disconnected, $M\alpha e \gamma f = M\delta f \beta e$, and $M\delta f$ is connected; or
 - (c) $M\beta e$ and $M\delta f$ are disconnected, and $M\alpha e \gamma f = M\alpha e \delta f$.

Proof. We may assume that $M\gamma f$ is disconnected otherwise (i) holds. As $M\alpha e \gamma f$ is connected, Lemma 4.2 implies that $\{e\}$ is a component of $M\gamma f$, and $M\gamma f \setminus e = M\gamma f / e$. Thus

$$M\alpha e \gamma f = M\gamma f \alpha e = M\gamma f \beta e = M\beta e \gamma f.$$

We may assume that $M\beta e$ is disconnected otherwise (ii)(a) holds. Then $M\beta e$ has $\{f\}$ as a component and $M\beta e \setminus f = M\beta e / f$. Thus

$$M\alpha e \gamma f = M\beta e \gamma f = M\beta e \delta f = M\delta f \beta e.$$

We may now assume that $M\delta f$ is disconnected otherwise (ii)(b) holds. Then $M\delta f \setminus e = M\delta f / e$. Thus $M\delta f \beta e = M\delta f \alpha e$. Hence $M\alpha e \gamma f = M\delta f \beta e = M\delta f \alpha e = M\alpha e \delta f$, and (ii)(c) holds. \square

We are now able to prove the main result of this section.

Proof of Theorem 4.1. We may assume that $|E(M) - E(N)| \geq 2$. Let $((\alpha, e), (\gamma, f), (\alpha_3, a_3) \dots, (\alpha_k, a_k))$ be a constrained admissible ordering of $E(M) - E(N)$. We use Lemma 4.3 to show that $E(M) - E(N)$ has a constrained admissible ordering $((\alpha_1, a_1), (\alpha_2, a_2), (\alpha_3, a_3), \dots, (\alpha_k, a_k))$ in which $((\alpha, e), (\gamma, f)) \neq ((\alpha_1, a_1), (\alpha_2, a_2))$ and $\{e, f\} = \{a_1, a_2\}$. If $M\gamma f$ is connected, then we can take $((\alpha_1, a_1), (\alpha_2, a_2))$ to be $((\gamma, f), (\alpha, e))$. Using the notation of Lemma 4.3, if $M\gamma f$ is disconnected but $M\beta e$ is connected, then we can take $((\alpha_1, a_1), (\alpha_2, a_2))$ to be $((\beta, e), (\gamma, f))$. Now suppose that $M\gamma f$ and $M\beta e$ are disconnected. If $M\delta f$ is connected, then we can take $((\alpha_1, a_1), (\alpha_2, a_2))$ to be $((\delta, f), (\beta, e))$. Finally, if $M\delta f$ is disconnected, then we can take $((\alpha_1, a_1), (\alpha_2, a_2))$ to be $((\alpha, e), (\delta, f))$. \square

REFERENCES

- [1] T. H. Brylawski, A decomposition for combinatorial geometries, *Trans. Amer. Math. Soc.* **171** (1972), 235–282.
- [2] D. Hall, Essential elements in connected k -polymatroids, *Adv. in Appl. Math.* **50** (2013), 281–291.
- [3] F. Matuš, Adhesivity of polymatroids, *Discrete Math.* **307** (2007) 2464–2477.

- [4] J. Oxley, *Matroid Theory*, Second edition, Oxford University Press, New York, 2011.
- [5] J. Oxley and G. Whittle, Connectivity of submodular functions, *Discrete Math.* **105**(1992), 173–184.
- [6] J. Oxley, C. Semple, and G. Whittle, The structure of the 3-separations of 3-connected matroids, *J. Combin. Theory Ser. B* **92** (2004), 257–293.
- [7] J. Oxley, C. Semple, and G. Whittle, A splitter theorem for 3-connected 2-polymatroids, submitted.
- [8] P. D. Seymour, A note on the production of matroid minors, *J. Combin. Theory Ser. B* **22** (1977), 289–295.
- [9] W. T. Tutte, Connectivity in matroids, *Canad. J. Math.* **18** (1966), 1301–1324.

MATHEMATICS DEPARTMENT, LOUISIANA STATE UNIVERSITY, BATON ROUGE, LOUISIANA
E-mail address: zgersh2@math.lsu.edu

MATHEMATICS DEPARTMENT, LOUISIANA STATE UNIVERSITY, BATON ROUGE, LOUISIANA
E-mail address: oxley@math.lsu.edu