

SUPER-EXPONENTIAL 2-DIMENSIONAL DEHN FUNCTIONS

JOSH BARNARD, NOEL BRADY, AND PALLAVI DANI

ABSTRACT. We produce examples of groups of type \mathcal{F}_3 with 2-dimensional Dehn functions of the form $\exp^n(x)$ (a tower of exponentials of height n), where n is any natural number.

CONTENTS

1. Introduction	2
2. Preliminaries	6
3. Overview and geometric intuition	8
3.1. A 1-dimensional Dehn function example and schematic diagrams	8
3.2. The 2-dimensional Dehn function schematic procedure	11
4. The groups	16
4.1. Inductive definition of the groups G_n and H_n	18
4.2. Further details about Table 1.	19
4.3. The cell structure in K_{G_n} and K_{H_n}	23
5. Lower bounds	24
5.1. Some spheres	24
5.2. Iteration notation	27
5.3. The van Kampen diagrams for moves	29
5.4. Geometry of the moves	32
5.5. Lower bound estimates	36
5.6. Induction steps	39
6. Upper bounds for graphs of groups	42
6.1. The 3-complex \widetilde{K}_G	43
6.2. Transverse maps	44

Date: October 13, 2009.

¹N. Brady was partially supported by NSF grant no. DMS-0505707

6.3.	General strategy for the proofs	46
6.4.	The admissible filling $\bar{\tau}$	48
6.5.	Filling volumes in graphs of groups	49
6.6.	Upper bounds for the super-exponential examples	51
7.	Proof of the area-distortion lemma	52
7.1.	Strategy	53
7.2.	Geometry of \mathbf{u} -corridors	53
7.3.	Construction of Δ	56
7.4.	Upper bound for $\text{Area}(\Delta)$	56
	References	61

1. INTRODUCTION

Dehn functions have a long and rich history in group theory and topology. The germ of the notion of Dehn function was expounded by Max Dehn in his solution to the word problem for the fundamental groups of closed hyperbolic surfaces in [10, 11]. Gromov [13, 14] further developed the notion of Dehn function as a filling invariant for a finitely presented group, and proposed the investigation of higher dimensional filling invariants.

A function $\delta : \mathbb{N} \rightarrow \mathbb{N}$ is called the (1-dimensional) Dehn function of a finite presentation if it is the minimal function with the following property. Every word of length at most x in the generators representing the identity element of the group can be expressed as a product of at most $\delta(x)$ conjugates of relators and their inverses. Stated more geometrically, every loop of combinatorial length at most x in the universal cover of a presentation 2-complex for the group, can be null-homotoped using at most $\delta(x)$ 2-cells. The adjective 1-dimensional refers to the fact that the function $\delta(x)$ measures the area of efficient disk fillings of 1-dimensional spheres. It is customary to drop the adjective 1-dimensional, and to simply talk about Dehn functions of finite presentations. It is known that, up to coarse Lipschitz equivalence, the Dehn function is a well-defined geometric invariant of a finitely presented group.

Dehn functions are intimately connected to the solution of the word problem in finitely presented groups. For example, a group has a solvable word problem if and only if its Dehn function is bounded by a recursive function. In particular, the existence of groups with unsolvable word problem implies that there are groups with Dehn function not bounded above by any recursive function. We now have a greater understanding, thanks to the intense research activity of the past two decades and in particular to the deep work of [16], of which functions can be Dehn functions of finitely presented groups. For example, combining the results of [5] and [16], we know that the set of exponents α for which x^α is coarse Lipschitz equivalent to a Dehn function is dense in $\{1\} \cup [2, \infty)$.

Following Gromov, for each integer $k \geq 1$ one can define k -dimensional Dehn functions for groups G of type \mathcal{F}_{k+1} (that is, G admits a $K(G, 1)$ with finite $(k+1)$ -skeleton). Roughly speaking, a k -dimensional Dehn function, $\delta_G^{(k)}(x)$, gives a minimal upper bound for the number of $(k+1)$ -cells needed in a $(k+1)$ -ball filling of a singular k -sphere in X of k -volume at most x . As in the case of ordinary Dehn functions, the coarse Lipschitz equivalence class of a k -dimensional Dehn function is an invariant of a group of type \mathcal{F}_{k+1} . The k -dimensional Dehn functions are examples of Gromov's higher dimensional filling invariants of groups. Precise definitions of the $\delta_G^{(k)}(x)$ are given in Section 2.

In recent years, there has been considerable progress in our understanding of which functions are (k -dimensional) Dehn functions of groups [1, 2, 4, 6, 8, 17]. Combining the results of [4, 6], one now knows that for each $k \geq 2$, the set of exponents α for which x^α is coarse Lipschitz equivalent to a k -dimensional Dehn function is dense in $[1, \infty)$. Young [18] has shown that for $k \geq 3$ there are groups with k -dimensional Dehn function not bounded above by any recursive function. What Young actually shows is that for $k \geq 2$ there exist groups for which the Dehn functions, defined in terms of admissible $(k+1)$ -dimensional manifold fillings of admissible k -tori, are not bounded above by any recursive function. For $k \geq 3$, he then uses the trick introduced in Remark 2.6(4) of [4] to conclude that the Dehn

functions $\delta^{(k)}$ of these groups have the same property. In contrast, Papasoglu [15] shows that 2-dimensional Dehn functions are all bounded above by recursive functions. In the case $k = 2$, Young's example together with Papasoglu's result show that the Dehn function based on admissible 3-manifold fillings of 2-tori is different from the Dehn function $\delta^{(2)}$ introduced above. In summary, Dehn functions based on admissible $(k + 1)$ -ball fillings of admissible k -spheres do not have to be bounded above by recursive functions for $k \neq 2$. The case $k = 2$ is indeed special.

Papasoglu asked if there were examples of groups with super-exponential 2-dimensional Dehn functions. See also the remarks by Gromov in 5.D.(6) on page 100 of [14]. Pride and Wang [17] produced an example of a group whose 2-dimensional Dehn function lies between $e^{\sqrt{x}}$ and e^x . The purpose of the current paper is to describe two families of groups $\{G_n\}_{n=2}^\infty$ and $\{H_n\}_{n=2}^\infty$ whose 2-dimensional Dehn functions display a range of super-exponential behavior.

Theorem 1.1. *For every $n > 0$, there exist groups H_n and G_n of type \mathcal{F}_3 , with $\delta_{H_n}^{(2)}(x) \simeq \exp^n(\sqrt{x})$ and $\delta_{G_n}^{(2)}(x) \simeq \exp^n(x)$.*

Here \exp^n denotes the n -fold composition of exponential functions.

Remark 1.2. The combination-subdivision techniques of the current paper are one way of producing groups with super-exponential 2-dimensional Dehn functions. There are other strategies that have the potential to produce groups of type \mathcal{F}_3 whose 2-dimensional Dehn functions have super-exponential behavior. The following was suggested to the second author by Steve Pride.

Take a group A of type \mathcal{F}_3 with polynomially bounded ordinary Dehn function and containing a finitely presented subgroup B whose ordinary Dehn function is some tower of exponentials. Now let G be the HNN extension with base group A , edge group B , and both edge maps are just the inclusion $B \rightarrow A$. Because of the finiteness properties of A and B , we know that G is of type \mathcal{F}_3 . Finally take a word in B that has very large area filling in B , and efficient filling in A . The product of this word with the HNN stable letter gives an annulus, whose ends can be capped off using the efficient A -fillings. This gives

a 2-sphere in the HNN space with small area but large 3-dimensional filling.

So the whole problem reduces to finding groups with subgroups whose area is highly distorted. It is tempting to use the remarkable constructions of Birget-Ol'shanskii-Rips-Sapir [3] to this end. However, it is likely that the ambient groups in [3] are not of type \mathcal{F}_3 . In particular, using the notation of [3], start with the Baumslag-Solitar group $BS(1, 2) = \langle a, t \mid tat^{-1} = a^2 \rangle$ as the group G_b , and then construct the group $H_N(S)$. The Cayley graph of G_b is contained in that of $H_N(S)$. Consider the loop $[a, t^n at^{-n}]$, which is filled by a disk of the form shown in Figure 6 of [3]. (This figure is viewed as a punctured sphere diagram, with the small loop labeled u_b as the boundary, where the disk fillings track the behavior of the S -machine that reduces $[a, t^n at^{-n}]$ to the identity.) This disk filling has polynomial area. On the other hand, this loop is a product of exponentially many loops of the form $tat^{-1}a^{-2}$, each of which admits a disk filling of the form in Figure 6. These two methods of filling the word $[a, t^n at^{-n}]$ produce a 2-sphere in the Cayley complex of $H_N(S)$, indicating the possibility that $H_N(S)$ is not of type \mathcal{F}_3 . Indeed, it is likely that the group $G_N(S)$ of [3] is not of type \mathcal{F}_3 either.

This paper is organized as follows. In Section 2 we give the definitions of k -dimensional Dehn functions, and remind the reader of a few standard techniques that are useful for establishing lower bounds. Section 3 gives a geometric overview of the groups, and the inductive construction of the ball-sphere pairs that are used to establish the lower bounds for the Dehn functions. Since the precise definitions of the groups are so involved, it is important that the reader keep the overview in mind when reading the later sections. In Section 4 we give the precise definitions of the groups H_n and G_n . These are defined recursively, and the details are recorded in Table 1 for the reader's convenience.

The proof of Theorem 1.1 is broken into two parts. We establish the appropriate lower bounds in Section 5 by exhibiting sequences of embedded ball-sphere pairs in the universal covers of 3-dimensional $K(G, 1)$ -complexes. In Section 6 we obtain upper bounds on 2-dimensional Dehn functions of general graphs of groups in terms of

information about their vertex and edge groups. These are then used to prove the desired upper bounds for $\delta_{H_n}^{(2)}$ and $\delta_{G_n}^{(2)}$. This proof depends on a key *area distortion* result, which is established in Section 7.

We thank the referee for helpful comments and suggestions.

2. PRELIMINARIES

In this section we define k -dimensional Dehn functions, and describe two useful techniques for establishing lower bounds for Dehn functions. As in the case of ordinary (or 1-dimensional) Dehn functions, one must work with coarse Lipschitz classes of functions in order to obtain a geometric invariant of a group.

We begin with a discussion of coarse Lipschitz equivalence. Given two functions $f, g: [0, \infty) \rightarrow [0, \infty)$ we define $f \preceq g$ to mean that there exists a positive constant C such that

$$f(x) \leq C g(Cx) + Cx$$

for all $x \geq 0$. If $f \preceq g$ and $g \preceq f$ then f and g are said to be *coarse Lipschitz equivalent* (or simply *equivalent*), denoted $f \simeq g$.

We work with the same definition of high-dimensional Dehn functions as in [4], which is equivalent to the notions in [8] and in [2]. In order to give a formal definition of $\delta^{(k)}(x)$, one needs to work with a suitable class of maps that facilitates the counting of cells. In [4] the class of *admissible maps* is used, and we use this class here.

If W is a compact k -dimensional manifold and X a CW complex, an *admissible map* is a continuous map $f: W \rightarrow X^{(k)} \subset X$ such that $f^{-1}(X^{(k)} - X^{(k-1)})$ is a disjoint union of open k -dimensional balls, each mapped by f homeomorphically onto a k -cell of X . The key fact about admissible maps is that every continuous map is homotopic to an admissible one.

Lemma 2.1 (Lemma 2.3 of [4]). *Let W be a compact manifold (smooth or PL) of dimension k and let X be a CW complex. Then every continuous map $f: W \rightarrow X$ is homotopic to an admissible map. If $f(\partial W) \subset X^{(k-1)}$ then the homotopy may be taken rel ∂W .*

Given an admissible map $f: W \rightarrow X$ from a compact k -manifold to a CW complex X , the k -volume of f , denoted $\text{Vol}_k(f)$, is defined to

be the number of open k -balls in W mapping to k -cells of X . We now have all the ingredients for a formal definition of the k -dimensional Dehn function.

Given a group G of type \mathcal{F}_{k+1} , fix an aspherical CW complex X with fundamental group G and finite $(k+1)$ -skeleton. Let \tilde{X} be the universal cover of X . If $f: S^k \rightarrow \tilde{X}$ is an admissible map, define the *filling volume* of f to be the minimal volume of an admissible extension of f to B^{k+1} :

$$\text{FVol}(f) = \min\{ \text{Vol}_{k+1}(g) \mid g: B^{k+1} \rightarrow \tilde{X}, g|_{\partial B^{k+1}} = f \}.$$

Note that extensions of f exist since \tilde{X} is contractible, and that we may assume these extensions are admissible by Lemma 2.1.

Now, the k -dimensional Dehn function of X is defined to be

$$\delta^{(k)}(x) = \sup\{ \text{FVol}(f) \mid f: S^k \rightarrow \tilde{X}, f \text{ is admissible, } \text{Vol}_k(f) \leq x \}.$$

This definition is the one given in [4], which is equivalent to the definitions given in [2, 8]. It is shown in [2] that, up to equivalence, $\delta^{(k)}(x)$ is a quasi-isometry invariant of \tilde{X} . In particular, the equivalence class of $\delta^{(k)}(x)$ does not depend on the particular $K(G, 1)$ complex used. Therefore, we will often write $\delta^{(k)}(x)$ as $\delta_G^{(k)}(x)$.

Remark 2.2. We are concerned in this paper primarily with 2- and 3-dimensional volume, which for clarity we write as $\text{Vol}_2(f) = \text{Area}(f)$ and $\text{Vol}_3(f) = \text{Vol}(f)$. We also abuse notation in the standard way by writing, for example, $\text{Vol}(Y)$ to mean $\text{Vol}(f)$ for some understood $f: Y \rightarrow X$.

We recall two very useful techniques from [4] for computing lower bounds for k -dimensional Dehn functions.

Remark 2.3. The first technique follows from the fact (see Remark 2.7 of [4]) that the volumes of top-dimensional embedded balls in a contractible complex are precisely the filling volumes of their boundary spheres. More precisely, if \tilde{X} is a contractible $(k+1)$ -complex, and $g: B^{k+1} \rightarrow \tilde{X}$ is an embedding, such that g and $g|_{S^k} = f$ are both admissible, then $\text{FVol}(f) = \text{Vol}_{k+1}(g)$. The proof is a standard homological argument; see [4] for details.

Remark 2.4. See Remark 2.1 of [4]. In order to establish the relation $f \preceq g$ between non-decreasing functions, it suffices to consider relatively sparse sequences of integers. For if (n_i) is an unbounded sequence of integers for which there is a constant $C > 0$ such that $n_0 = 1$ and $n_{i+1} \leq Cn_i$ for all i , and if $f(n_i) \leq g(n_i)$ for all i , then $f \preceq g$. Indeed, given $x \in [0, \infty)$ there is an index i such that $n_i \leq x \leq n_{i+1}$, whence $f(x) \leq f(n_{i+1}) \leq g(n_{i+1}) \leq g(Cn_i) \leq g(Cx)$.

In the case of the groups H_j in this paper, we produce a sequence of embedded balls $\{B_r^3\}_{r=1}^\infty$ and spheres $\{S_r^2\}_{r=1}^\infty$ in the universal cover of a 3-dimensional, aspherical $K(H_j, 1)$ complex. By Remark 2.3 $\text{Vol}(B_r^3)$ provides a lower bound for $\delta^{(2)}$ evaluated at $\text{Area}(S_r^2)$. In Lemma 5.17 we prove that there exist constants A_1, A_2 so that

$$A_1 r^2 \leq \text{Area}(S_r^2) \leq A_2 r^2.$$

The sequence of numbers $\{\text{Area}(S_r^2)\}_{r=1}^\infty$ satisfies the conditions of Remark 2.4, and so one obtains a lower bound for the function $\delta_{H_j}^{(2)}(x)$.

The situation is similar for the groups G_j , although the inequalities are a bit more subtle. We show directly in Lemma 5.18 that the sequence $\{\text{Area}(S_r^2)\}_{r=1}^\infty$ satisfies the conditions of Remark 2.4, and so lower bounds for $\delta_{G_j}^{(2)}(x)$ evaluated at $\text{Area}(S_r^2)$ (given by the 3-volume of the corresponding embedded 3-balls), translate into a general lower bound for the function $\delta_{G_j}^{(2)}(x)$.

3. OVERVIEW AND GEOMETRIC INTUITION

In Section 4 we define a sequence of groups G_n and H_n (for $n \geq 1$) with $\delta_{G_n}^{(2)}(x) \simeq \exp^n(x)$ and $\delta_{H_n}^{(2)}(x) \simeq \exp^n(\sqrt{x})$. In Section 5 lower bounds for these Dehn functions are established by exhibiting sequences of embedded 3-balls and boundary 2-spheres in the universal covers of 3-dimensional $K(G_n, 1)$ and $K(H_n, 1)$ complexes. Some of the details in these two sections may appear a little daunting on first reading. In order to motivate these groups and the sequences of 3-balls, we first investigate a classical construction for producing groups whose 1-dimensional Dehn functions are iterated exponentials.

3.1. A 1-dimensional Dehn function example and schematic diagrams. Consider how one may construct finitely presented groups

whose 1-dimensional Dehn functions are iterated exponential functions. The following examples are part of the folklore, but only appear to have been written down relatively recently [7]. Consider the family of finitely presented groups BS_n (for each positive integer n), defined by taking BS_1 to be the Baumslag-Solitar group

$$BS_1 = \langle a, t_1 \mid t_1 a t_1^{-1} = a^2 \rangle$$

and defining the BS_n ($n \geq 2$) as iterated HNN extensions

$$BS_{n+1} = \langle BS_n, t_{n+1} \mid t_{n+1} t_n t_{n+1}^{-1} = t_n^2 \rangle.$$

The group BS_n has Dehn function equivalent to the iterated exponential function $\exp^n(x)$.

The intuition is as follows. The commutator word

$$w_{1,m_1} = [t_1^{m_1} a t_1^{-m_1}, a]$$

has linear length in m_1 and has area $\exp(m_1)$ in BS_1 .

Now $BS_1 < BS_2$, and the extra relation in the presentation of BS_2 ensures that one can replace the segments $t_1^{m_1}$ in the commutator word above by words of the form $t_2^{m_2} t_1 t_2^{-m_2}$ of length approximately $\log_2(m_1)$. Thus, the word

$$w_{2,m_2} = [(t_2^{m_2} t_1 t_2^{-m_2}) a (t_2^{m_2} t_1 t_2^{-m_2})^{-1}, a]$$

has length a linear function of m_2 , but area $\exp^2(m_2)$ in BS_2 .

Continuing in this fashion, one can keep replacing subwords of the form $t_j^{m_j}$ by $t_{j+1}^{m_{j+1}} t_j t_{j+1}^{-m_{j+1}}$ where m_{j+1} is approximately $\log(m_j)$. One obtains a word w_{n,m_n} representing the identity in BS_n whose length is a linear function of m_n and whose area is $\exp^n(m_n)$. The lower bound on the area of the word w_{n,m_n} is obtained by realizing w_{n,m_n} as the boundary of an embedded van Kampen disk in the universal cover of the standard presentation 2-complex for BS_n . One verifies that the area of the embedded disk is $\exp^n(m_n)$. Since this universal cover is contractible, the standard homological argument from Remark 2.3 enables one to conclude that any van Kampen filling disk for w_{n,m_n} has at least as much area as the embedded one.

Figure 3.1 contains schematic diagrams of these embedded disks. In these schematics, we ignore all syllables of length 1, so the original




FIGURE 3.1. Schematic van Kampen disks for the words w_{i,m_i} .

commutator w_{1,m_1} appears as a square. Each replacement rule

$$t_j^{m_j} \mapsto t_{j+1}^{m_{j+1}} t_j t_{j+1}^{-m_{j+1}}$$

is represented by a triangle with base attached along an edge of the previous schematic diagram. The original square schematic for w_{1,m_1} becomes the octagon schematic for w_{2,m_2} , and this becomes the 16-gon schematic for w_{3,m_3} , and so on.

We reformulate the geometric intuition and combinatorics of these examples in terms of schematic diagrams. Start with a quadrilateral whose area is an exponential function of its perimeter length. Now take logs of the perimeter length by attaching triangles via their base edges. Metrically, we think of the two remaining edges of each triangle as having length that is approximately a log of the length of the base edge. We attach a generation of 4 triangles, then a second generation of 8 triangles, a third generation of 16 triangles and so on. Repeat this procedure as often as necessary to obtain a diagram whose area is an n -fold composite of exponential functions of its perimeter length.




FIGURE 3.2. The subdivision rule and associated triangular disk.

The combinatorics of the boundary circles of the schematic diagrams changes in a very simple fashion. Edges get subdivided into two at each stage. Topologically this subdivision is achieved by attaching triangular 2-disks along their base edges. This is shown in Figure 3.2.

Remark 3.1. Note that for each i the schematic diagram for w_{i,m_i} is a *template* for an infinite family of embedded van Kampen diagrams in the universal cover of the presentation 2-complex for BS_i . The infinite family is obtained by letting the integer m_i range from 1 to infinity.

3.2. The 2-dimensional Dehn function schematic procedure.

What complications arise when one tries to mimic the procedure of the previous subsection in higher dimensions? We will focus on the case of 2-dimensional Dehn functions. Start with a schematic 3-dimensional ball and then try to reduce the areas of portions of its boundary 2-sphere by attaching new 3-balls along its boundary. Each new 3-ball attaches along a 2-disk. The boundary of such a disc is a circle that survives on the boundary of the new 3-ball. A *collar neighborhood* of such a circle contributes to the area of the new boundary 2-sphere.

In order to be able to reduce area repeatedly by attaching successive generations of 3-balls, we will need to ensure that future generations of 3-balls attach in such a way as to cover up (large portions of) the boundary circles of the attaching disks of a given generation of 3-balls. This is a very basic complication in the combinatorics of attaching 3-balls that does not occur in the case of 1-dimensional Dehn functions and the examples of the previous subsection. In the 1-dimensional case the new 2-disks attach along 1-disks, the boundary of a 1-disk is a 0-sphere, and a collar neighborhood of a 0-sphere makes no significant contributions to length of the new boundary circle.

We first describe the combinatorics of our approach for dealing with the difficulty outlined in the preceding paragraph. We construct a sequence of schematic 3-balls B_i whose boundary 2-spheres S_i have the following properties.

- (i) Each S_i has a tiling into triangular regions.
- (ii) The boundary edges of these regions have two colors, labeled $(i - 1)$ and i .
- (iii) Each tile contains exactly one $(i - 1)$ -edge.

Now S_{i+1} is obtained from S_i by a *combination-subdivision* move as shown in Figure 3.3. Unlike the 1-dimensional case, where a simple subdivision procedure was sufficient, in this case one combines two cells together to form a quadrilateral region, and then one subdivides that region.

Each edge of color $(i-1)$ is the diagonal of a quadrilateral (combine two adjacent triangular regions). Delete this diagonal edge and retriangulate this quadrilateral by adding a barycenter and coning to the four vertices. Each new edge is of color $(i+1)$. Note that this gives a triangulation of the sphere S_{i+1} with just two colors, i and $(i+1)$, and now each triangle contains exactly one edge of color i . Thus one can repeat this procedure inductively.




FIGURE 3.3. The combination-subdivision rule and associated 3-ball.

We think of each sphere S_i as being the boundary of a 3-ball B_i . We think of each old quadrilateral as the back half of a 2-sphere and each new (subdivided) quadrilateral as the front half of the same 2-sphere. This 2-sphere bounds a 3-ball, as indicated in the lower half of Figure 3.3. The 3-ball B_{i+1} is obtained from B_i by attaching a

collection of such 3-balls along the quadrilateral neighborhoods of the edges with color $(i - 1)$.




FIGURE 3.4. Geometric versions of the combination-subdivision rule.

As in the case of the 1-dimensional Dehn function examples, the combination-subdivision moves have geometric content. Now we are concerned with areas as well as lengths. There are two geometric versions of the combination-subdivision rule. These are indicated in Figure 3.4, and details are given in Section 5.

- (i) In a Type I rule, edges of color i have length equal to m , edges of color $(i - 1)$ have length e^m , and the new edges of color $(i + 1)$ have length m . Each of the two old triangles have area e^m and the new triangles each have area m^2 .
- (ii) In a Type II rule, edges of color i and $(i - 1)$ have length equal to m , and the new edges of color $(i + 1)$ have length $\log(m)$. Each of the two old triangles have area m^2 and the new triangles each have area of order m .

These two types are designed so that one can alternate families of each type. In Section 5 we construct families of embedded balls and boundary spheres with the following schematic diagrams. The base




FIGURE 3.5. Alternating the combination-subdivision rules.

2-sphere S_1 is the join $S^0 * S^0 * S^0$, with the equator $S^0 * S^0$ being comprised of 4 edges of color 0, and the remaining 8 edges having color 1. We alternate these versions of the geometric combination-subdivision rule as we progress along the sequence of schematic 2-spheres. Figure 3.5 shows three fourths of the sphere S_1 , and the effect of the first few combination-subdivision rules.

The geometry of these families is as follows. The sphere S_1 bounds a 3-ball whose volume is a quadratic function of the area of S_1 . Applying a generation of Type I moves, we see that the sphere S_2 bounds a 3-ball whose volume is the square of $\exp(\sqrt{\text{Area}(S_2)})$, which is equivalent to $\exp(\sqrt{\text{Area}(S_2)})$. The sphere S_3 is obtained by applying Type II moves, and so bounds a 3-ball whose volume is $\exp(\text{Area}(S_3))$, and so on.

Remark 3.2. These schematic diagrams are related to the groups G_i and H_i ($i \geq 0$) of Section 4 as follows. For each integer $i \geq 0$, the

sphere S_{2i+1} is a template for 2-spheres in the universal cover of a 3-dimensional $K(G_i, 1)$ -complex, and the sphere S_{2i+2} is a template for 2-spheres in the universal cover of a 3-dimensional $K(H_i, 1)$ -complex.

Remark 3.3. In the case of the 1-dimensional Dehn function the schematic diagrams ignored syllables of length one. So a vertex may correspond to an edge of length 1 in a corresponding van Kampen diagram.

Similarly, vertices in these schematic sphere diagrams S_i may correspond to (several) 2-cells of the corresponding embedded spheres, and an edge may correspond to corridors on the corresponding spheres.

Remark 3.4. By analogy with the 1-dimensional case, one might be tempted in constructing the groups G_i and H_i to use Baumslag-Solitar type relations to provide the exponential scaling required by the Type I moves. For example, thinking of each of the triangles in the base quadrilateral of a Type I move as similar to the triangle in Figure 3.2, we could take the base quadrilateral to have boundary word $t^n a t^{-n} s^n a^{-1} s^{-n}$ in the group

$$B = \langle a, t, s \mid a^s = a^t = a^2 \rangle.$$

This quadrilateral is composed of two triangles (each with area $\exp(n)$) with boundary words $t^n a t^{-n} = a^{2^n}$ and $s^n a s^{-n} = a^{2^n}$.

The Type I move is executed by adding a stable letter u that acts on B via the endomorphism φ given by: $s \mapsto s, t \mapsto t, a \mapsto a^2$. The word $t^n a t^{-n} s^n a^{-1} s^{-n}$ is the boundary of a new quadrilateral, composed of four triangles (each of area n^2), and the new-color edges are u^n , $(s^{-1}u)^n$, u^n , and $(t^{-1}u)^n$. (See Figure 3.3.)

Observe, however, that the group

$$P = \langle B, u \mid g^u = \varphi(g), (g \in B) \rangle$$

is not an HNN extension with base B since the map φ is not injective. For instance $t^{-1} a t s^{-1} a^{-1} s \in \text{Ker}(\varphi)$.

It is important that each of groups in the present paper admits a graph-of-groups decomposition. This facet of our construction is exploited heavily in Section 5 where the graph of groups structure is used to conclude the existence of 3-dimensional classifying spaces, and

these 3–dimensional spaces are used to establish lower bounds for the 2–dimensional Dehn functions.

In Section 4 below the graph-of-groups structure is guaranteed by using mapping tori of free groups of rank 2 in place of Baumslag-Solitar groups.

4. THE GROUPS

Let F_2 denote the free group on two generators. To construct the two families of groups G_n and H_n , we start with the base group $F_2 \times F_2 \times F_2$ and then alternate two procedures, coning and attaching suspended wings, which are described below. The base group is of type \mathcal{F}_3 , and we shall see from the definitions of the two procedures that the groups obtained at each stage are also of type \mathcal{F}_3 .

As indicated in Remark 3.4 we will need to use an exponentially growing automorphism φ of F_2 repeatedly in the construction. It will be important (in Section 5.3) that φ be palindromic.

Definition 4.1 (The palindromic automorphism φ). Setting $F_2 = \langle \xi, \nu \rangle$, we define the automorphism $\varphi : F_2 \rightarrow F_2$ by

$$\varphi(\xi) = \xi\nu\xi \text{ and } \varphi(\nu) = \xi.$$

Remark 4.2 (Vector notation for free group bases). We use the vector notation \mathbf{y} to denote the basis $\{y_1, y_2\}$ for a free group of rank two. In Table 1, the vectors themselves may have subscripts. For example, \mathbf{u}_1 denotes the basis $\{u_{11}, u_{12}\}$ and \mathbf{a}_{23} denotes the basis $\{a_{231}, a_{232}\}$. Furthermore, an ordered list of k vectors describes an ordered basis of F_{2k} . For example, in Table 1, $\langle \mathbf{u}_0, \mathbf{y} \rangle$ denotes F_4 with the ordered basis $\{u_{01}, u_{02}, y_1, y_2\}$.

We also use product notation to denote coordinatewise multiplication of basis elements. For example, $\mathbf{u}_0^{-1}\mathbf{a}_{21}$ denotes the basis $\{u_{01}^{-1}a_{211}, u_{02}^{-1}a_{212}\}$.

We say that two vectors \mathbf{v} and \mathbf{w} commute if the two basis elements represented by \mathbf{v} commute with the two represented by \mathbf{w} .

The following definition provides the algebraic framework for the Type II moves from Section 3. We shall see explicit examples of these moves in Section 5.1.

Definition 4.3 (Coning). Suppose G is a group with a subgroup Γ isomorphic to $(F_2 * F_2 * \cdots * F_2) \times F_2$. We define the *cone of G over Γ* to be the fundamental group of the graph of groups in Figure 4.1.




FIGURE 4.1. The cone of G over Γ .

For each edge group, one of the edge maps is inclusion and the other edge map is $(\varphi * \varphi * \cdots * \varphi) \times \varphi$. This map is clearly injective, as φ is injective. Thus the cone of G over Γ is simply a double HNN extension of G . If G is of type \mathcal{F}_3 , then the cone of G over Γ is also of type \mathcal{F}_3 .

Example 4.4. In the first step of the inductive procedure, we start with $G = F_2 \times F_2 \times F_2$, where the three F_2 factors are generated by the 2-vectors \mathbf{a}_{01} , \mathbf{a}_{02} and \mathbf{y} . The subgroup Γ is $F_2 \times F_2 = \langle \mathbf{a}_{01} \rangle \times \langle \mathbf{a}_{02} \rangle$. Thus the presentation for $G_0 =$ the cone of G over Γ is given by

$$G_0 = \langle G, \mathbf{u}_0 \mid (g, h)^{u_{01}} = (g, h)^{u_{02}} = (\varphi(g), \varphi(h)), \forall (g, h) \in \Gamma \rangle$$

The next definition provides the algebraic framework for the Type I moves from Section 3. We shall see explicit examples of these moves in Section 5.1.

Definition 4.5 (Attaching suspended wings). Let G be a group containing a collection of subgroups $\{\Gamma_1, \dots, \Gamma_n\}$, each isomorphic to $F_2 \rtimes_{\theta} F_4$, where the map $\theta: F_4 \rightarrow \text{Aut}(F_2)$ is defined on the ordered basis $\{x_1, x_2, x_3, x_4\}$ as follows:

$$x_1 \mapsto \varphi, \quad x_2 \mapsto \varphi, \quad x_3 \mapsto \text{id}, \quad x_4 \mapsto \text{id}.$$

Consider the fundamental group of the graph of groups in Figure 4.2. For each edge group $F_2 \rtimes_{\theta} F_4$, the two edge maps are the inclusion map and the map $\varphi \times \text{id}$, which acts by φ on the F_2 factor and the identity on the F_4 factor. It is easy to check that the latter map is an injective homomorphism. We say that this group is obtained from G by *attaching suspended wings* to the collection of subgroups Γ_i .




FIGURE 4.2. Attaching suspended wings to the subgroups Γ_i .

Note that the edge groups Γ_i are all of type \mathcal{F}_2 . Thus, if G is of type \mathcal{F}_3 , then the new group is also of type \mathcal{F}_3 .

Example 4.6. In the group G_0 above, the subgroup $\Gamma_1 = \langle \mathbf{a}_{01}, \mathbf{u}_0, \mathbf{y} \rangle$ is isomorphic to the group $F_2 \rtimes_{\theta} F_4$ from Definition 4.5. Then the group obtained from G_0 by attaching suspended wings to Γ_1 has the presentation

$$\left\langle G_0, \mathbf{a}_{11} \mid \begin{array}{ll} g^{a_{111}} = g^{a_{112}} = \varphi(g) & \forall g \in F(\mathbf{a}_{01}), \\ g^{a_{111}} = g^{a_{112}} = g & \forall g \in F(\mathbf{u}_0) * F(\mathbf{y}) \end{array} \right\rangle$$

Example 4.7. The group H_1 in the inductive construction is obtained from G_0 by attaching suspended wings to Γ_1 and Γ_2 , where $\Gamma_1 = F_2 \rtimes_{\theta} F_4 = \langle \mathbf{a}_{01} \rangle \rtimes_{\theta} \langle \mathbf{u}_0, \mathbf{y} \rangle$ as in Example 4.6, and $\Gamma_2 = F_2 \rtimes_{\theta} F_4 = \langle \mathbf{a}_{02} \rangle \rtimes_{\theta} \langle \mathbf{u}_0, \mathbf{y} \rangle$. The pair of stable letters in H_1 corresponding to the edge group Γ_1 is denoted by \mathbf{a}_{11} and the pair of stable letters corresponding to the edge group Γ_2 is denoted by \mathbf{a}_{12} . This is summarized in the H_1 -row of Table 1.

4.1. Inductive definition of the groups G_n and H_n . In Example 4.4, the group G_0 was defined by applying the coning procedure to the base group $G = F_2 \times F_2 \times F_2$. In Example 4.7 the group H_1 was obtained from G_0 by attaching suspended wings. The groups G_n , $n \geq 1$ and H_n , $n \geq 2$ are defined inductively by alternating these two procedures.

The group G_n is obtained from H_n by coning over a subgroup of H_n that is isomorphic to $F_{2^{n+1}} \times F_2$, where $F_{2^{n+1}}$ is a free product

of 2^n copies of F_2 . The pair of new stable letters is denoted by \mathbf{u}_n . The F_2 factor of the subgroup $F_{2^{n+1}} \times F_2$ is generated by \mathbf{u}_{n-1} . The generators of the $F_{2^{n+1}}$ factor are listed in the G_n -row of Table 1.

The group H_n is obtained from G_{n-1} by attaching suspended wings to a collection of 2^n subgroups of G_{n-1} , each of which is isomorphic to $F_2 \rtimes_{\theta} F_4$ from Definition 4.5. Note that each of these subgroups labels two edges of the rose in Figure 4.2, so that there are actually 2^{n+1} new stable letters in H_n . These are labeled by 2^n vectors in the H_n -row of Table 1.

Lemma 4.11 establishes that the edge groups listed in Column 3 of Table 1 are indeed isomorphic to $F_{2^i} \times F_2$ or $F_2 \rtimes_{\theta} F_4$, depending on the case. By induction, the groups G_n and H_n are of type \mathcal{F}_3 for all n . Table 1 also includes the 2-dimensional Dehn functions of the resulting groups.

4.2. Further details about Table 1. This subsection provides a precise description of the edge groups $F_2 \rtimes_{\theta} F_4$ in the definition of H_n . On a first reading the reader may wish to focus on the first few groups in Table 1 (up to G_2). The arguments for the lower bounds for the first few groups (Section 5.1) and for the upper bounds in general (Sections 6 and 7) can be worked through without a thorough knowledge of the general labeling.

Recall that H_n is obtained from G_n by attaching suspended wings along a collection of 2^n subgroups of the form $F_2 \rtimes_{\theta} F_4$. These are labeled as follows:

$$\langle \mathbf{a}_{(n-1)i} \rangle \rtimes_{\theta} \langle \mathbf{u}_{n-1}, \mathcal{L}_n(i) \rangle, \quad \text{with } 1 \leq i \leq 2^{n-1}$$

and

$$\langle \mathbf{u}_{n-2}^{-1} \mathbf{a}_{(n-1)j} \rangle \rtimes_{\theta} \langle \mathbf{u}_{n-1}, \mathcal{L}_n(j + 2^{n-1}) \rangle, \quad \text{with } 1 \leq j \leq 2^{n-1}.$$

Lemma 4.11 establishes that these groups are indeed isomorphic to $F_2 \rtimes_{\theta} F_4$. Note that the generators of the F_2 factors are either the stable letters of H_{n-1} or are \mathbf{u}_{n-2}^{-1} times these stable letters. Note also that two of the generators of the F_4 factor are always \mathbf{u}_{n-1} , the stable letters in the definition of G_{n-1} . The remaining generators of the F_4 factor are described using the ordered list \mathcal{L}_n , which is defined

TABLE 1.

Group	Stable letters	Edge groups	$\delta^{(2)}(s)$
H_0	$\mathbf{a}_{01}, \mathbf{a}_{02}, \mathbf{y}$		$x^{3/2}$
G_0	\mathbf{u}_0	$\langle \mathbf{a}_{01} \rangle \times \langle \mathbf{a}_{02} \rangle$	x^2
H_1	\mathbf{a}_{11}	$\langle \mathbf{a}_{01} \rangle \rtimes_{\theta} \langle \mathbf{u}_0, \mathbf{y} \rangle$	$e^{\sqrt{x}}$
	\mathbf{a}_{12}	$\langle \mathbf{a}_{02} \rangle \rtimes_{\theta} \langle \mathbf{u}_0, \mathbf{y} \rangle$	
G_1	\mathbf{u}_1	$\langle \mathbf{a}_{11}, \mathbf{a}_{12} \rangle \times \langle \mathbf{u}_0 \rangle$	e^x
H_2	\mathbf{a}_{21}	$\langle \mathbf{a}_{11} \rangle \rtimes_{\theta} \langle \mathbf{u}_1, \mathbf{y} \rangle$	$e^{e\sqrt{x}}$
	\mathbf{a}_{22}	$\langle \mathbf{a}_{12} \rangle \rtimes_{\theta} \langle \mathbf{u}_1, \mathbf{y} \rangle$	
	\mathbf{a}_{23}	$\langle \mathbf{u}_0^{-1} \mathbf{a}_{11} \rangle \rtimes_{\theta} \langle \mathbf{u}_1, \mathbf{a}_{01} \rangle$	
	\mathbf{a}_{24}	$\langle \mathbf{u}_0^{-1} \mathbf{a}_{12} \rangle \rtimes_{\theta} \langle \mathbf{u}_1, \mathbf{a}_{02} \rangle$	
G_2	\mathbf{u}_2	$\langle \mathbf{a}_{21}, \mathbf{a}_{22}, \mathbf{a}_{23}, \mathbf{a}_{24} \rangle \times \langle \mathbf{u}_1 \rangle$	e^{e^x}
H_3	\mathbf{a}_{31}	$\langle \mathbf{a}_{21} \rangle \rtimes_{\theta} \langle \mathbf{u}_2, \mathbf{y} \rangle$	$e^{e^{e\sqrt{x}}}$
	\mathbf{a}_{32}	$\langle \mathbf{a}_{22} \rangle \rtimes_{\theta} \langle \mathbf{u}_2, \mathbf{y} \rangle$	
	\mathbf{a}_{33}	$\langle \mathbf{a}_{23} \rangle \rtimes_{\theta} \langle \mathbf{u}_2, \mathbf{a}_{01} \rangle$	
	\mathbf{a}_{34}	$\langle \mathbf{a}_{24} \rangle \rtimes_{\theta} \langle \mathbf{u}_2, \mathbf{a}_{02} \rangle$	
	\mathbf{a}_{35}	$\langle \mathbf{u}_1^{-1} \mathbf{a}_{21} \rangle \rtimes_{\theta} \langle \mathbf{u}_2, \mathbf{a}_{11} \rangle$	
	\mathbf{a}_{36}	$\langle \mathbf{u}_1^{-1} \mathbf{a}_{22} \rangle \rtimes_{\theta} \langle \mathbf{u}_2, \mathbf{a}_{12} \rangle$	
	\mathbf{a}_{37}	$\langle \mathbf{u}_1^{-1} \mathbf{a}_{23} \rangle \rtimes_{\theta} \langle \mathbf{u}_2, \mathbf{u}_0^{-1} \mathbf{a}_{11} \rangle$	
	\mathbf{a}_{38}	$\langle \mathbf{u}_1^{-1} \mathbf{a}_{24} \rangle \rtimes_{\theta} \langle \mathbf{u}_2, \mathbf{u}_0^{-1} \mathbf{a}_{12} \rangle$	
\vdots	\vdots	\vdots	\vdots
H_n ($n \geq 1$)	\mathbf{a}_{ni} $1 \leq i \leq 2^{n-1}$	$\langle \mathbf{a}_{(n-1)i} \rangle \rtimes_{\theta} \langle \mathbf{u}_{n-1}, \mathcal{L}_n(i) \rangle$ $1 \leq i \leq 2^{n-1}$	$\exp^n \sqrt{x}$
	\mathbf{a}_{ni} $2^{n-1} < i \leq 2^n$	$\langle \mathbf{u}_{n-2}^{-1} \mathbf{a}_{(n-1)j} \rangle \rtimes_{\theta} \langle \mathbf{u}_{n-1}, \mathcal{L}_n(j + 2^{n-1}) \rangle$ $1 \leq j \leq 2^{n-1}, j = i - 2^{n-1}$	
G_n ($n \geq 1$)	\mathbf{u}_n	$\langle \mathbf{a}_{nj} \rangle_{j=1}^{2^n} \times \langle \mathbf{u}_{n-1} \rangle$	$\exp^n x$

recursively as follows:

$$\mathcal{L}_1 = \{\mathbf{y}, \mathbf{y}\}$$

$$\mathcal{L}_2 = \{\mathbf{y}, \mathbf{y}, \mathbf{a}_{01}, \mathbf{a}_{02}\}$$

$$\mathcal{L}_n = \{\mathcal{L}_{n-1}, \mathbf{a}_{(n-2)1}, \dots, \mathbf{a}_{(n-2)2^{n-2}}, \mathbf{u}_{n-3}^{-1}\mathbf{a}_{(n-2)1}, \dots, \mathbf{u}_{n-3}^{-1}\mathbf{a}_{(n-2)2^{n-2}}\},$$

where this last equation is for $n \geq 3$. We let $\mathcal{L}_n(i)$ denote the i th element of \mathcal{L}_n . For $n = 1, 2$, and 3 these labels are given explicitly in Table 1.

We now establish a few facts that will be used repeatedly in the proof of Lemma 4.11. Let $M(G, \mathbf{t})$ denote a multiple HNN-extension with base group G and stable letters represented by the vector \mathbf{t} (which may have more than two coordinates).

Observation 4.8. If H is a subgroup of G such that the intersection of H with any of the edge groups of $M(G, \mathbf{t})$ is trivial, then the subgroup $\langle H, \mathbf{t} \rangle$ inside $M(G, \mathbf{t})$ is isomorphic to $H * \langle \mathbf{t} \rangle$.

In the following lemma, we use the notation of Remark 4.2.

Lemma 4.9. *Let $M(M(G, \mathbf{t}), \mathbf{s})$ be an iterated multiple HNN-extension. Let \mathbf{b} and \mathbf{c} be vectors consisting of elements of G with the following properties.*

- (i) *The intersection of $\langle \mathbf{c} \rangle$ with each edge group of $M(M(G, \mathbf{t}), \mathbf{s})$ is trivial.*
- (ii) *The vector \mathbf{b} has the same number of coordinates as \mathbf{t} .*

Then $\langle \mathbf{bt} \rangle$ is a free group whose rank equals the number of coordinates of \mathbf{t} and

$$\langle \mathbf{bt} \rangle \cap \langle \mathbf{s}, \mathbf{c} \rangle = 1.$$

Proof. There is a retraction ρ of $M(G, \mathbf{t})$ onto the free group F on \mathbf{t} , such that $\rho(g) = 1$ for all $g \in G$. The subgroup $\langle \mathbf{bt} \rangle$ of $M(G, \mathbf{t})$ has the same number of generators as F and maps onto F under this retraction. It is therefore free.

Now suppose $w_1 = w_1(\mathbf{s}, \mathbf{c})$ and $w_2 = w_2(\mathbf{bt})$ are words that represent the same element of $M(M(G, \mathbf{t}), \mathbf{s})$. Note that w_1 is contained in $\langle \mathbf{s}, \mathbf{c} \rangle$, which is isomorphic to $\langle \mathbf{s} \rangle * \langle \mathbf{c} \rangle$, by Condition (i) above and Observation 4.8. Moreover, w_1 represents an element of the base group for $M(M(G, \mathbf{t}), \mathbf{s})$, since w_2 does. These two facts imply that w_1 can

be freely reduced to a word that does not involve any coordinates of \mathbf{s} , say to $w_3 = w_3(\mathbf{c})$. Note that w_3 actually represents an element of G , so that $\rho(w_3) = 1$. This implies that $\rho(w_2) = 1$. Since w_2 is a word in \mathbf{bt} , this can only happen if w_2 is trivial. \square

Remark 4.10. The subgroup $\langle \mathbf{t} \rangle$ of $M(G, \mathbf{t})$ is free. Since $M(G, \mathbf{t}) \rightarrow M(M(G, \mathbf{t}), \mathbf{s})$ is an injection, the stable letters \mathbf{t} also generate a free group in $M(M(G, \mathbf{t}), \mathbf{s})$. This will be used repeatedly in the proof of Lemma 4.11.

Lemma 4.11 (Structure of edge groups). *The subgroup $\langle \mathbf{a}_{0i}, \mathbf{u}_0, \mathbf{y} \rangle$ (where $i = 1$ or 2) of G_0 is isomorphic to $F_2 \rtimes_{\theta} F_4$. Further, for each $n \geq 1$, the following statements hold.*

Coning:

- (1) *The subgroup $\langle \mathbf{a}_{n1}, \dots, \mathbf{a}_{n2^n}, \mathbf{u}_{n-1} \rangle$ of H_n is isomorphic to $F_{2^{n+1}} \times F_2$, where the F_2 factor has basis \mathbf{u}_{n-1} .*

Suspended wings:

- (2) *For $1 \leq i \leq 2^n$ the subgroup $\langle \mathbf{a}_{ni}, \mathbf{u}_n, \mathcal{L}_{n+1}(i) \rangle$ of G_n is isomorphic to $F_2 \rtimes_{\theta} F_4$, where F_2 has basis \mathbf{a}_{ni} and F_4 has ordered basis $\{\mathbf{u}_n, \mathcal{L}_{n+1}(i)\}$.*
- (3) *For $1 \leq j \leq 2^n$, the subgroup $\langle \mathbf{u}_{n-1}^{-1} \mathbf{a}_{nj}, \mathbf{u}_n, \mathcal{L}_{n+1}(j + 2^n) \rangle$ of G_n is isomorphic to $F_2 \rtimes_{\theta} F_4$, where F_2 has basis $\mathbf{u}_{n-1}^{-1} \mathbf{a}_{nj}$ and F_4 has ordered basis $\{\mathbf{u}_n, \mathcal{L}_{n+1}(j + 2^n)\}$.*

Proof. The proof is by induction on n .

For the base case, (the subgroup $\langle \mathbf{a}_{0i}, \mathbf{u}_0, \mathbf{y} \rangle$ of G_0 with $i = 1$ or 2) note that the subgroups $\langle \mathbf{a}_{0i} \rangle$, and $\langle \mathbf{y} \rangle$ are free factors of the base group $(F_2)^3$ (see Table 1). By Remark 4.10, $\langle \mathbf{u}_0 \rangle \simeq F_2$ since it is generated by the stable letters of G_0 . Now $\langle \mathbf{u}_0, \mathbf{y} \rangle$ is isomorphic to F_4 by Observation 4.8. The fact that $\langle \mathbf{u}_0, \mathbf{y} \rangle$ acts on $\langle \mathbf{a}_{0i} \rangle$ via θ is obvious by construction. Note that G_0 can be thought of as a single multiple HNN extension $M(F_2 \times F_2, \mathbf{y}, \mathbf{u}_0)$. Then $\langle \mathbf{u}_0, \mathbf{y} \rangle$, consisting of stable letters, intersects the subgroup $\langle \mathbf{a}_{0i} \rangle$ of the edge group trivially. This shows that $\langle \mathbf{a}_{0i}, \mathbf{u}_0, \mathbf{y} \rangle$ is isomorphic to $F_2 \rtimes_{\theta} F_4$.

Inductive step for (1): The group H_n is a multiple HNN extension $M(G_{n-1}, \mathbf{a}_{n1}, \dots, \mathbf{a}_{n2^n})$. So by Remark 4.10, the subgroup $\langle \mathbf{a}_{n1}, \dots, \mathbf{a}_{n2^n} \rangle$ is isomorphic to $F_{2^{n+1}}$. Further, it intersects G_{n-1} , and

hence $\langle \mathbf{u}_{n-1} \rangle$, trivially. Lastly, $\langle \mathbf{u}_{n-1} \rangle$ is isomorphic to F_2 since it consists of stable letters introduced in the construction of G_{n-1} . It follows that $\langle \mathbf{a}_{n1}, \dots, \mathbf{a}_{n2^n}, \mathbf{u}_{n-1} \rangle$ of H_n is isomorphic to $F_{2^{n+1}} \times F_2$, where the F_2 factor has basis \mathbf{u}_{n-1} .

Suspended wings: We first analyze the subgroups $\langle \mathcal{L}_{n+1}(i) \rangle$. Note that for $k \leq n$, the subgroups $\langle \mathbf{a}_{ki} \rangle$, where $1 \leq i \leq 2^k$, and $\langle \mathbf{u}_k \rangle$ of G_n are isomorphic to F_2 by Remark 4.10. Then the diagonal subgroups $\langle \mathbf{u}_{k-1}^{-1} \mathbf{a}_{ki} \rangle$ are isomorphic to F_2 as well. Since every vector $\mathcal{L}_{n+1}(i)$ is either \mathbf{a}_{ki} or $\mathbf{u}_{k-1}^{-1} \mathbf{a}_{ki}$ for some $k \leq n-1$, the subgroup $\langle \mathcal{L}_{n+1}(i) \rangle$ of G_n is isomorphic to F_2 for $1 \leq i \leq 2^{n+1}$.

Note that, as shown above, the edge group in the construction of G_n is isomorphic to $\langle \mathbf{a}_{nj} \rangle_{j=1}^{2^n} \times \langle \mathbf{u}_{n-1} \rangle$. In particular, $\langle \mathcal{L}_{n+1}(i) \rangle$ does not intersect the edge group, and Observation 4.8 implies that $\langle \mathbf{u}_n, \mathcal{L}_{n+1}(i) \rangle$ is isomorphic to F_4 for all $1 \leq i \leq 2^{n+1}$.

Inductive step for (2): Let $1 \leq i \leq 2^n$. By construction, the stable letters \mathbf{u}_n for G_n act on \mathbf{a}_{ni} via φ . The vector $\mathcal{L}_{n+1}(i)$ is the same as $\mathcal{L}_n(i)$, and the fact that $\mathcal{L}_n(i)$ commutes with \mathbf{a}_{ni} follows from the construction of H_n . This shows that $\langle \mathbf{u}_n, \mathcal{L}_{n+1}(i) \rangle$ acts on $\langle \mathbf{a}_{ni} \rangle$ via θ . Since $G_n = M(M(G_{n-1}, \mathbf{a}_{n1} \dots \mathbf{a}_{n2^n}), \mathbf{u}_n)$, Lemma 4.9 (with \mathbf{b} trivial and $\mathbf{c} = \mathcal{L}_{n+1}(i)$) shows that $\langle \mathbf{u}_n, \mathcal{L}_{n+1}(i) \rangle \cap \langle \mathbf{a}_{ni} \rangle$ is trivial. Thus the subgroup $\langle \mathbf{a}_{ni}, \mathbf{u}_n, \mathcal{L}_{n+1}(i) \rangle$ of G_n is isomorphic to $F_2 \rtimes_{\theta} F_4$.

Inductive step for (3): Let $1 \leq j \leq 2^n$. Then \mathbf{u}_n acts on \mathbf{a}_{nj} and \mathbf{u}_{n-1} (and hence on $\mathbf{u}_{n-1}^{-1} \mathbf{a}_{nj}$) via φ . Recall that

$$\mathcal{L}_{n+1}(j + 2^n) = \begin{cases} \mathbf{a}_{(n-1)j} & \text{for } 1 \leq j \leq 2^{n-1} \\ \mathbf{u}_{n-2}^{-1} \mathbf{a}_{(n-1)(j-2^{n-1})} & \text{for } 2^{n-1} < j \leq 2^n \end{cases}$$

and observe (from the definitions of G_{n-1} and H_n) that both \mathbf{u}_{n-1} and \mathbf{a}_{nj} act on $\mathcal{L}_{n+1}(j + 2^n)$ via φ . It follows that $\mathcal{L}_{n+1}(j + 2^n)$ commutes with $\mathbf{u}_{n-1}^{-1} \mathbf{a}_{(n)j}$. Thus $\langle \mathbf{u}_n, \mathcal{L}_{n+1}(j + 2^n) \rangle$ acts on $\langle \mathbf{u}_{n-1}^{-1} \mathbf{a}_{nj} \rangle$ via θ . The fact that these two groups have trivial intersection follows from Lemma 4.9 (taking $\mathbf{b} = \mathbf{u}_{n-1}^{-1}$ and $\mathbf{c} = \mathcal{L}_{n+1}(j + 2^n)$). Thus $\langle \mathbf{u}_{n-1}^{-1} \mathbf{a}_{nj}, \mathbf{u}_n, \mathcal{L}_{n+1}(j + 2^n) \rangle$ is isomorphic to $F_2 \rtimes_{\theta} F_4$. \square

4.3. The cell structure in K_{G_n} and K_{H_n} . In this section we describe 3-dimensional $K(\pi, 1)$'s for G_n and H_n . Start with the standard cell structure (one 0-cell and two 1-cells) on a bouquet of two circles.

The product of three copies of this with the product cell structure is defined to be K_{H_0} .

Since G_n and H_n are defined recursively as graphs of groups with 2-dimensional edge groups and 3-dimensional vertex groups, the complexes K_{G_n} and K_{H_n} are constructed inductively as total spaces of graphs of 3-dimensional vertex spaces and 2-dimensional edge spaces.

Here is the model situation. Let $X^{(3)}$ be a 3-dimensional cell complex and let $Y_1^{(2)}, \dots, Y_k^{(2)} \subset X^{(3)}$ be π_1 -injective 2-dimensional subcomplexes. For each i , let $\Phi_i: Y_i^{(2)} \rightarrow Y_i^{(2)}$ be a cellular map inducing an automorphism in π_1 . We define a new 3-complex $Z^{(3)}$ by

$$Z^{(3)} = X^{(3)} \sqcup \left(\sqcup_i Y_i^{(2)} \times [0, 1] \right) / \sim$$

where \sim identifies $Y_i^{(2)} \times 0$ with $Y_i^{(2)}$ (via the inclusion map) and $Y_i^{(2)} \times 1$ with $\Phi(Y_i^{(2)})$ for $1 \leq i \leq k$. Each $Y_i^{(2)} \times [0, 1]$ is given the product cell structure with a subdivision at the $Y_i^{(2)} \times 1$ end, obtained by pulling back the standard cell structure on $Y_i^{(2)}$ via Φ_i .

To get K_{G_n} from K_{H_n} we apply this model situation, taking $X^{(3)}$ to be K_{H_n} and the $Y_i^{(2)}$ to be the subcomplexes corresponding to the edge groups listed in the G_n -row of the third column of Table 1.

To get K_{H_n} from $K_{G_{n-1}}$ we first apply this model situation taking $X^{(3)}$ to be $K_{G_{n-1}}$ and the $Y_i^{(2)}$ to be the subcomplexes corresponding to the edge groups listed in the H_n -row of the third column of Table 1. We subdivide the resulting 3-complex by introducing edges labeled $\mathbf{u}_{n-1}^{-1} \mathbf{a}_{nj}$ and 2-cells corresponding to the commutation relations $[\mathbf{u}_{n-1}^{-1} \mathbf{a}_{nj}, \mathcal{L}_{n+1}(j + 2^n)] = 1$. (Refer to the edge groups in the H_{n+1} -row of Table 1 for these commutation relations.) Note that the \mathbf{a}_{nj} run over the stable letters of H_n . Figure 4.3 demonstrates such subdivisions in the case of K_{H_1} .

5. LOWER BOUNDS

5.1. Some spheres. As preparation for the general case, we explicitly describe some spheres in the first few groups. The computations of area and volume will be only roughly sketched, to help give a general idea. Formal verification of these computations will come later.




FIGURE 4.3. Typical new 1- and 2-cells in \widetilde{K}_{H_1} .

Spheres in G_0 . Fix $N \in \mathbb{N}$ and consider the slab in \widetilde{K}_{H_0} (the universal cover of K_{H_0}) of the form $\varphi^N(a_{011}) \times \varphi^N(a_{021}) \times y_1$ (the equatorial slab in Figure 5.1 below). The volume of this slab is $|\varphi^N(\xi)|^2$ (cf. Lemma 5.9).

Each of the top and bottom faces of this slab consists of 2-cells corresponding to relations in the edge group $\langle \mathbf{a}_{01} \rangle \times \langle \mathbf{a}_{02} \rangle$ in the definition of G_0 . Allowing u_{01}^{-N} to act on both faces produces the 3-ball shown in Figure 5.1. Let us call the boundary sphere of this ball




FIGURE 5.1. The sphere $S_1(N)$ in \widetilde{K}_{G_0} .

$S_1(N)$ in correspondence to Figure 3.5. Each of the eight trapezoidal faces of $S_1(N)$ is a van Kampen diagram for an equality of the form $\varphi^N(a_{0j1}) = u_{01}^N a_{0j1} u_{01}^{-N}$. In particular, each vertical side of the trapezoid has length N , the top has length one, and the bottom has length $|\varphi^N(\xi)|$, which is exponential in N . Thus the transition from the slab

to $S_1(N)$ corresponds to two Type II moves, one on each face of the slab.

We can think of each trapezoidal face on $S_1(N)$ as a stack of horizontal strips. Because of the exponential growth of φ , the lengths of these strips approximate a geometric series (Lemma 5.4). Thus the area of a trapezoid is approximately equal to the length of its base edge, which is the same as the length of one piece of the equatorial band about the sphere. We deduce that $\text{Area}(S_1(N)) \simeq |\varphi^N(\xi)|$ (Lemma 5.15). The volume of the 3-ball is bounded below by the volume of the slab, which is $|\varphi^N(\xi)|^2$, from which it follows that $\delta_{G_0}^{(2)}(x) \succeq x^2$ (cf. Remarks 2.3 and 2.4).

Spheres in H_1 . Looking at $S_1(N)$ from a different point of view, we see that it consists of four belted-trapezoid pairs, each of which has the geometry marking the beginning of a Type I move (see Figure 3.4). Each such trapezoid pair is contained in a group of the form $\langle \mathbf{a}_{0j} \rangle \rtimes_{\theta} \langle \mathbf{u}_0, \mathbf{y} \rangle$, an edge group in H_1 . We again think of each trapezoid as comprised of a stack of horizontal strips with boundary labeled $u_{01} \varphi^{i-1}(a_{0j1}) u_{01}^{-1} \varphi^i(a_{0j1}^{-1})$, where $1 \leq i \leq N$. The result of allowing a_{1j1}^{-i} to act on each such strip and a_{1j1}^{-N} to act on the four belts is the 3-ball shown in Figure 5.2. We call the boundary of this ball $S_2(N)$.




FIGURE 5.2. The sphere $S_2(N)$ in \widetilde{K}_{H_1} .

The long sides of the horizontal strips in Figure 5.2 represent words of the form a_{1j1}^N , giving each such strip an area equal to N . The angled strips connected to the poles have long sides representing words of the form $(a_{1j1}^{-1}u_{01})^N$, so that these strips also have area equal to N (recall from Section 4.3 that edges labeled $a_{1j1}^{-1}u_{01}$ are introduced into the cell structure on K_{H_1}). Each triangular region is a van Kampen diagram for an equation of the form $a_{1j1}^{-N}u_{01}^N = (a_{1j1}^{-1}u_{01})^N$ (recall that a_{1j1} commutes with u_{01}). The area of such a triangle is essentially the number of commutation relations required to achieve this equality, which is of order N^2 (Lemma 5.2). Thus the area of $S_2(N)$ is of order N^2 . On the other hand, the volume of the 3-ball is again bounded below by the volume of the slab, which is $|\varphi^N(\xi)|^2 \simeq (e^N)^2 \simeq e^N$. It follows that $\delta_{H_1}^{(2)}(x) \succeq e^{\sqrt{x}}$.

Spheres in G_1 . To form $S_3(N)$ in \widetilde{K}_{G_1} , we alter the previous constructions slightly. We still begin with a slab of the form $\varphi^N(a_{011}) \times \varphi^N(a_{021}) \times y_1$, but now we set $N = |\varphi^M(\xi)|$ for some M , so that the volume of the slab, as a function of M , is $|\varphi^{|\varphi^M(\xi)|}| \simeq e^{e^M}$. We next allow the word $\varphi^M(u_{01}^{-1})$ (instead of u_{01}^{-N}) to act on the top and bottom faces of the slab, producing a sphere analogous to the one in Figure 5.1. Performing Type I moves as before, but with $\varphi^M(a_{1j1})$ acting on the belts instead of a_{1j1}^N , produces a sphere analogous to the one in Figure 5.2, but with labels changed as indicated in Figure 5.3. The triangles in this sphere can be grouped in pairs to form eight quadrilaterals, each with the geometry marking the beginning of a Type II move. Each of these quadrilaterals is a van Kampen diagram labeled by elements of the group $\langle \mathbf{a}_{11}, \mathbf{a}_{12} \rangle \times \langle \mathbf{u}_0 \rangle$, the edge group for G_1 . We now allow u_{11}^{-M} to act on each of these quadrilaterals. As in G_0 , this will produce trapezoids representing equalities of the form $u_{11}^M g u_{11}^{-M} = \varphi^M(g)$, where g is of the form either a_{1j1} or $a_{1j1}^{-1}u_{01}$. These relations have area of order $|\varphi^M(\xi)| \simeq e^M$, which gives the area of the sphere. The volume of the slab is $\simeq e^{e^M}$, so that $\delta_{G_1}^{(2)}(x) \succeq e^x$.

5.2. Iteration notation. Before moving on to the general case, we pause briefly to develop the iteration notation necessary to avoid unwieldy towers of exponents.




FIGURE 5.3. A portion of a variation of the sphere in \widetilde{K}_{H_1} shown in Figure 5.2. The triangle pair admits an action by u_{11}^{-M} .

Recall that the map $\varphi : \langle \xi, \nu \rangle \rightarrow \langle \xi, \nu \rangle$ has the form $\varphi(\xi) = \xi\nu\xi$ and $\varphi(\nu) = \xi$. We inductively define functions $w_n : \mathbb{N} \rightarrow \mathbb{N}$, setting $w_0(r) = r$ for all $r \in \mathbb{N}$, and

$$w_n(r) = |\varphi^{w_{n-1}(r)}(\xi)|.$$

The growth of $|\varphi^N(\xi)|$ is determined by the matrix $\begin{bmatrix} 2 & 1 \\ 1 & 0 \end{bmatrix}$, which has positive eigenvalue $1 + \sqrt{2}$. More precisely, we have that

$$|\varphi^N(\xi)| = \left\| \begin{bmatrix} 2 & 1 \\ 1 & 0 \end{bmatrix}^N \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right\|_1 \approx \frac{1}{2}(1 + \sqrt{2})^{N+1}.$$

This implies that

$$\lim_{k \rightarrow \infty} \frac{|\varphi^{k+1}(\xi)|}{|\varphi^k(\xi)|} = 1 + \sqrt{2}.$$

It follows easily that

$$\sum_{k=0}^N |\varphi^k(\xi)| \simeq |\varphi^N(\xi)|,$$

as this sum is approximately geometric. From this discussion we deduce the following.

Lemma 5.1 (Growth of w_n). *For each n we have*

$$\sum_{i=0}^{w_{n-1}(r)} |\varphi^i(\xi)| \simeq w_n(r) \simeq \exp^n(r).$$

5.3. The van Kampen diagrams for moves. We now describe the van Kampen diagrams that will make up the surface of our spheres, and we explain how they correspond to the schematic diagrams and the subdivision-combination moves.

Given two words $W_1 = b_1 b_2 \cdots b_m$ and $W_2 = c_1 c_2 \cdots c_m$, we let $\delta(W_1, W_2)$ denote the word $b_1 c_1 b_2 c_2 \cdots b_m c_m$. With this notation, we define $\Delta_{ij}^n(w_k(r))$, for $k \geq 1$, $n \geq 0$, and $1 \leq i, j \leq 2^n$, to be the van Kampen diagram in \widetilde{K}_{H_n} shown in Figure 5.4.




FIGURE 5.4. The van Kampen diagram $\Delta_{ij}^n(w_k(r))$.

Such a van Kampen diagram will be found in the initial stage of a Type II move. The following lemma establishes the existence and geometry of such van Kampen diagrams.

Lemma 5.2. *The van Kampen diagram $\Delta_{ij}^n(w_k(r))$ exists. Moreover, $\text{Area}(\Delta_{ij}^n(w_k(r))) \simeq [w_k(r)]^2$.*

Proof. Because φ is palindromic, we may write

$$\varphi^{w_{k-1}(r)}(a_{ni1}) = b_1 \cdots b_p b_{p+1} b_p \cdots b_1,$$

where for $j \neq p+1$ we have $b_j \in \{a_{ni1}^\pm, a_{ni2}^\pm\}$, and $b_{p+1} \in \{a_{ni1}^\pm, a_{ni2}^\pm, e\}$. We similarly write

$$\varphi^{w_{k-1}(r)}(u_{(n-1)1}) = c_1 \cdots c_p c_{p+1} c_p \cdots c_1.$$

Recall from Table 1 that a_{nij} commutes with $u_{(n-1)j}$ for $1 \leq i \leq 2^n$ and $j = 1, 2$. Then

$$\begin{aligned} \varphi^{w_{k-1}(r)}(a_{ni1}^{-1})\varphi^{w_{k-1}(r)}(u_{(n-1)1}) &= (b_1 \cdots b_p b_{p+1} b_p \cdots b_1)^{-1} (c_1 \cdots c_p c_{p+1} c_p \cdots c_1) \\ &= (b_1^{-1} c_1) \cdots (b_p^{-1} c_p) (b_{p+1}^{-1} c_{p+1}) (b_p^{-1} c_p) \cdots (b_1^{-1} c_1) \\ &= \delta(\varphi^{w_{k-1}(r)}(a_{ni1}^{-1}), \varphi^{w_{k-1}(r)}(u_{(n-1)1})). \end{aligned}$$

The area claim follows by counting the number of commutation relations involved in this equality, remembering that each such relation provides two 2-cells due to the diagonal subdivision in the cell structure. We obtain

$$\begin{aligned} \text{Area}(\Delta_{ij}^n(w_k(r))) &= 2 \left(\sum_{j=1}^{2p} j + (2p+1) + \sum_{j=1}^{2p} j \right) \\ &= 2(2p+1)^2 = 2|\varphi^{w_{k-1}(r)}(g_1)|^2 \simeq [w_k(r)]^2. \end{aligned}$$

□

Remark 5.3. The reason φ is chosen to be palindromic is to ensure the existence of such van Kampen diagrams. To understand the mechanics behind $\Delta_{ij}^n(N)$, consider the subgroup $\langle g_1^{-1}h_1, g_2^{-1}h_2 \rangle \cong F_2$ of the group $\langle g_1, g_2 \rangle \times \langle h_1, h_2 \rangle \cong F_2 \times F_2$. The action of $\varphi \times \varphi$ on $F_2 \times F_2$ restricts to an automorphism of this subgroup exactly when φ is palindromic, and in this case the induced action is the same as the action of φ coming from the isomorphism $\langle g_1^{-1}h_1, g_2^{-1}h_2 \rangle \simeq F_2$. Note that the diagonal sides of $\Delta_{ij}^n(w_k(r))$ are labeled by the generators of such subgroups, and are in fact of the form $\varphi^{w_{k-1}(r)}(g_1^{-1}h_1)$. This fact is crucial for performing future generations of the iterative procedure.

When $k = 0$ we define $\Delta_{ij}^n(w_0(r))$ to be the van Kampen diagram obtained from the one in Figure 5.4 by replacing all labels of the form $\varphi^{w_{k-1}(r)}(g)$ by g^r . Also the notation $\Delta_{12}^0(w_k(r))$ will make sense with the definition above if we define $\mathbf{u}_{-1} = \mathbf{a}_{01}\mathbf{a}_{02}^{-1}$. For

then we have $\delta(\varphi^{w_{k-1}(r)}(a_{011}^{-1}), \varphi^{w_{k-1}(r)}(a_{011}a_{021}^{-1})) = \varphi^{w_{k-1}(r)}(a_{021}^{-1}) = [\varphi^{w_{k-1}(r)}(a_{021})]^{-1}$. Each of these diagrams has geometry identical to that of the others, up to \simeq -equivalence. See Figure 5.5.




FIGURE 5.5. The van Kampen diagrams $\Delta_{12}^0(w_k(r))$ and $\Delta_{12}^0(w_0(2))$.

For the other van Kampen diagram with which we will be concerned, we first recall that $\mathcal{L}_n(i)$ denotes the i th element of the list \mathcal{L}_n . This element is a 2-vector $\langle \xi, \nu \rangle$. It will be useful in what follows to let $\mathcal{L}_n^{(1)}(i)$ denote the first element ξ of this vector. With this notation we define $\Theta_i^n(w_k(r))$, for $k \geq 1$, $n \geq 0$, and $1 \leq i \leq 2^n$, to be the van Kampen diagram in $\widetilde{K_{G_n}}$ shown in Figure 5.6. (Recall that each vector in the list \mathcal{L}_{n+2} consists of elements in G_n .) When $k = 0$ we define $\Theta_i^n(w_0(r))$ to be the van Kampen diagram obtained from that in Figure 5.6 by replacing all labels of the form $\varphi^{w_{k-1}(r)}(g)$ by g^r .

We establish the existence and geometry of such diagrams in the following lemma.

Lemma 5.4. *The van Kampen diagram $\Theta_i^n(w_k(r))$ exists. Moreover, $\text{Area}(\Theta_i^n(w_k(r))) \simeq w_{k+1}(r)$.*

For the intuition behind this result, see $\Theta_2^1(w_0(1))$ in Figure 5.6. The crucial ingredient for the area claim is the exponential growth of φ .




FIGURE 5.6. The van Kampen diagrams $\Theta_i^n(w_k(r))$ and $\Theta_2^1(w_0(1))$.

Proof. To establish the existence of the van Kampen diagram, note that each of u_{n1} and u_{n2} acts on $\mathcal{L}_{n+2}^{(1)}(i+2^{n+1})$ via φ . Thus as elements of G_n , the word $\varphi^{w_{k-1}(r)}(u_{n1})\mathcal{L}_{n+2}^{(1)}(i+2^{n+1})\varphi^{w_{k-1}(r)}(u_{n1}^{-1})$ and the word labeling the long sides of the central strip in $\Theta_i^n(w_k(r))$ are equal. This establishes the top and bottom trapezoids of $\Theta_i^n(w_k(r))$.

For the middle strip, we simply need to observe that $\mathcal{L}_{n+2}^{(1)}(i)$ commutes with $\mathcal{L}_{n+2}(i+2^{n+1})$. To see this, note that each $\mathcal{L}_{n+2}(i+2^{n+1})$ term is also the first factor in an edge group for H_{n+1} , while each $\mathcal{L}_{n+2}^{(1)}(i)$ term is the third factor in the corresponding group. These commute, by definition of the action of θ .

We now compute the area of $\Theta_i^n(w_k(r))$. Note that the area obtained when φ acts on a positive word W is exactly $|W|$. Since φ is applied $|\varphi^{w_{k-1}(r)}(\xi)| \simeq w_k(r)$ times in each trapezoid, its area is

$$1 + |\varphi(\xi)| + |\varphi^2(\xi)| + \cdots + |\varphi^{w_k(r)-1}(\xi)|,$$

so that $\text{Area}(\Theta_i^n(w_k(r))) \simeq |\varphi^{w_k(r)}(\xi)| = w_{k+1}(r)$, as required. \square

The geometry of the van Kampen diagrams can be summarized as shown in Figure 5.7, where the labels on the edges indicate lengths.

5.4. Geometry of the moves.

5.4.1. *Type II.* As indicated above, the geometry of the van Kampen diagram $\Delta_{ij}^n(w_k(r))$ is that of the schematic quadrilateral with which




FIGURE 5.7. The geometry of $\Delta_{ij}^n(w_k(r))$ and $\Theta_i^n(w_k(r))$.

a Type II move begins, in that each side has equivalent length, and the area is equivalent to the square of this length. We now explain how such a move is performed in the group G_n . Note that $\Delta_{ij}^n(w_k(r))$ is contained in the edge group for G_n . As such it admits an action by the stable letters \mathbf{u}_n .

Lemma 5.5. *Assume $k \geq 1$, and suppose W is a positive word in the u_{nj} of length $w_{k-1}(r)$, and let B denote the abstract 3-ball obtained by allowing W^{-1} to act on $\Delta_{ij}^n(w_k(r))$. Let S denote the 2-sphere boundary of B . Then B is embedded in \widetilde{K}_{G_n} . Moreover*

$$\text{Vol}(B) \succeq w_k^2(r)$$

and

$$\text{Area}(S - \Delta_{ij}^n(w_k(r))) \simeq w_k(r).$$

Proof. Note that $w_k^2(r)$ is exactly the area of the van Kampen diagram $\Delta_{ij}^n(w_k(r))$. In the bottom layer of the 3-ball there is at least one 3-cell for every nine 2-cells in $\Delta_{ij}^n(w_k(r))$. The volume claim follows.

The boundary of $\Delta_{ij}^n(w_k(r))$ consists of four words of the form $\varphi^{w_{k-1}(r)}(g)$ for some $g \in \{a_{ni1}, a_{nj1}, a_{ni1}^{-1}u_{(n-1)1}, a_{nj1}^{-1}u_{(n-1)1}\}$. The action of W^{-1} on each of these words has the geometry of one trapezoid of a $\Theta_i^n(w_{k-1}(r))$. As such the area claim follows from the argument in the proof of Lemma 5.4.

Finally, suppose for contradiction that v_1 and v_2 are vertices of B that are identified in \widetilde{K}_{G_n} , let σ be an edge path in B joining them. Note that B is contained in a subgroup of G_n of the form $(F_4 \times F_2) \rtimes F_2$,




FIGURE 5.8. A view from above of the action of W on $\Delta_{ij}^n(w_k(r))$, as well as the resulting 3-ball in the case that $W = \varphi^{w_{k-2}(r)}(u_{n1})$. Note that each vertical quadrilateral on the boundary of the 3-ball is identical to a portion of $\Theta_i^n(w_{k-1}(r))$.

where the $F_4 \times F_2$ factor is the edge group and the final F_2 is generated by the stable letters in the entry for G_n in Table 1.

If w is the word corresponding to σ , then we may write $w =_{G_n} w_1 w_2 w_3$, where w_1 , w_2 , and w_3 are words in the generators of the three factors F_4 , $F_2 = \langle \mathbf{u}_{n-1} \rangle$, and $F_2 = \langle \mathbf{u}_n \rangle$, respectively (the stable letters \mathbf{u}_n act by automorphism, so that we may move them to the right, giving w_3 ; commutation relations in the base group then separate the \mathbf{a}_{nj} from the \mathbf{u}_{n-1}). There is a corresponding proper homotopy of σ to a path σ' corresponding to $w_1 w_2 w_3$.

Now the fact that v_1 and v_2 are identified in \widetilde{K}_{G_n} means that w represents the trivial element in G_n , and this implies that each of the subwords w_1 , w_2 , and w_3 is trivial in its corresponding group, which, being free, means that each w_i freely reduces to the empty word. Thus σ' is homotopic *with endpoints fixed* to a point. It follows that $v_1 = v_2$, so that B is embedded in \widetilde{K}_{G_n} . \square

Remark 5.6. When $k \geq 2$ and $W = \varphi^{w_{k-2}(r)}(u_{n1})$, the action of W^{-1} on each of the four boundary words of $\Delta_{ij}^n(w_k(r))$ produces precisely a trapezoid of $\Theta_i^n(w_{k-1}(r))$, as shown in Figure 5.8. When $k = 1$ and $W = u_{n1}^r$, it produces a trapezoid of $\Theta_i^n(w_0(r))$.

5.4.2. *Type I.* The geometry of $\Theta_i^n(w_k(r))$ is that of the schematic quadrilateral with which a Type I move begins, in that each side of the quadrilateral has equivalent length, while the length of the diagonal and the total area are both exponential in the outer edge length. We now explain how a Type I move is performed on such a van Kampen diagram in H_{n+1} . Note that each $\Theta_i^n(w_k(r))$ is contained in some single edge group for H_{n+1} (depending on i). As such it admits an action by the stable letters $\mathbf{a}_{(n+1)j}$, for some j depending on i . In particular, given a word V in the $a_{(n+1)ji}^{-1}$, we may perform the procedure analogous to the one performed on each belted trapezoid pair of $S_1(N)$ in the construction of $S_2(N)$ (cf. Section 5.1). In the following lemma, this is referred to as *allowing V to act on $\Theta_i^n(w_k(r))$* .

Lemma 5.7. *Suppose W is a positive word in the a_{nji} of length $w_k(r)$, and let B denote the 3-ball in H_{n+1} obtained by allowing W^{-1} to act on $\Theta_i^n(w_k(r))$. Let S denote the 2-sphere boundary of B . Then B is embedded in $\widetilde{K}_{H_{n+1}}$. Moreover*

$$\text{Vol}(B) \succeq w_{k+1}(r),$$

and

$$\text{Area}(S - \Theta_i^n(w_k(r))) \simeq [w_k(r)]^2.$$

Proof. The lower volume bound is trivial, as this is the length of the central strip, and there is at least one 3-cell for every three 2-cells in this strip.

Suppose first that $k \geq 1$. Then the boundary of $\Theta_i^n(w_k(r))$ consists of four copies of the word $\varphi^{w_{k-1}(r)}(u_{n1})$ along with four words of length one (those from \mathcal{L}_{n+2}). The action of $\varphi^{w_{k-1}(r)}(a_{nj1}^{-1})$ on $\varphi^{w_{k-1}(r)}(u_{n1})$ has the geometry of one-half of a $\Delta_{ij}^n(w_k(r))$ diagram. As such the area of each of the corresponding pieces of $S - \Theta_i^n(w_k(r))$ is $\simeq [w_k(r)]^2$. Because each letter in $\varphi^{w_{k-1}(r)}(a_{nj1}^{-1})$ commutes with each of the letters in \mathcal{L}_{n+2} , the area resulting from the action on these words is $\simeq w_k(r)$. The area claim follows for this case.




FIGURE 5.9. A view from above of the action of W on $\Theta_i^n(w_k(r))$, as well as the resulting 3-ball in the case that $W = \varphi^{w_{k-2}(r)}(u_{n1})$. Note that each triangle on the boundary of the 3-ball is identical to a portion of $\Delta_{ij}^{n+1}(w_k(r))$.

For the case that $k = 0$, the same argument applies with $\varphi^{w_{k-1}(r)}(g)$ replaced by g^r .

The proof that B is embedded in $\widetilde{K_{H_{n+1}}}$ is essentially the same as in the case of $\widetilde{K_{G_n}}$. The ball B is contained in a subgroup of H_{n+1} of the form $(F_2 \rtimes_{\theta} F_4) \rtimes F_2$, where the $F_2 \rtimes_{\theta} F_4$ factor is an edge group and the final F_2 is generated by the corresponding stable letters in the entry for H_{n+1} in Table1.

A path joining two vertices v_1 and v_2 in B that are identified in $\widetilde{K_{H_{n+1}}}$ corresponds to a word w with $w =_{H_{n+1}} w_1 w_2 w_3$, where each of w_1 , w_2 , and w_3 freely reduces to the empty word. This implies as before that $v_1 = v_2$. \square

Remark 5.8. When $W = \varphi^{w_{k-1}(r)}(a_{nj1})$, the action of W^{-1} on $\varphi^{w_{k-1}(r)}(u_{n1})$ produces precisely one half of $\Delta_{ij}^n(w_k(r))$.

5.5. Lower bound estimates. Base Cases. We now inductively compute lower bounds for $\delta_{G_n}^{(2)}(x)$ and $\delta_{H_n}^{(2)}(x)$. For the base cases,

we return to the spheres described in Section 5.1 and point out how the claims made there are verified by the computations just obtained. The only ingredient we lack is the following.

Lemma 5.9 (Slab volume). *The volume of the slab $\varphi^{w_k(r)}(a_{011}) \times \varphi^{w_k(r)}(a_{021}) \times y_1$ is $[w_{k+1}(r)]^2$.*

Proof. As a cell-complex this slab is isomorphic to $\Delta_{12}^0(w_{k+1}(r)) \times [0, 1]$ with the product cell structure. The result then follows from Lemma 5.2. \square

For G_0 we take $k = 0$, and so begin with the slab $\varphi^r(a_{011}) \times \varphi^r(a_{021}) \times y_1$ with volume $[w_1(r)]^2$. The boundary of this slab is a topological sphere containing two copies of $\Delta_{12}^0(w_1(r))$. Let $B_{G_0}(r)$ denote the 3-ball obtained by allowing u_{01}^{-r} to act on both of these. Let $S_{G_0}(r)$ denote the 2-sphere boundary of $B_{G_0}(r)$.

Lemma 5.10 (Area and volume in G_0). *Area($S_{G_0}(r)$) $\simeq w_1(r)$ and Vol($B_{G_0}(r)$) $\succeq [w_1(r)]^2$.*

Proof. The volume claim follows immediately from the volume of the slab, computed above. For the area, we note that the band around the slab consists of four pieces, each with area $|\varphi^r(\xi)| = w_1(r)$. The remaining area is shown to be $\simeq w_1(r)$ by applying Lemma 5.5 with $k = 0$. \square

We now consider H_1 . Note that $S_{G_0}(r)$ consists of four copies of $\Theta_i^0(w_0(r))$ (two with $i = 1$, two with $i = 2$) plus two additional 2-cells. We define $B_{H_1}(r)$ to be the 3-ball obtained by allowing a_{111}^r and a_{121}^r to act on the corresponding van Kampen diagrams. We let $S_{H_1}(r)$ denote the 2-sphere boundary of this ball.

Proposition 5.11 (Area and volume in H_1). *Area($S_{H_1}(r)$) $\simeq r^2$ and Vol($B_{H_1}(r)$) $\succeq [w_1(r)]^2$.*

Proof. We again obtain Vol($B_{H_1}(r)$) $\succeq [w_1(r)]^2$ from the volume of the central slab. For the area we use Lemma 5.7 to compute

$$\text{Area}(S_{H_1}(r)) = 8\text{Area}(\Delta_{12}^1(r)) + 16r + 6 \simeq r^2.$$

\square

Proposition 5.12 (Lower bounds for G_0 and H_1).

$$\delta_{G_0}^{(2)}(x) \succeq x^2 \quad \delta_{H_1}^{(2)}(x) \succeq e^{\sqrt{x}}$$

Proof. By Remark 2.3, (and assuming Lemmas 5.17 and 5.18 below), it suffices now to show that $B_{G_0}(r)$ and $B_{H_1}(r)$ are embedded in \widetilde{K}_{G_0} and \widetilde{K}_{H_1} , respectively (but see Remark 5.13 below).

Note that $B_{G_0}(r)$ consists of three component balls: the initial equatorial slab and the two balls resulting from the two Type II moves. Call these latter two *Type II balls*. Each of these three balls is embedded in \widetilde{K}_{G_0} , by Lemma 5.5. If we label a vertex in the slab with the identity element $e \in G_0$, the other vertices in $B_{G_0}(r)$ naturally inherit labels by group elements. To show that $B_{G_0}(r)$ is embedded, it suffices to show that any two vertices in the cones that carry the same label must lie in the same Type II ball.

Elements $g \in G_0$ labeling vertices in a Type II ball satisfy $g = hs$, for some $h \in H_0$ and $s \in \langle \mathbf{u}_0 \rangle$ (this is because $\langle \mathbf{u}_0 \rangle$ acts by automorphisms). If two such elements g_1 and g_2 are equal in G_0 , then $h_1 s_1 = h_2 s_2$. By considering the natural retraction $G_0 \rightarrow \langle \mathbf{u}_0 \rangle$, we see then that $s_1 = s_2$, from which it follows that $h_1 = h_2$. Note that h_1 and h_2 label vertices on the embedded sphere boundary of the slab. Because distinct Type II balls are attached to this sphere along disjoint disks, $h_1 = h_2$ implies that the vertices labeled by g_1 and g_2 lie in the same Type II ball.

For $B_{H_1}(r)$ the argument is similar. We think of $B_{H_1}(r)$ as having five component balls: $B_{G_0}(r)$ and four *Type I balls*. As before, it suffices to show that any two vertices in the Type I balls carrying the same label must lie in the same Type I ball.

Elements h labelling vertices in a Type I ball satisfy $h = gs$, where $g \in G_0$ and $h \in \langle \mathbf{a}_{1j} \rangle$. If two such elements h_1 and h_2 are equal in H_1 , we deduce as before that $s_1 = s_2$. In particular, the two corresponding vertices lie in Type I balls corresponding to the same stable group. Such balls are attached along disjoint disks on the embedded sphere $S_{G_0}(r)$. As before we deduce that $h_1 = h_2$ only if the corresponding vertices lie in the same Type I ball. \square

Remark 5.13. The equivalence relation \simeq behaves poorly under composition. In particular, one cannot always deduce the nature of volume

as a function of area given (in)equivalencies of each as functions of r . It is not difficult to show, however, that these difficulties do not arise for the particular pairs of functions we encounter.

5.6. Induction steps. Here we put together the pieces of the previous subsections to compute lower bounds for $\delta_{H_n}^{(2)}(x)$ and $\delta_{G_n}^{(2)}(x)$.

Lower Bounds in H_n . Begin with a slab in $(F_2)^3$ of the form $\varphi^{w_{n-1}(r)}(a_{011}) \times \varphi^{w_{n-1}(r)}(a_{021}) \times y_1$, and note that the volume of this slab is $[w_n(r)]^2$. Assume by induction that a sphere $S_{G_{n-1}}(r)$ has been built upon this slab via alternating Type I and Type II moves, so that $S_{G_{n-1}}(r)$ consists of

- 2^{2n} copies of $\Theta_i^{n-1}(w_0(r))$; and
- $\sum_{k=1}^{2n-1} 2^k$ individual 2-cells.

These hypotheses are verified above for G_0 ($n = 1$). Let $B_{H_n}(r)$ denote the 3-ball obtained by applying a Type II move using a_{nij}^{-r} to each copy of $\Theta_i^{n-1}(w_0(r))$, and let $S_{H_n}(r)$ denote its boundary 2-sphere.

Lemma 5.14. *The 2-sphere $S_{H_n}(r)$ consists of*

- 2^{2n+1} copies of $\Delta_{ij}^n(w_0(r))$;
- 2^{2n+2} strips with dimensions $r \times 1$;
- $\sum_{k=1}^{2n} 2^k$ individual 2-cells.

Moreover, we have the following estimates:

$$\text{Vol}(B_{H_n}(r)) \succeq [w_n(r)]^2 \quad \text{Area}(S_{H_n}(r)) \simeq [w_0(r)]^2 = r^2.$$

Proof. The volume claim is immediate from the volume of the slab. For the area, we first note that each Type I move produces four halves of some $\Delta_{ij}^n(w_0(r))$, as pointed out in Lemma 5.7. There are as many Type I moves involved in creating $S_{H_n}(r)$ as there are copies of $\Theta_i^{n-1}(w_0(r))$ in $S_{G_{n-1}}(r)$, namely 2^{2n} . This gives the correct number of copies of $\Delta_{ij}^n(w_0(r))$ assuming we can show that these half-diagrams join up in pairs to form genuine copies of $\Delta_{ij}^n(w_0(r))$. But this follows immediately from the fact that any two adjacent half-diagrams are adjacent along an edge of the form $(u_{(n-1)1})^r$ lying along the boundary of some original $\Theta_i^{n-1}(w_0(r))$ on $S_{G_{n-1}}(r)$. See Figure 5.9.

On $S_{G_{n-1}}(r)$ there is one strip in the middle of each $\Theta_i^{n-1}(w_0(r))$, each of which is covered by a Type I move. Each Type I move gives rise to four new strips with dimensions $r \times 1$.

Finally we have the original $\sum_{k=1}^{2n-1} 2^k$ individual 2-cells from $S_{G_{n-1}}(r)$, plus one more for each Type I move. Thus the number of individual 2-cells in $S_{H_n}(r)$ is $\sum_{k=1}^{2n} 2^k$.

Adding up the areas of these pieces, we obtain

$$\text{Area}(S_{H_n}(r)) \simeq 2^{2n+1}r^2 + 2^{2n+2}r + \sum_{k=1}^{2n} 2^k \simeq r^2.$$

□

Lower Bounds in G_n . Begin with a slab in $(F_2)^3$ of the form $\varphi^{w_n(r)}(a_{011}) \times \varphi^{w_n(r)}(a_{021}) \times y$, and note that the volume of this slab is $[w_{n+1}(r)]^2$. Assume by induction that a sphere $S_{H_n}(r)$ has been built upon this slab via alternating Type I and Type II moves so that $S_{H_n}(r)$ consists, as shown in the previous lemma, of

- 2^{2n+1} copies of $\Delta_{ij}^n(w_1(r))$;
- 2^{2n+2} strips with dimensions $r \times 1$;
- $\sum_{k=1}^{2n} 2^k$ individual 2-cells.

See Figure 5.2 for confirmation of these hypotheses in the case $n = 1$.

Let $B_{G_n}(r)$ denote the 3-ball obtained by applying a Type I move using u_{ni}^{-r} to each copy of $\Delta_{ij}^n(w_1(r))$, and let $S_{G_n}(r)$ denote its boundary 2-sphere.

Lemma 5.15. *The 2-sphere $S_{G_n}(r)$ consists of*

- 2^{2n+1} copies of $\Theta_i^n(w_0(r))$;
- $\sum_{k=1}^{2n+1} 2^k$ individual 2-cells.

Moreover, we have the following estimates:

$$\text{Vol}(B_{G_n}(r)) \succeq [w_{n+1}(r)]^2 \quad \text{Area}(S_{G_n}(r)) \simeq w_1(r).$$

Proof. The volume claim is immediate from the volume of the slab. For the area, we first note that each Type II move produces four halves of some $\Theta_i^n(w_0(r))$ (minus the equatorial strip), as pointed out in Lemma 5.5. There are as many Type II moves involved in creating $S_{G_n}(r)$ as there are copies of $\Delta_{ij}^{n-1}(w_1(r))$ in $S_{H_n}(r)$, namely 2^{2n+1} . This gives the correct number of copies of $\Theta_i^n(w_0(r))$ assuming we can show that these half-diagrams join up in pairs across strips to form genuine copies of $\Theta_i^n(w_0(r))$. But this follows immediately from the

fact that any two adjacent half-diagrams are adjacent across a strip from some previous Type I move. See Figure 5.8.

Finally we have the original $\sum_{k=1}^{2n} 2^k$ individual 2-cells from $S_{H_n}(r)$, plus one more for each Type II move. Thus the number of individual 2-cells in $S_{G_n}(r)$ is $\sum_{k=1}^{2n+1} 2^k$.

Adding up the area of these pieces, we obtain

$$\text{Area}(S_{G_n}(r)) \simeq 2^{2n+1}w_1(r) + \sum_{k=1}^{2n+1} 2^k \simeq w_1(r).$$

□

Remark 5.16. Note that the constants involved in verifying lower bounds for the \simeq -class of $\delta^{(2)}(x)$ for G_n and H_n grow exponentially as functions of n . This is due to the exponential increase in the number of faces in the spheres created in these groups.

The proof that these balls are embedded in their respective spaces follows exactly as in the base cases, Proposition 5.12, because Type II balls and Type I balls with identical stable groups lie in distinct cosets of the stable group. Thus it remains only to show that the estimates obtained for these particular spheres suffice to establish the lower bounds (cf. Remark 2.4).

Lemma 5.17. *Let $S_{H_n}(r)$ denote the sphere defined above in K_{H_n} with parameter r . For each n , there exist constants A_1, A_2 so that*

$$A_1r^2 \leq \text{Area}(S_{H_n}(r)) \leq A_2r^2.$$

Proof. We use the combinatorial structure of $S_{H_n}(r)$ described in Lemma 5.14. From the proof of Lemma 5.2, we have that $\text{Area}(\Delta_{ij}^n(w_0(r))) = 2r^2$, where this is an actual equality. It follows that

$$\text{Area}(S_{H_n}(r)) = 2^{2n+2}r^2 + 2^{2n+2}r + 2^{2n+1} - 2.$$

It follows (since $r \geq 1$) that we may choose $A_1 = 2^{2n+2}$ and $A_2 = 3 \cdot 2^{2n+2}$. □

Lemma 5.18. *Let $S_{G_n}(r)$ denote the sphere defined above in K_{G_n} with parameter r . For each n , there exists a constant $C > 0$ so that*

$$\text{Area}(S_{G_n}(r + 1)) \leq C \text{Area}(S_{G_n}(r)).$$

Proof. The proof of Lemma 5.4 can be refined to give the estimate

$$w_1(r) \leq \text{Area}(\Theta_i^n(w_0(r))) \leq 3w_1(r).$$

Combining this with the combinatorial structure of S_{G_n} , we then have

$$2^{2n+1}w_1(r) + 2^{2n+2} - 2 \leq \text{Area}(S_{G_n}(r)) \leq 3 \cdot 2^{2n+1}w_1(r) + 2^{2n+2} - 2.$$

From this we compute

$$\begin{aligned} \text{Area}(S_{G_n}(r+1)) &\leq 3 \cdot 2^{2n+1}w_1(r+1) + 2^{2n+2} - 2 \\ &\leq 3 \cdot 2^{2n+2}w_1(r+1) = 3 \cdot 2^{2n+2} \frac{w_1(r+1)}{w_1(r)} w_1(r) \\ &\leq 3 \cdot 2^{2n+2} \frac{w_1(r+1)}{w_1(r)} w_1(r) \\ &\leq 3 \cdot 2 \frac{w_1(r+1)}{w_1(r)} \text{Area}(S_{G_n}(r)). \end{aligned}$$

Note that the ratio $w_1(r+1)/w_1(r)$ is never larger than three ($w_1(k)$ is the length of $\varphi^k(\xi)$, and each application of φ multiplies length by no more than three). It follows that

$$\text{Area}(S_{G_n}(r+1)) \leq 18 \text{Area}(S_{G_n}(r))$$

as required. □

We are finally able to deduce lower bounds for the 2-dimensional Dehn functions of H_n and G_n .

Proposition 5.19 (Lower bounds). *For $n \geq 1$ we have $\delta_{H_n}^{(2)}(x) \succeq \exp^n(\sqrt{x})$ and $\delta_{G_n}^{(2)}(x) \succeq \exp^n(x)$.*

6. UPPER BOUNDS FOR GRAPHS OF GROUPS

In this section we show that the 2-dimensional Dehn functions of the groups G_n and H_n are bounded above by the functions listed in Table 1. In the inductive construction, each group is obtained from the the previous group as a multiple HNN extension. The upper bounds are obtained inductively using two general results about graphs of groups (Propositions 6.1 and 6.2 below). A version of Proposition 6.1 appears in [17]. We give a different proof using admissible maps.

Proposition 6.1. *Let G be the fundamental group of a graph of groups with the following properties:*

- (i) *All the vertex groups are of type \mathcal{F}_3 and their 2-dimensional Dehn functions are bounded above by the superadditive increasing function f .*
- (ii) *All the edge groups are of type \mathcal{F}_2 and their 1-dimensional Dehn functions are bounded above by the superadditive increasing function g .*

Then $\delta_G^{(2)}(x) \preceq (f \circ g)(x)$.

The upper bound obtained in the above proposition may not always be the optimal one. Proposition 6.2 below gives another way to obtain an upper bound. It relies on the existence of a function bounding the area-distortions of the edge groups in G .

Proposition 6.2. *Let G be the fundamental group of a graph of groups with the following properties:*

- (i) *All the vertex groups are of type \mathcal{F}_3 and their 2-dimensional Dehn functions are bounded above by the superadditive increasing function f .*
- (ii) *All the edge groups are of type \mathcal{F}_2 .*
- (iii) *There exists a function h such that if w is a word representing 1 in an edge group Γ , then $\text{Area}_\Gamma(w) \leq h(\text{Area}_G(w))$.*

Then $\delta_G^{(2)}(x) \preceq f(xh(x))$.

Sections 6.1 and 6.2 contain some preliminaries for the proofs of these propositions. The proofs are contained in Sections 6.3-6.5. The propositions are used to prove upper bounds for $\delta_{G_n}^{(2)}(x)$ and $\delta_{H_n}^{(2)}(x)$ in Section 6.6.

6.1. The 3-complex \widetilde{K}_G . We start with the standard total space K_G of the graph of spaces associated with G . Recall that K_G is a quotient map

$$(6.1) \quad q : \sqcup K_v \sqcup (\sqcup K_e \times [0, 1]) \rightarrow K_G,$$

where the K_v are indexed by the vertex set of the graph and the K_e are indexed by its edge set. Note that q identifies each $K_e \times i$ (where

$i = 0, 1$) with its image in a vertex space K_v under a map induced by the edge inclusion $G_e \rightarrow G_v$.

The vertices of \widetilde{K}_G are exactly the vertices of the \widetilde{K}_v . For $1 \leq i \leq 3$, an i -cell of \widetilde{K}_G is either an i -cell of some \widetilde{K}_v or of the form $c \times [0, 1]$, where c is an $(i - 1)$ -cell of some \widetilde{K}_e .

Labeling on \widetilde{K}_G . Each 1-cell of \widetilde{K}_G of the first type mentioned above is labeled by a generator of the corresponding vertex group. All 1-cells of the second type are labeled by the letter e .

The boundary of a 2-cell of the first type is labeled by a relation in the corresponding vertex group. A 2-cell of the second type is homeomorphic to $E \times [0, 1]$, where E is an edge of some \widetilde{K}_e . Then E corresponds a generator, say g , of the edge group G_e . Let G_u and G_v be the vertex groups adjacent to G_e , and let X and Y be words in the generators of G_u and G_v respectively, which represent the images of g under the edge inclusion maps. Then the 2-cell $E \times [0, 1]$ has boundary label $eXe^{-1}Y^{-1}$. Note that any boundary label that contains e is of this form.

Within each piece of \widetilde{K}_G of the form $\widetilde{K}_e \times [0, 1]$, we identify the slice $\widetilde{K}_e \times 1/2$ with \widetilde{K}_e . Every cell $c \times [0, 1]$ in $\widetilde{K}_e \times [0, 1]$ contains an embedded copy $c \times 1/2$ of a cell of K_e . If c is a 1-cell, we label $c \times 1/2$ by the generator of the edge group G_e that labels c . We use this additional cell structure and labeling in the course of the construction below (to define central words of annuli), but we do not think of these as cells of \widetilde{K}_G .

6.2. Transverse maps. In proving upper bounds for 1-dimensional Dehn functions, one often uses the notion of an e -corridor or e -annulus in a van Kampen diagram. To facilitate the definition of an analogous object in the present setting, we require maps $f : S^2 \rightarrow \widetilde{K}_G$ to be *transverse*. Transversality, a condition more stringent than admissibility, gives rise to a *generalized handle decomposition* of S^2 . We summarize a few essential facts here and refer to [6] for more details.

An *index i handle* of dimension n is a product $\Sigma^i \times D^{n-i}$, where Σ^i is a compact, connected i -dimensional manifold with boundary, and D^{n-i} is a closed disk. A *generalized handle decomposition* of an n -dimensional manifold M is a filtration $\emptyset = M^{(-1)} \subset M^{(0)} \subset$

$\dots \subset M^{(n)} = M$ by codimension-zero submanifolds, such that for each i , $M^{(i)}$ is obtained from $M^{(i-1)}$ by attaching finitely many index i handles. Each i -handle $\Sigma^i \times D^{n-i}$ is attached via an embedding $\partial\Sigma^i \times D^{n-i} \rightarrow M^{(i-1)}$. A map f from M to a CW complex X is *transverse* to the cell structure of X if M has a generalized handle decomposition such that the restriction of f to each handle is given by projection onto the second factor, followed by the characteristic map of a cell of X . We say that X is a *transverse CW complex* if the attaching map of every cell is transverse to the cell structure of the skeleton to which it is attached. We will need the following result of Buoncrisiano, Rourke, and Sanderson.

Theorem 6.3 ([9]). *Let M be a compact smooth manifold and $f : M \rightarrow X$ a continuous map into a transverse CW-complex. Suppose $f|_{\partial M}$ is transverse. Then f is homotopic rel ∂M to a transverse map $g : M \rightarrow X$.*

In order to apply this theorem, one needs a transverse complex. The complex K_G defined above can be made transverse by inductively applying the theorem to attaching maps of cells. This procedure can be done in way that preserves the homeomorphism type of the complex and its partition into cells. (See Section 3 of [6] for the details of this procedure.)

Lastly, if one applies the theorem to an admissible map to make it transverse, its combinatorial volume does not change. (This is because the preimages of n -cells do not change during the course of the homotopy, except possibly by shrinking slightly)

Existence of e -annuli. We are concerned with transverse maps $\tau : S^2 \rightarrow \widetilde{K}_G$. Such a map induces a decomposition of S^2 into handles of index 0, 1, and 2, and each i -handle maps to a $(2-i)$ -dimensional cell of \widetilde{K}_G . Each 1-handle inherits the label of the 1-cell that it maps onto. If there is a 1-handle labeled e , there are two possibilities: either the 1-handle is of the form $S^1 \times I$ (an annulus) or it is adjacent to a 0-handle. In the latter case, the 0-handle is mapped homeomorphically to a 2-cell in K_G whose boundary is labeled by a word of the form $eXe^{-1}Y^{-1}$, and is therefore adjacent to exactly one other 1-handle labeled e . This new 1-handle is, in turn, adjacent to another 0-handle. Continuing

this process, we obtain a finite concatenation of alternating 1-handles (labeled e) and 0-handles, which forms an annulus. In either case, we call the annulus an e -annulus. Note that an e -annulus can consist of a single 1-handle of the form $S^1 \times I$.

The only handles adjacent to an e -annulus are 1-handles that are not labeled e and 2-handles. Thus e -annuli never intersect each other. (In particular, such an annulus never intersects itself.)

The central and boundary words of an e -annulus. Any 0-handle in an e -annulus is mapped homeomorphically to a 2-cell of the form $E \times [0, 1]$, where E is an edge in an edge space, say K_e , and $E \times 1/2$ is labeled by a generator of the edge group G_e . Concatenating such labels from successive 0-handles, we obtain the *central word* of the e -annulus. We think of the central word as labeling a *central circle* C that runs through the interior of the annulus. The circle C is simply the union of the segments $[0, 1] \times \{1/2\}$ through the center of each 1-handle of the annulus and the curves $\tau^{-1}(E \times 1/2)$ in each 0-handle. Note that $\tau(C) \subset K_e \times 1/2$.

The boundary of the 2-cell is labeled by $eXe^{-1}Y^{-1}$, where X and Y are words in the generators of the two adjacent vertex groups. Concatenating such words from successive 0-handles in the e -annulus gives the two *boundary words* of the annulus.

6.3. General strategy for the proofs. We assume that \widetilde{K}_G has been made transverse by applying the procedure mentioned after Theorem 6.3. Given any admissible map $\sigma : S^2 \rightarrow \widetilde{K}_G^{(2)}$ with $\text{Area}(\sigma) = x$, we construct an admissible filling $\bar{\sigma} : D^3 \rightarrow \widetilde{K}_G$ whose volume is bounded above by a function \simeq -equivalent to $f(g(x))$ (for Proposition 6.1) or $f(xh(x))$ (for Proposition 6.2).

To construct the admissible filling $\bar{\sigma}$, we think of D^3 as a cone $S^2 \times [0, 1]/S^2 \times 1$ and decompose it into two pieces, $A = S^2 \times [0, 1/2]$ and a closed ball $B = S^2 \times [1/2, 1]/S^2 \times 1$. It follows from Theorem 6.3 that there is a homotopy $\Psi : S^2 \times [0, 1/2] \rightarrow \widetilde{K}_G^{(2)}$ such that $\Psi_0 = \sigma$, and $\Psi_{1/2} = \tau$ is a transverse map with $\text{Area}(\tau) = \text{Area}(\sigma)$. In Section 6.4



FIGURE 6.1. Filling the Admissible map $\sigma : S^2 \rightarrow \widetilde{K}_G$.

we construct an admissible filling $\bar{\tau} : B \rightarrow \widetilde{K}_G$. Then

$$(6.2) \quad \bar{\sigma} = \begin{cases} \Psi & \text{on } A \\ \bar{\tau} & \text{on } B \end{cases}$$

(See Figure 6.1.) Note that $\bar{\sigma}$ is admissible and that $\text{Vol}(\bar{\sigma}) = \text{Vol}(\bar{\tau})$.

To construct the filling $\bar{\tau}$ we decompose the ball B into two families of balls with disjoint interiors. The first family consists of “slabs” homeomorphic to $D^2 \times I$. The lateral boundary $\partial D^2 \times I$ of each slab lies on ∂B . The space formed by deleting the interiors and lateral boundaries (i.e. $D^2 \times (0, 1)$) of the slabs is a disjoint union of balls, which make up the second family. We construct admissible maps on each of the component balls in the two families in such a way that they agree on common boundaries and agree with τ on ∂B .

Under the standard map from \widetilde{K}_G to the associated Bass-Serre tree, the image in \widetilde{K}_G of a slab from the first family above maps to an edge of the tree, while the image of a ball from the second family maps to a vertex of the tree. The volume of a slab is related to the areas of its boundary disks $D^2 \times 0$ and $D^2 \times 1$. The volume of a ball of the second type is controlled using the 2-dimensional Dehn function of the corresponding vertex group. The details of the volume estimates are given in Section 6.5.

6.4. The admissible filling $\bar{\tau}$. An admissible filling of the transverse map $\tau : \partial B \rightarrow \widetilde{K}_G^{(2)}$ is constructed as follows. If ∂B has no e -annuli, the image of τ lies completely in one of the (3-dimensional) vertex spaces K_v . Since f is an upper bound for the 2-dimensional Dehn functions of the vertex groups (in both propositions), there exists an admissible filling $\bar{\tau} : B \rightarrow \widetilde{K}_v$, with $\text{Vol}(\bar{\tau}) \preceq f(\text{Area}(\tau))$.

If ∂B has e -annuli, then $\bar{\tau}$ is constructed inductively. The induction is on the number of e -annuli. If A_1 is an e -annulus, form a new space B_1 , by gluing a slab $D^2 \times [0, 1]$ to ∂B along A_1 , i.e.

$$B_1 = \partial B \sqcup (D^2 \times [0, 1]) / A_1 \sim (\partial D^2 \times [0, 1]).$$

If A_1 consists of a single 1-handle, then it is homeomorphic to $S^1 \times [0, 1]$. With this identification, the restriction of τ to A_1 is simply projection onto the second factor, followed by the characteristic map of a 1-cell. Then τ extends to a map on $D^2 \times [0, 1]$ whose image is the same 1-cell. (The map is constant on each disk $D^2 \times t$.) This defines an admissible map $\tau_1 : B_1 \rightarrow \widetilde{K}_G$

Otherwise let w_1 and C_1 be the central word and central circle of A_1 . Now $\tau|_{C_1}$ is an admissible map into some \widetilde{K}_ϵ . The circle C_1 divides ∂B into two components, and the restriction of τ to the closure of one of these components is a filling of $\tau|_{C_1}$. Thus w_1 represents the identity in G (and hence in the edge group G_ϵ). It follows that w_1 can be filled in \widetilde{K}_ϵ , i.e., there is an admissible map $D^2 \rightarrow \widetilde{K}_\epsilon$ that agrees with $\tau|_{C_1}$ on ∂D^2 . Extend this to a map $\eta : D^2 \times [0, 1] \rightarrow \widetilde{K}_\epsilon \times [0, 1]$ (by defining it to be the identity on the second factor). Restrict the quotient map q from Section 6.1 to $K_\epsilon \times [0, 1]$ and let \tilde{q} denote the unique lift to universal covers so that $\tilde{q} \circ \eta$ agrees with τ on A_1 .

Now define $\tau_1 : B_1 \rightarrow \widetilde{K}_G$ by

$$\tau_1 = \begin{cases} \tau & \text{on } \partial B \\ \tilde{q} \circ \eta & \text{on } D^2 \times I \end{cases}.$$

Note that τ_1 is an admissible map of B_1 into \widetilde{K}_G (i.e. τ_1 restricted to a component of the inverse image of an open 3-cell is a homeomorphism). Moreover, the restrictions of τ_1 to the boundary disks $D^2 \times i$, where $i = 0, 1$, are admissible maps into $\widetilde{K}_G^{(2)}$. (In fact the boundary disks are mapped into the 2-skeletons of vertex spaces.)

The complement $B_1 \setminus D^2 \times (0, 1)$ is a disjoint union of two spheres, which we call the *complementary* spheres. The restrictions of τ_1 to the complementary spheres are admissible maps into $\widetilde{K}_G^{(2)}$. Note that even though τ was a transverse map, we do not require τ_1 to be transverse. The transversality of τ was only used to conclude that the e -annuli on ∂B are all well-defined, embedded, and mutually disjoint. This is clearly the case for the restrictions of τ_1 to each of the complementary spheres. Note that, since the boundary disks $D^2 \times i$ do not contain any 2-cells labeled e , no new e -annuli are added in the course of the above procedure. Now we repeat the procedure with τ replaced by each of the restrictions of τ_1 to the complementary spheres.

Let N be the total number of e -annuli on ∂B . By induction we obtain an admissible map $\tau_N : B_N \rightarrow \widetilde{K}_G$, and complementary spheres S_1, \dots, S_{N+1} (since in each step one of the complementary spheres is divided into two). Let τ_{N_i} denote the restriction of τ_N to S_i . Then τ_{N_i} is an admissible map into the 2-skeleton of some vertex space.

As before, for each i , there is an admissible extension $\bar{\tau}_{N_i}$ of τ_{N_i} defined on D_i^3 , a 3-dimensional ball with boundary S_i , such that $\text{Vol}(\bar{\tau}_{N_i}) \leq f(\text{Area}(\tau_{N_i}))$. Note that the ball $B = \cup_{i=1}^{N+1} D_i^3 \cup B_N$, and define $\bar{\tau} : B \rightarrow \widetilde{K}_G$ by

$$\bar{\tau} = \begin{cases} \tau_N & \text{on } B_N \\ \bar{\tau}_{N_i} & \text{on } D_i^3, \text{ for } 1 \leq i \leq N + 1 \end{cases}$$

Note that $\bar{\tau}$ is an admissible extension of τ . Finally, the admissible filling $\bar{\sigma}$ of the original map σ is defined as in Equation 6.2.

6.5. Filling volumes in graphs of groups. Let A_1, \dots, A_N , w_1, \dots, w_N and $\widetilde{K}_{e_1}, \dots, \widetilde{K}_{e_N}$ denote the e -annuli on ∂B , their central words and the corresponding edge spaces respectively. For any A_i , the restriction of τ_N to the slab that it bounds has volume equal to $\text{Area}_{K_{e_i}}(w_i)$. (If A_i is an e -annulus of the form $S^1 \times [0, 1]$ for some i , then w_i is empty and $\text{Area}_{K_{e_i}}(w_i) = 0$.)

There exists a constant M , which depends only on G , such that if h is a boundary word of one of the e -annuli, say A_i , (so that h is a word in the generators of a vertex group, say G_v), then $\text{Area}_{K_v}(h) \leq M \text{Area}_{K_{e_i}}(w_i)$. This ensures that at the i th step of the procedure,

(i.e., when a slab is glued in along A_i to form B_i), the *total* area of the complementary spheres increases by at most $2M \text{Area}_{K_{e_i}}(w_i)$. Using the superadditivity of f , and the fact that the fillings $\bar{\tau}_{Ni}$ of the complementary spheres were chosen so that $\text{Vol}(\bar{\tau}_{Ni}) \leq f(\text{Area}(\tau)_{Ni})$, we have

$$\begin{aligned}
(6.3) \quad \text{Vol}(\bar{\sigma}) &= \text{Vol}(\bar{\tau}) \\
&= \text{Total volume of slabs} + \text{Total volume of balls} \\
&\leq \sum_{i=1}^N \text{Area}_{K_{e_i}}(w_i) + \sum_{i=1}^{N+1} f(\text{Area}(\tau_{Ni})) \\
&\leq \sum_{i=1}^N \text{Area}_{K_{e_i}}(w_i) + f\left(\sum_{i=1}^{N+1} \text{Area}(\tau_{Ni})\right) \\
&\leq \sum_{i=1}^N \text{Area}_{K_{e_i}}(w_i) + f\left(\text{Area}(\tau) + 2M \sum_{i=1}^N \text{Area}_{K_{e_i}}(w_i)\right)
\end{aligned}$$

We are now ready to complete the proofs of Propositions 6.1 and 6.2. We retain the notation developed above.

Proof of Proposition 6.1. Let $\sigma : S^2 \rightarrow \widetilde{K}_G^{(2)}$ be an admissible map with area x . We obtain an admissible filling $\bar{\sigma} : D^3 \rightarrow \widetilde{K}_G$ as described in the procedure above. Equation (6.3) estimates the volume of this filling. By the assumption on the 1-dimensional Dehn functions of the edge groups, we have that $\text{Area}_{K_{e_i}}(w_{A_i}) \leq g(|w_{A_i}|)$. So by the superadditivity of g , we have $\sum \text{Area}_{K_{e_i}}(w_{A_i}) \leq g(\sum |w_{A_i}|) \leq g(x)$, since the total lengths of central words of annuli cannot be more than the total area of σ . So the estimate for the volume is $\text{Vol}(\bar{\sigma}) \leq g(x) + f(x + 2Mg(x))$. This gives

$$\delta_G^{(2)}(x) \preceq g(x) + f(x + 2Mg(x)).$$

However, since $x \preceq f(x)$, and $x \preceq g(x)$, we have $g(x) + f(x + 2Mg(x)) \preceq f(g(x))$. This is the required upper bound. \square

Proof of Proposition 6.2. As in the previous proof, let $\sigma : S^2 \rightarrow \widetilde{K}_G^{(2)}$ be an admissible map with area x and let $\bar{\sigma}$ be the admissible filling from the procedure above. This time, we have the condition

$\text{Area}_{K_{e_i}}(w_{A_i}) \leq h(\text{Area}_{K_G}(w_{A_i}))$ for each i . The restriction of τ to a ‘‘hemispherical’’ piece of ∂B gives a filling of w_{A_i} of size at most x . Since there are at most x annuli, the estimate in Equation 6.3 becomes

$$\begin{aligned} \text{Vol}(\bar{\sigma}) &\leq \sum_{i=1}^N \text{Area}_{K_{e_i}}(w_{A_i}) + f\left(x + 2M \sum_{i=1}^N \text{Area}_{K_{e_i}}(w_{A_i})\right) \\ &\leq xh(x) + f(x + 2Mxh(x)). \end{aligned}$$

This give $\delta_G^{(2)}(x) \preceq f(xh(x))$. \square

6.6. Upper bounds for the super-exponential examples. The proof of the upper bounds for the super-exponential examples is by induction. We have an alternating sequence of groups

$$H_0 < G_0 < H_1 < G_1 < H_2 < G_2 \cdots$$

where $H_0 = F_2 \times F_2$. (Note that this is different from the H_0 defined in Table 1. In Section 4, example 4.4, the group G_0 was obtained from $F_2 \times F_2 \times F_2$ by coning over $F_2 \times F_2$. This can also be viewed as a multiple (4-fold) HNN extension with base $F_2 \times F_2 = H_0$, where two stable letters act via the identity and two act via $\varphi \times \varphi$.)

All of the above groups are of type \mathcal{F}_3 . Apart from H_0 , they all have 3-dimensional $K(\pi, 1)$'s. As described in Section 4.1, each group in the sequence is a multiple HNN extension of the previous one.

Proof of upper bounds for $\delta_{H_n}^{(2)}$ and $\delta_{G_n}^{(2)}$. Since $F_2 \times F_2$ has a 2-dimensional $K(\pi, 1)$ we have $\delta_{H_0}^{(2)}(x) = \delta_{F_2 \times F_2}^{(2)}(x) \simeq x$. This starts the induction.

Step 1. Deducing $\delta_{G_n}^{(2)}(x)$ upper bounds from $\delta_{H_n}^{(2)}(x)$ upper bounds ($n \geq 0$). This is a straightforward application of Proposition 6.1. The group G_n is the fundamental group of a graph of groups where the underlying graph is a bouquet of two circles (four if $n = 0$), the vertex group is H_n and the edge groups are $F_{2^{n+1}} \times F_2$. Further,

- (i) $\delta_{H_0}^{(2)}(x) \simeq x$ (base case) and $\delta_{H_n}^{(2)}(x) \simeq \exp^n(\sqrt{x})$ for $n > 0$ (induction hypothesis).
- (ii) $\delta_{F_{2^{n+1}} \times F_2}^{(1)}(x) \simeq x^2$.

Proposition 6.1 now implies that $\delta_{G_0}^{(2)}(x) \preceq x^2$ and $\delta_{G_n}^{(2)}(x) \preceq \exp^n(x)$ for $n > 0$.

Step 2. Deducing $\delta_{H_n}^{(2)}(x)$ upper bounds from $\delta_{G_{n-1}}^{(2)}(x)$ upper bounds ($n > 0$). This is more involved than the previous step. The group H_n is the fundamental group of a graph of groups where the underlying graph is a wedge of 2^{n+1} circles, the vertex group is G_{n-1} and the edge groups are all isomorphic to $F_2 \rtimes F_4$. The result will follow from Proposition 6.2 together with the following lemma, which describes how areas in the edge groups get distorted in H_n .

Lemma 6.4 (Area-distortion of Γ in H_n). *Let Γ be an edge group in the graph of groups description of H_n . Then there exists a constant β_n , which depends only on n , such that for any word w in the generators of Γ that represents 1,*

$$(6.4) \quad \text{Area}_\Gamma(w) \leq (\beta_n \text{Area}_{H_n}(w) e^{\sqrt{\beta_n \text{Area}_{H_n}(w)}})^2.$$

Section 7 is devoted to the proof of this lemma. We now have:

- (i) $\delta_{G_0}^{(2)}(x) \preceq x^2$ (Step 1) and $\delta_{G_{n-1}}^{(2)}(x) \preceq \exp^{n-1}(x)$ for $n > 1$ (induction hypothesis)
- (ii) By Lemma 6.4, the function $h(x) = (\beta_n x e^{\sqrt{\beta_n x}})^2$ satisfies the third condition in the statement of Proposition 6.2.

Proposition 6.2 and the fact that $h(x) \simeq e^{\sqrt{x}}$ now imply that $\delta_{H_1}^{(2)}(x) \preceq (x e^{\sqrt{x}})^2$ and $\delta_{H_n}^{(2)}(x) \preceq \exp^{n-1}(x e^{\sqrt{x}})$ for $n > 1$. Since $x e^{\sqrt{x}} \simeq e^{\sqrt{x}}$ and $(e^{\sqrt{x}})^2 \simeq e^{\sqrt{x}}$, we have $\delta_{H_n}^{(2)}(x) \preceq \exp^n(\sqrt{x})$ for all $n \geq 1$.

This completes the proof of the upper bounds for the super-exponential examples. \square

7. PROOF OF THE AREA-DISTORTION LEMMA

This section provides a detailed proof of Lemma 6.4. Let Γ be any of the 2^n groups isomorphic to $F_2 \rtimes_\theta F_4$ listed in the row corresponding to H_n in Table 1. In this section we will use the following notation for Γ :

$$\Gamma = \langle \mathbf{a} \rangle \rtimes_\theta \langle \mathbf{u}, \mathbf{b} \rangle$$

Here $\mathbf{u} = \mathbf{u}_{n-1}$. (Note that each of the 2^n edge groups has $\langle \mathbf{u}_{n-1} \rangle$ as a subgroup.) Either $\mathbf{a} = \mathbf{a}_{(n-1)i}$ with $\mathbf{b} = \mathcal{L}(i)$ or $\mathbf{a} = \mathbf{u}_{n-2}^{-1} \mathbf{a}_{(n-1)i}$ with $\mathbf{b} = \mathcal{L}(i + 2^{n-1})$ for some $1 \leq i \leq 2^{n-1}$. The F_4 generated by \mathbf{u} and \mathbf{b} acts on $\langle \mathbf{a} \rangle$ via θ , as defined in Definition 4.5.

As we are concerned only with areas in this section, we work with van Kampen diagrams, rather than admissible or transverse maps, and combinatorial complexes rather than transverse ones. Since we are working with fixed presentations, we will use the phrase “van Kampen diagram over G ” to mean “van Kampen diagram over the fixed presentation for G .”

7.1. Strategy. Let w be a word in the generators of Γ that represents 1. Then there exists a van Kampen diagram D for w over H_n , which is area-minimizing, i.e. $\text{Area}(D) = \text{Area}_{H_n}(w)$. First realize D as a union of \mathbf{u} -corridors and complementary regions. Next, use this structure to produce a van Kampen diagram Δ for w over Γ , which has the same combinatorial decomposition into \mathbf{u} -corridors and complementary regions. Finally show that $\text{Area}(\Delta)$, and hence $\text{Area}_\Gamma(w)$, is bounded above by the quantity on the right hand side of inequality (6.4).

Remark 7.1. For the rest of this section, we assume that the van Kampen diagram D has the property that every edge on the boundary of D belongs to a 2-cell in D . We can restrict to this case using the superadditivity of the function $(Cx3^{\sqrt{Cx}})^2$.

Remark 7.2. Throughout this proof we abuse notation and use bold-faced letters to denote either a pair of generators for a free group or a single one of these generators; it will be clear from the context which of these we mean. For example, “ \mathbf{u} -corridor” is used to mean $u_{(n-1)i}$ -corridor, where i is 1 or 2, as we do not need to distinguish between these. When we refer to, say, “the word $\mathbf{u}x\mathbf{u}^{-1}$ ” it is understood that both instances of \mathbf{u} in the word refer to the same generator (either $u_{(n-1)1}$ or $u_{(n-1)2}$).

7.2. Geometry of \mathbf{u} -corridors. The following is a complete list of the relations involving the generators $\mathbf{u} = \mathbf{u}_{n-1}$ in the presentation for H_n .

- (i) Let $E = \langle \mathbf{a}_{(n-1)1}, \dots, \mathbf{a}_{(n-1)2^{n-1}}, \mathbf{u}_{n-2} \rangle \simeq F_{2^n} \times F_2$. Recall that G_{n-1} is the cone of H_{n-1} over E with stable letters \mathbf{u} and relations

$$\mathbf{u}g\mathbf{u}^{-1}(\varphi(g))^{-1} = 1 \quad \text{where } g = \text{any generator for } E.$$

- (ii) Recall that H_n is a multiple HNN-extension of G_{n-1} with stable letters \mathbf{a}_{nj} , with $1 \leq j \leq 2^n$. The new relations involving $\mathbf{u}_{(n-1)}$ are the commuting relations:

$$\mathbf{u} \mathbf{a}_{nj} \mathbf{u}^{-1} \mathbf{a}_{nj}^{-1} = 1 \quad \text{for } 1 \leq j \leq 2^n.$$

Since these are all the relations involving the \mathbf{u} , it makes sense to talk about \mathbf{u} -corridors and \mathbf{u} -annuli in van Kampen diagrams over H_n . The reader may refer to Section 7.2 of [7] for the definitions and properties of corridors and annuli (called rings in [7]). By the assumption on D in Remark 7.1, every edge labeled \mathbf{u} in ∂D is part of a non-trivial \mathbf{u} -corridor in D . Although D may contain \mathbf{u} -annuli, only \mathbf{u} -corridors will play a key role in the argument below.

Area-length inequality for \mathbf{u} -corridors in D and Δ . The boundary of a \mathbf{u} -corridor \mathcal{C} over H_n or Γ is labeled by a word of the form $\mathbf{u}X_1\mathbf{u}^{-1}X_2$. We call the X_i the *horizontal boundary words* of \mathcal{C} . From the presentations of H_n and Γ , we see that a horizontal boundary of a single 2-cell involving \mathbf{u} has length at most 3. (For Γ we measure lengths in the intrinsic metric, which may differ from the inherited one by a factor of 2.) Thus

$$(7.1) \quad \frac{|X_i|}{3} \leq \text{Area}(\mathcal{C}) \leq |X_i|, \quad \text{for } i = 1, 2.$$

Exponential distortion of corridors. The following lemma produces a \mathbf{u} -corridor over Γ corresponding to a given \mathbf{u} -corridor in D , and relates their lengths.

Lemma 7.3 (Exponential distortion of corridors). *Let \mathcal{C}_D be a \mathbf{u} -corridor in D with boundary label $\mathbf{u}X_1\mathbf{u}^{-1}X_2$. Then there exists a \mathbf{u} -corridor \mathcal{C}_Γ over Γ , with boundary label $\mathbf{u}Y_1\mathbf{u}^{-1}Y_2$ such that the Y_i are words in \mathbf{a} with $Y_i =_{H_n} X_i$ for $i = 1, 2$. Furthermore,*

$$(7.2) \quad |Y_i| \leq 3(3^{|X_i|}).$$

Proof. Let $\Gamma' = \langle \mathbf{a}' \rangle \rtimes \langle \mathbf{u}, \mathbf{b}' \rangle$ be a group from the list of edge groups for H_n (possibly different from Γ), and let E be as defined in the beginning of subsection 7.2. The following two properties will be used repeatedly to construct \mathcal{C}_Γ .

Property 1. $E \cap \Gamma' = \langle \mathbf{a}' \rangle$

Proof. It is easy to see (using the normal form for semidirect products) that this is equivalent to the statement $E \cap \langle \mathbf{u}, \mathbf{b}' \rangle = 1$. The latter statement follows easily from an elementary argument using \mathbf{u} -corridors. \square

Property 2. $\langle \mathbf{a}' \rangle$ is a retract of E .

Proof. If $\langle \mathbf{a}' \rangle$ is generated by $\mathbf{a}_{(n-1)i}$ for some i , then the retraction is simply the projection of E onto $\langle \mathbf{a}' \rangle$. If it is a subgroup generated by $\mathbf{u}_{n-1}^{-1} \mathbf{a}_{(n-1)i}$ for some i , then observe that the following map is a retraction:

$$\mathbf{a}_{(n-1)j} \mapsto 1 \ (j \neq i); \quad \mathbf{a}_{(n-1)i} \mapsto \mathbf{u}_{n-1}^{-1} \mathbf{a}_{(n-1)i}; \quad \mathbf{u}_{n-1} \mapsto 1.$$

\square

Since the horizontal boundary X_1 of \mathcal{C}_D has endpoints on the boundary of D , there is a subword W of w such that $W^{-1}X_1$ represents 1. By Britton's Lemma applied to the multiple HNN extension description of H_n in Table 1, we conclude that X_1 must have an innermost subword of the form $\mathbf{a}_{ni}v\mathbf{a}_{ni}^{-1}$ or $\mathbf{a}_{ni}^{-1}v\mathbf{a}_{ni}$ for some i . Here v is a word in the generators of E that represents an element of the edge group corresponding to \mathbf{a}_{ni} , say $\Gamma' = \langle \mathbf{a}' \rangle \rtimes \langle \mathbf{u}, \mathbf{b}' \rangle$. By Property 1, there exists a (reduced) word v' in the generators \mathbf{a}' representing the same element as v . By Property 2 we have $|v'| \leq |v|$ (since retracts are length-non-increasing). Then if the innermost subword is of the form $\mathbf{a}_{ni}v\mathbf{a}_{ni}^{-1}$, it can be replaced with $\varphi(v')$, and if it is of the form $\mathbf{a}_{ni}^{-1}v\mathbf{a}_{ni}$, it can be replaced with $\varphi^{-1}(v')$. Both $\varphi(v')$ and $\varphi^{-1}(v')$ have length at most $3|v|$.

Repeat this procedure (at most $|X_1|/2$ times) until all instances of \mathbf{a}_{ni} have been eliminated. We obtain a word X'_1 in the generators of E of length $|X'_1| \leq 3^{|X_1|/2} \leq 3^{|X_1|}$.

Since X'_1 represents the same word as W , which is a word in Γ , Properties 1 and 2 again apply to produce a word Y_1 in the generators \mathbf{a} , representing the same group element as X'_1 . Furthermore, $|Y_1| \leq |X'_1| \leq 3^{|X_1|}$.

Let Y_2 be the word $\varphi(Y_1)$ (where φ simply acts individually on each generator). Then $\mathbf{u}Y_1\mathbf{u}^{-1}Y_2$ is the boundary of a corridor (i.e. a van Kampen diagram consisting of a single corridor) over Γ , which we call

\mathcal{C}_Γ . Since $|Y_2| \leq 3|Y_1|$ and $|X_1| \leq 3|X_2|$, we have $|Y_2| \leq 3(3^{3|X_2|})$. Clearly X_1 and Y_1 also satisfy this inequality. \square

7.3. Construction of Δ . A van Kampen diagram Δ for w over Γ is obtained from D by the following sequence of moves.

- (i) Remove from D all of the open cells of $D \setminus \partial D$ except for open 1-cells labeled \mathbf{u} and open 2-cells whose closures have an edge labeled \mathbf{u} . The result is a circle labeled w , with a finite collection of “open” \mathbf{u} -corridors attached. Each open \mathbf{u} -corridor is topologically $[0, 1] \times (0, 1)$, with $0 \times (0, 1)$ and $1 \times (0, 1)$ identified with open 1-cells in the circle. Complete this to get a band complex, i.e. a circle with a collection of closed \mathbf{u} -corridor bands attached. The \mathbf{u} -corridor structure on the closed bands is obtained by pulling back the cell structure and labeling from D as in Section 7.2 of [7].
- (ii) Replace each \mathbf{u} -corridor in this band complex with the corresponding corridor over Γ , guaranteed by Lemma 7.3. The result is another band complex, which we denote by B .
- (iii) Remove the open 1-cells labeled \mathbf{u} and the open 2-cells of B to obtain a disjoint union of circles. The labels of these circles are called *complementary words*.

The complementary words w_i are words in the generators \mathbf{a} and \mathbf{b} that represent the trivial element of Γ . Let Δ_i be an area-minimizing van Kampen diagram over Γ for w_i .

- (iv) Define

$$\Delta = (B \sqcup (\sqcup_i \Delta_i)) / \sim$$

where \sim identifies the loop corresponding to w_i in B with $\partial\Delta_i$ for each i .

Note that if there are no \mathbf{u} -corridors in D , the band complex B is just a circle, and there is just one complementary word $w_1 = w$.

7.4. Upper bound for $\text{Area}(\Delta)$. The area of Δ is simply the sum of the areas of the \mathbf{u} -corridors and the areas of the Δ_i .

We first obtain an upper bound on the total area contribution of the \mathbf{u} -corridors. Define

$$L = \begin{cases} \max \{|v| \mid v \text{ a horizontal boundary word of a } \mathbf{u}\text{-corridor in } \Delta\} \\ 1 \text{ if there are no } \mathbf{u}\text{-corridors} \end{cases}$$

It follows from inequality (7.1) that $\text{Area}(\mathcal{C}) \leq L$ for any \mathbf{u} -corridor \mathcal{C} . Since each \mathbf{u} -corridor intersects the boundary of Δ in two edges labeled \mathbf{u} , there are at most $|w|/2$ such corridors. Thus

$$(7.3) \quad \text{total area of the } \mathbf{u}\text{-corridors} \leq \frac{|w|L}{2}.$$

Since $\langle \mathbf{a}, \mathbf{b} \rangle < \Gamma$ is isomorphic to $F_2 \times F_2$, and the complementary words w_i are words in \mathbf{a} and \mathbf{b} , we have

$$(7.4) \quad \sum_i \text{Area}(w_i) \leq \sum_i (|w_i|)^2 \leq \left(\sum_i |w_i| \right)^2$$

From the definition of the w_i , we have:

$$(7.5) \quad \sum_i |w_i| \leq 2(\#\{\mathbf{u}\text{-corridors}\})L + (|w| - 2\#\{\mathbf{u}\text{-corridors}\}) \leq |w|L,$$

Putting together the inequalities (7.3), (7.4) and (7.5), we have

$$\text{Area}(\Delta) \leq \frac{|w|L}{2} + (|w|L)^2 \leq (2|w|L)^2.$$

The proof of Lemma 6.4 will now follow easily from the above estimate, together with the following two facts. (Just take $\beta_n = 20\beta$, where β is the constant from Fact 2, and recall that $\text{Area}(D) = \text{Area}_{H_n}(w)$.)

Fact 1. $|w| \leq 10\text{Area}(D)$.

Proof. The assumption that D is a topological disk implies that each edge of ∂D is part of the boundary of a 2-cell in D . Since the maximum length of a relation in H_n is 10, the area of D is at least $\frac{|w|}{10}$. \square

Fact 2. There exists a constant β , independent of w , such that

$$(7.6) \quad L \leq \beta 3\sqrt{\beta \text{Area}(D)}.$$

Proof. There are two cases, depending on the relative sizes of $|w|$ and L .

Case (i): $|w| > (\log_3 L)^2$.

By Fact 1 we have $L < 3\sqrt{|w|} \leq 3\sqrt{10\text{Area}(D)}$, and so inequality (7.6) holds with $\beta \geq 10$.

Case (ii): $|w| \leq (\log_3 L)^2$.

In this case, we establish inequality (7.6) by obtaining a lower bound on $\text{Area}(D)$. More precisely, we show that for sufficiently large L , $\text{Area}(D) \geq (\log_3 L)^2/144$. Recall that L is the maximal length of a horizontal boundary word of a \mathbf{u} -corridor in Δ . The idea of the proof is as follows: since $|\partial\Delta| = |w|$ is relatively small compared to L , the existence of a \mathbf{u} -corridor of length L forces Δ to have a large number of long \mathbf{u} -corridors. This implies that D also has a large number (order $\log_3 L$) of long (length order $\log_3 L$) \mathbf{u} -corridors in D . The areas of these corridors of D add up to the required lower bound on $\text{Area}(D)$.

We first introduce the notion of the level of a \mathbf{u} -corridor in order to compare lengths of corridors and obtain the above estimates.

Levels for \mathbf{u} -corridors. We first define levels of \mathbf{u} -corridors in Δ . Note that Γ can be expressed as a multiple HNN extension with base group $F_2 \times F_2 = \langle \mathbf{a} \rangle \times \langle \mathbf{b} \rangle$, edge group $\langle \mathbf{a} \rangle$, and stable letters \mathbf{u} . Let T denote the corresponding Bass-Serre tree. There is a natural equivariant map from the Cayley 2-complex of Γ to T . Let $\eta : \Delta \rightarrow T$ be the composition of this map with a standard combinatorial map from Δ into the Cayley complex. Note that η takes 1-cells labeled \mathbf{u} in Δ to 1-cells of T and maps all other 1-cells to vertices. Thus η maps a \mathbf{u} -corridor of Δ to an edge of T .

Choose a base corridor \mathcal{C}_0 of Δ that has a horizontal boundary word of maximal length L . Then $\eta(\mathcal{C}_0)$ is an edge of T . Use the distance function d_T on T (assume each edge of T has length 1) to define a Morse function

$$h : T \rightarrow \mathbb{R} : x \mapsto d_T(x, \eta(\mathcal{C}_0)).$$

Now we say that a vertex v in T is at *level* $h(v)$ and an edge $[v, w]$ of T is at *level* $\max\{h(v), h(w)\}$. We say that a subset of Δ is at *level* i if its image under η is either a vertex or an edge at level i . (Examples

of such subsets will be \mathbf{u} -corridors and horizontal boundaries of \mathbf{u} -corridors.) Finally, we define the *level* of a \mathbf{u} -corridor in D to be the level of the corresponding \mathbf{u} -corridor in Δ .

By the above definitions, the edge $\eta(\mathcal{C}_0)$ of T is at level 0, so that the corridor \mathcal{C}_0 of Δ is at level 0 as well. Note that there may be other \mathbf{u} -corridors at level 0, but these can be ignored: the presence of \mathcal{C}_0 is already sufficient to guarantee the existence of enough levels of long corridors.

Iterated scaling inequality. Let L_i denote the number of horizontal edges of Δ at level i that are part of the boundary of a \mathbf{u} -corridor at level i . We also call this the *total \mathbf{u} -corridor length* at level i . Note that $L \leq L_0$ and that

$$\frac{L_i}{3} \leq \text{total area of } \mathbf{u}\text{-corridors at level } i \leq L_i.$$

Let B_i denote the number of edges at level i that are part of $\partial\Delta$.

Since conjugation by $\mathbf{u}^{\pm 1}$ scales \mathbf{a} -words by at most a factor of 3, we have $L_{i-1} \leq 3L_i + B_{i-1}$. By iterating this we obtain:

$$L \leq L_0 \leq 3^i L_i + \sum_{j=0}^{i-1} 3^j B_j, \quad \text{for } i > 0.$$

Since $\sum_{j=0}^{i-1} B_j \leq |w|$ for each i , this implies

$$(7.7) \quad L \leq 3^i L_i + 3^i |w|.$$

Large L implies many long levels. Now we show that if L is sufficiently large, then there are at least $\frac{\log_3 L}{4}$ levels in Δ , each with total \mathbf{u} -corridor length at least \sqrt{L} .

Since $\lim_{L \rightarrow \infty} \frac{(\log_3 L)^2}{\sqrt{L}} = 0$, there exists $P > 0$ such that $(\log_3 L)^2 \leq \sqrt{L}$ for all $L \geq P$. Note that P depends only on the functions $(\log_3 x)^2$ and \sqrt{x} and not on w . For $L \geq P$, inequality (7.7) and the base inequality of Case (ii) give

$$L \leq 3^i L_i + 3^i |w| \leq 3^i L_i + 3^i (\log_3 L)^2 \leq 3^i L_i + 3^i \sqrt{L}.$$

Rearranging gives $L_i \geq 3^{-i} L - \sqrt{L}$. The reader can now verify that if $i \leq \frac{\log_3 L}{4}$ and $L \geq 16$, then $3^{-i} L \geq 2\sqrt{L}$. As a consequence, we

have:

$$L_i \geq \sqrt{L}.$$

In summary, if $L \geq \max\{P, 16\}$, then there are at least $\frac{\log_3 L}{4}$ levels in Δ , each with total \mathbf{u} -corridor length at least \sqrt{L} .

Relating $\text{Area}(D)$ and $(\log_3 L)^2$ for large L . Now we can estimate the area of D from below using \mathbf{u} -corridors.

$$\begin{aligned} \text{Area}(D) &\geq \text{total area of } u\text{-corridors in } D \\ &\geq \text{total area of } u\text{-corridors in } D \text{ at or below level } \frac{\log_3 L}{4} \\ &\geq \left(\frac{\log_3 L}{4}\right) \left(\frac{\log_3 \sqrt{L}/3}{9}\right) \\ &= \frac{(\log_3 L)^2}{72} - \frac{\log_3 L}{36} \\ &\geq \frac{(\log_3 L)^2}{144} \quad \text{provided } L \geq 3^4. \end{aligned}$$

The second term in the third inequality above is obtained as follows. Note that \sqrt{L} is a lower bound for the total \mathbf{u} -corridor length at level i in Δ , and hence is a lower bound for a sum $\sum |Y_j|$, where j runs over an index set for all \mathbf{u} -corridors at level i , and the Y_j are horizontal boundary words of these corridors. Thus, by inequality (7.2),

$$\sqrt{L} \leq \sum |Y_j| \leq \sum 3(3^{3|X_j|}) \leq 3(3^{3\sum |X_j|}),$$

and so the total \mathbf{u} -corridor length at level i in D (which is $\sum |X_j|$ in the inequality above) is at least $\log_3(\sqrt{L}/3)/3$. Finally, by inequality (7.1), the total area of \mathbf{u} -corridors at level i in D is at least $\log_3(\sqrt{L}/3)/9$.

Summary. We have shown that in Case (i), we have $L \leq 3\sqrt{10\text{Area}(D)}$, and in Case (ii), we have $L \leq 3\sqrt{144\text{Area}(D)}$, provided $L \geq \max\{3^4, 16, P\}$. Thus inequality (7.6) holds for all L , provided we take $\beta = \max\{10, 144, 3^4, 16, P\} = \max\{144, P\}$. Since P was independent of w , so is β . \square

Remark 7.4. The notion of area distortion as a group invariant is defined in Section 2 of [12]. Lemma 6.4 provides an upper bound for the area distortion of Γ in H_n . The reader can verify that the

boundary words for the van Kampen diagrams $\Theta_i^n(N)$ from Section 5 establish the lower bounds for this distortion. Thus the distortion of Γ in H_n is $f(x) \simeq e^{\sqrt{x}}$. It would be interesting to find other pairs (G, H) of type $(\mathcal{F}_3, \mathcal{F}_2)$ where area distortion can be explicitly computed. For example, the subgroup of the group H_n which is generated by $\{\mathbf{a}_{ki} \mid k \leq n, i = 1, 2\}$ should have area distortion $\exp^n \sqrt{x}$.

REFERENCES

- [1] J. M. ALONSO, W. A. BOGLEY, R. M. BURTON, S. J. PRIDE, AND X. WANG, *Second order Dehn functions of groups*, Quart. J. Math. Oxford Ser. (2), 49 (1998), pp. 1–30.
- [2] J. M. ALONSO, X. WANG, AND S. J. PRIDE, *Higher-dimensional isoperimetric (or Dehn) functions of groups*, J. Group Theory, 2 (1999), pp. 81–112.
- [3] J.-C. BIRGET, A. Y. OL'SHANSKII, E. RIPS, AND M. V. SAPIR, *Isoperimetric functions of groups and computational complexity of the word problem*, Ann. of Math. (2), 156 (2002), pp. 467–518.
- [4] N. BRADY, M. BRIDSON, M. FORESTER, AND K. SHANKAR, *Snowflake groups, Perron-Frobenius eigenvalues, and isoperimetric spectra*, Geom. Topol., 13 (2009), pp. 141–187.
- [5] N. BRADY AND M. R. BRIDSON, *There is only one gap in the isoperimetric spectrum*, Geom. Funct. Anal., 10 (2000), pp. 1053–1070.
- [6] N. BRADY AND M. FORESTER, *Density of isoperimetric spectra*. arXiv:0812.1036, 2008.
- [7] M. R. BRIDSON, *The geometry of the word problem*, in *Invitations to geometry and topology*, vol. 7 of Oxf. Grad. Texts Math., Oxford Univ. Press, Oxford, 2002, pp. 29–91.
- [8] ———, *Polynomial Dehn functions and the length of asynchronously automatic structures*, Proc. London Math. Soc. (3), 85 (2002), pp. 441–466.
- [9] S. BUONCRISTIANO, C. P. ROURKE, AND B. J. SANDERSON, *A geometric approach to homology theory*, Cambridge University Press, Cambridge, 1976. London Mathematical Society Lecture Note Series, No. 18.
- [10] M. DEHN, *Über unendliche diskontinuierliche Gruppen*, Math. Ann., 71 (1911), pp. 116–144.
- [11] ———, *Transformation der Kurven auf zweiseitigen Flächen*, Math. Ann., 72 (1912), pp. 413–421.
- [12] S. M. GERSTEN, *Preservation and distortion of area in finitely presented groups*, Geom. Funct. Anal., 6 (1996), pp. 301–345.
- [13] M. GROMOV, *Hyperbolic groups*, in *Essays in group theory*, vol. 8 of Math. Sci. Res. Inst. Publ., Springer, New York, 1987, pp. 75–263.

- [14] ———, *Asymptotic invariants of infinite groups*, in Geometric group theory, Vol. 2 (Sussex, 1991), vol. 182 of London Math. Soc. Lecture Note Ser., Cambridge Univ. Press, Cambridge, 1993, pp. 1–295.
- [15] P. PAPASOGLU, *Isodiametric and isoperimetric inequalities for complexes and groups*, J. London Math. Soc. (2), 62 (2000), pp. 97–106.
- [16] M. V. SAPIR, J.-C. BIRGET, AND E. RIPS, *Isoperimetric and isodiametric functions of groups*, Ann. of Math. (2), 156 (2002), pp. 345–466.
- [17] X. WANG AND S. J. PRIDE, *Second order Dehn functions and HNN-extensions*, J. Austral. Math. Soc. Ser. A, 67 (1999), pp. 272–288.
- [18] R. YOUNG, *A note on higher order filling functions*. arXiv:0805.0584, to appear.

DEPT. OF MATHEMATICS & STATISTICS, UNIVERSITY OF SOUTH ALABAMA,
MOBILE, AL 36688

E-mail address: `jbarnard@jaguar1.usouthal.edu`

DEPT. OF MATHEMATICS, UNIVERSITY OF OKLAHOMA, NORMAN, OK 73019

E-mail address: `nbrady@math.ou.edu`

DEPT. OF MATHEMATICS, LOUISIANA STATE UNIVERSITY, BATON ROUGE,
LA 70803

E-mail address: `pdani@math.lsu.edu`