

# Super-Teichmüller spaces and related structures

Anton M. Zeitlin

Louisiana State University, Department of Mathematics

Columbia University

New York

February 1, 2019

Outline

Introduction

Cast of characters

Coordinates on  
Super-Teichmüller  
space

$\mathcal{N} = 2$   
Super-Teichmüller  
theory

Further work

Open problems



Introduction

Cast of characters

Coordinates on Super-Teichmüller space

$\mathcal{N} = 2$  Super-Teichmüller theory

Further work

Open problems

## Outline

Introduction

Cast of characters


Coordinates on  
Super-Teichmüller  
space

$\mathcal{N} = 2$   
Super-Teichmüller  
theory

Further work

Open problems

Let  $F_s^g \equiv F$  be the Riemann surface of genus  $g$  and  $s$  punctures.  
We assume  $s > 0$  and  $2 - 2g - s < 0$ .



Teichmüller space  $T(F)$  has many incarnations:

- ▶ {complex structures on  $F$ }/isotopy
- ▶ {conformal structures on  $F$ }/isotopy
- ▶ {hyperbolic structures on  $F$ }/isotopy

Isotopy here stands for diffeomorphisms isotopic to identity.

Outline

**Introduction**

Cast of characters


Coordinates on  
Super-Teichmüller  
space

$\mathcal{N} = 2$   
Super-Teichmüller  
theory

Further work

Open problems

Let  $F_s^g \equiv F$  be the Riemann surface of genus  $g$  and  $s$  punctures.  
We assume  $s > 0$  and  $2 - 2g - s < 0$ .



Teichmüller space  $T(F)$  has many incarnations:

- ▶ {complex structures on  $F$ }/isotopy
- ▶ {conformal structures on  $F$ }/isotopy
- ▶ {hyperbolic structures on  $F$ }/isotopy

Isotopy here stands for diffeomorphisms isotopic to identity.

Outline

**Introduction**

Cast of characters


Coordinates on  
Super-Teichmüller  
space

$\mathcal{N} = 2$   
Super-Teichmüller  
theory

Further work

Open problems

Let  $F_s^g \equiv F$  be the Riemann surface of genus  $g$  and  $s$  punctures.  
We assume  $s > 0$  and  $2 - 2g - s < 0$ .



Teichmüller space  $T(F)$  has many incarnations:

- ▶  $\{\text{complex structures on } F\}/\text{isotopy}$
- ▶  $\{\text{conformal structures on } F\}/\text{isotopy}$
- ▶  $\{\text{hyperbolic structures on } F\}/\text{isotopy}$

Isotopy here stands for diffeomorphisms isotopic to identity.

[Outline](#)[Introduction](#)[Cast of characters](#)[Coordinates on  
Super-Teichmüller  
space](#) [\$\mathcal{N} = 2\$   
Super-Teichmüller  
theory](#)[Further work](#)[Open problems](#)

Representation-theoretic definition:

$$T(F) = \text{Hom}'(\pi_1(F), PSL(2, \mathbb{R})) / PSL(2, \mathbb{R}),$$

where  $\rho \in \text{Hom}'$  if

- ▶  $\rho$  is injective
- ▶ identity in  $PSL(2, \mathbb{R})$  is not an accumulation point of the image of  $\rho$ , i.e.  $\rho$  is discrete
- ▶ the group elements corresponding to loops around punctures are parabolic ( $|\text{tr}| = 2$ )

Representation-theoretic definition:

$$T(F) = \text{Hom}'(\pi_1(F), PSL(2, \mathbb{R})) / PSL(2, \mathbb{R}),$$

where  $\rho \in \text{Hom}'$  if

- ▶  $\rho$  is injective
- ▶ identity in  $PSL(2, \mathbb{R})$  is not an accumulation point of the image of  $\rho$ , i.e.  $\rho$  is discrete
- ▶ the group elements corresponding to loops around punctures are parabolic ( $|\text{tr}| = 2$ )

The image  $\Gamma \in PSL(2, \mathbb{R})$  is a *Fuchsian group*.

By Poincaré uniformization we have  $F = H^+/\Gamma$ , where  $PSL(2, \mathbb{R})$  acts on the hyperbolic upper-half plane  $H^+$  as oriented isometries, given by fractional-linear transformations:

$$z \rightarrow \frac{az + b}{cz + d}.$$

The punctures of  $\tilde{F} = H^+$  belong to the real line  $\partial H^+$ , which is called *absolute*.



The image  $\Gamma \in PSL(2, \mathbb{R})$  is a *Fuchsian group*.

By Poincaré uniformization we have  $F = H^+/\Gamma$ , where  $PSL(2, \mathbb{R})$  acts on the hyperbolic upper-half plane  $H^+$  as oriented isometries, given by fractional-linear transformations:

$$z \rightarrow \frac{az + b}{cz + d}.$$

The punctures of  $\tilde{F} = H^+$  belong to the real line  $\partial H^+$ , which is called *absolute*.

The image  $\Gamma \in PSL(2, \mathbb{R})$  is a *Fuchsian group*.

By Poincaré uniformization we have  $F = H^+/\Gamma$ , where  $PSL(2, \mathbb{R})$  acts on the hyperbolic upper-half plane  $H^+$  as oriented isometries, given by fractional-linear transformations:

$$z \rightarrow \frac{az + b}{cz + d}.$$

The punctures of  $\tilde{F} = H^+$  belong to the real line  $\partial H^+$ , which is called *absolute*.

The image  $\Gamma \in PSL(2, \mathbb{R})$  is a *Fuchsian group*.

By Poincaré uniformization we have  $F = H^+/\Gamma$ , where  $PSL(2, \mathbb{R})$  acts on the hyperbolic upper-half plane  $H^+$  as oriented isometries, given by fractional-linear transformations:

$$z \rightarrow \frac{az + b}{cz + d}.$$

The punctures of  $\tilde{F} = H^+$  belong to the real line  $\partial H^+$ , which is called *absolute*.

The primary object of interest in many areas of mathematics is the *moduli space*:

$$M(F) = T(F)/MC(F).$$

The *mapping class group*  $MC(F)$ : a group of the homotopy classes of orientation preserving homeomorphisms.

$MC(F)$  acts on  $T(F)$  by outer automorphisms of  $\pi_1(F)$ .

The goal is to find a system of coordinates on  $T(F)$ , so that the action of  $MC(F)$  is realized in the simplest possible way.

The primary object of interest in many areas of mathematics is the *moduli space*:

$$M(F) = T(F)/MC(F).$$

The *mapping class group*  $MC(F)$ : a group of the homotopy classes of orientation preserving homeomorphisms.

$MC(F)$  acts on  $T(F)$  by outer automorphisms of  $\pi_1(F)$ .

The goal is to find a system of coordinates on  $T(F)$ , so that the action of  $MC(F)$  is realized in the simplest possible way.

The primary object of interest in many areas of mathematics is the *moduli space*:

$$M(F) = T(F)/MC(F).$$

The *mapping class group*  $MC(F)$ : a group of the homotopy classes of orientation preserving homeomorphisms.

$MC(F)$  acts on  $T(F)$  by outer automorphisms of  $\pi_1(F)$ .

The goal is to find a system of coordinates on  $T(F)$ , so that the action of  $MC(F)$  is realized in the simplest possible way.

The primary object of interest in many areas of mathematics is the *moduli space*:


$$M(F) = T(F)/MC(F).$$

The *mapping class group*  $MC(F)$ : a group of the homotopy classes of orientation preserving homeomorphisms.

$MC(F)$  acts on  $T(F)$  by outer automorphisms of  $\pi_1(F)$ .

The goal is to find a system of coordinates on  $T(F)$ , so that the action of  $MC(F)$  is realized in the simplest possible way.

R. Penner's work in the 1980s: a construction of coordinates associated to the ideal triangulation of  $F$ :



so that one assigns one positive number  $\lambda$ -length for every edge.

This provides coordinates for the decorated Teichmüller space:

$$\tilde{T}(F) = \mathbb{R}_+^5 \times T(F)$$

- Positive parameters correspond to the "renormalized" geodesic lengths ( $\lambda = e^{\delta/2}$ )
- $\mathbb{R}_+^5$ -fiber provides cut-off parameter (determining the size of the horocycle) for every puncture.

Outline

Introduction

Cast of characters

Coordinates on  
Super-Teichmüller  
space

$\mathcal{N} = 2$   
Super-Teichmüller  
theory

Further work


Open problems







R. Penner's work in the 1980s: a construction of coordinates associated to the ideal triangulation of  $F$ :




so that one assigns one positive number  $\lambda$ -length for every edge.

This provides coordinates for the decorated Teichmüller space:


$$\tilde{T}(F) = \mathbb{R}_+^s \times T(F)$$

- Positive parameters correspond to the "renormalized" geodesic lengths ( $\lambda = e^{\delta/2}$ )



- $\mathbb{R}_+^s$ -fiber provides cut-off parameter (determining the size of the horocycle) for every puncture.


The action of  $MC(F)$  can be described combinatorially using elementary transformations called flips:



Ptolemy relation :  $ef = ac + bd$

In order to obtain coordinates on  $T(F)$ , one has to consider *shear coordinates*  $z_e = \log\left(\frac{ac}{bd}\right)$ , which are subjects to certain linear constraints.


The action of  $MC(F)$  can be described combinatorially using elementary transformations called flips:



Ptolemy relation :  $ef = ac + bd$

In order to obtain coordinates on  $T(F)$ , one has to consider *shear coordinates*  $z_e = \log\left(\frac{ac}{bd}\right)$ , which are subjects to certain linear constraints.

The action of  $MC(F)$  can be described combinatorially using elementary transformations called flips:



Ptolemy relation :  $ef = ac + bd$

In order to obtain coordinates on  $T(F)$ , one has to consider *shear coordinates*  $z_e = \log\left(\frac{ac}{bd}\right)$ , which are subjects to certain linear constraints.

Transformation of coordinates via the triangulation change is therefore governed by Ptolemy relations. This leads to the prominent geometric example of *cluster algebra*, introduced by [S. Fomin](#) and [A. Zelevinsky](#) in the early 2000s.

Penner's coordinates can be used for the quantization of  $T(F)$  ([L. Chekhov](#), [V. Fock](#), [R. Kashaev](#), late 90s, early 2000s).

Higher Teichmüller spaces:  $PSL(2, \mathbb{R})$  is replaced by some split semisimple real Lie group  $G$ .

In the case of real reductive groups  $G$  the construction of coordinates was given by [V. Fock](#) and [A. Goncharov](#) (2003) and sparked a lot of applications in various areas of mathematics/mathematical physics.

Transformation of coordinates via the triangulation change is therefore governed by Ptolemy relations. This leads to the prominent geometric example of *cluster algebra*, introduced by [S. Fomin](#) and [A. Zelevinsky](#) in the early 2000s.

Penner's coordinates can be used for the quantization of  $T(F)$  ([L. Chekhov](#), [V. Fock](#), [R. Kashaev](#), late 90s, early 2000s).

Higher Teichmüller spaces:  $PSL(2, \mathbb{R})$  is replaced by some split semisimple real Lie group  $G$ .

In the case of real reductive groups  $G$  the construction of coordinates was given by [V. Fock](#) and [A. Goncharov](#) (2003) and sparked a lot of applications in various areas of mathematics/mathematical physics.



Transformation of coordinates via the triangulation change is therefore governed by Ptolemy relations. This leads to the prominent geometric example of *cluster algebra*, introduced by [S. Fomin](#) and [A. Zelevinsky](#) in the early 2000s.

Penner's coordinates can be used for the quantization of  $T(F)$  ([L. Chekhov](#), [V. Fock](#), [R. Kashaev](#), late 90s, early 2000s).

Higher Teichmüller spaces:  $PSL(2, \mathbb{R})$  is replaced by some split semisimple real Lie group  $G$ .

In the case of real reductive groups  $G$  the construction of coordinates was given by [V. Fock](#) and [A. Goncharov](#) (2003) and sparked a lot of applications in various areas of mathematics/mathematical physics.

Transformation of coordinates via the triangulation change is therefore governed by Ptolemy relations. This leads to the prominent geometric example of *cluster algebra*, introduced by [S. Fomin](#) and [A. Zelevinsky](#) in the early 2000s.

Penner's coordinates can be used for the quantization of  $T(F)$  ([L. Chekhov](#), [V. Fock](#), [R. Kashaev](#), late 90s, early 2000s).

Higher Teichmüller spaces:  $PSL(2, \mathbb{R})$  is replaced by some split semisimple real Lie group  $G$ .

In the case of real reductive groups  $G$  the construction of coordinates was given by [V. Fock](#) and [A. Goncharov](#) (2003) and sparked a lot of applications in various areas of mathematics/mathematical physics.

String theory: propagating closed strings generate Riemann surfaces:



*Superstrings*, which, according to string theory, are the fundamental objects for the description of our world, carry extra anticommuting parameters  $\theta^i$ , called *fermions*:

$$\theta^i \theta^j = -\theta^j \theta^i$$

That can be interpreted as strings propagating along *supermanifolds* called *super Riemann surfaces*.

Outline

**Introduction**

Cast of characters


Coordinates on  
Super-Teichmüller  
space

$\mathcal{N} = 2$   
Super-Teichmüller  
theory

Further work

Open problems

String theory: propagating closed strings generate Riemann surfaces:




*Superstrings*, which, according to string theory, are the fundamental objects for the description of our world, carry extra anticommuting parameters  $\theta^i$ , called *fermions*:

$$\theta^i \theta^j = -\theta^j \theta^i$$

That can be interpreted as strings propagating along *supermanifolds* called *super Riemann surfaces*.

[Outline](#)[Introduction](#)[Cast of characters](#)[Coordinates on  
Super-Teichmüller  
space](#)[N = 2  
Super-Teichmüller  
theory](#)[Further work](#)[Open problems](#)

String theory: propagating closed strings generate Riemann surfaces:



*Superstrings*, which, according to string theory, are the fundamental objects for the description of our world, carry extra anticommuting parameters  $\theta^i$ , called *fermions*:

$$\theta^i \theta^j = -\theta^j \theta^i$$

That can be interpreted as strings propagating along *supermanifolds* called *super Riemann surfaces*.

[Outline](#)[Introduction](#)[Cast of characters](#)[Coordinates on  
Super-Teichmüller  
space](#) [\$\mathcal{N} = 2\$   
Super-Teichmüller  
theory](#)[Further work](#)[Open problems](#)

That leads to generalizations of Teichmüller spaces, relevant for string theory, called  $\mathcal{N} = 1$  and  $\mathcal{N} = 2$  super-Teichmüller spaces  $ST(F)$ , depending on the number of extra fermionic degrees of freedom.

The corresponding supermoduli spaces were intensively studied by various physicists and mathematicians [L. Crane](#), [J. Rabin](#), [E. D'Hoker](#), [D. Phong](#), [A. Schwarz](#), [A. Voronov](#)...

Not so long ago [R. Donagi](#) and [E. Witten](#) showed that in the higher genus supermoduli spaces are very much involved:

[R. Donagi, E. Witten, \*Supermoduli Space Is Not Projected\*, arXiv:1304.7798](#)

That leads to generalizations of Teichmüller spaces, relevant for string theory, called  $\mathcal{N} = 1$  and  $\mathcal{N} = 2$  super-Teichmüller spaces  $ST(F)$ , depending on the number of extra fermionic degrees of freedom.

The corresponding supermoduli spaces were intensively studied by various physicists and mathematicians [L. Crane](#), [J. Rabin](#), [E. D'Hoker](#), [D. Phong](#), [A. Schwarz](#), [A. Voronov](#)...

Not so long ago [R. Donagi](#) and [E. Witten](#) showed that in the higher genus supermoduli spaces are very much involved:

[R. Donagi, E. Witten, \*Supermoduli Space Is Not Projected\*, arXiv:1304.7798](#)

That leads to generalizations of Teichmüller spaces, relevant for string theory, called  $\mathcal{N} = 1$  and  $\mathcal{N} = 2$  super-Teichmüller spaces  $ST(F)$ , depending on the number of extra fermionic degrees of freedom.

The corresponding supermoduli spaces were intensively studied by various physicists and mathematicians [L. Crane](#), [J. Rabin](#), [E. D'Hoker](#), [D. Phong](#), [A. Schwarz](#), [A. Voronov](#)...

Not so long ago [R. Donagi](#) and [E. Witten](#) showed that in the higher genus supermoduli spaces are very much involved:

[R. Donagi, E. Witten, \*Supermoduli Space Is Not Projected\*, arXiv:1304.7798](#)



That leads to generalizations of Teichmüller spaces, relevant for string theory, called  $\mathcal{N} = 1$  and  $\mathcal{N} = 2$  super-Teichmüller spaces  $ST(F)$ , depending on the number of extra fermionic degrees of freedom.

The corresponding supermoduli spaces were intensively studied by various physicists and mathematicians [L. Crane](#), [J. Rabin](#), [E. D'Hoker](#), [D. Phong](#), [A. Schwarz](#), [A. Voronov](#)...

Not so long ago [R. Donagi](#) and [E. Witten](#) showed that in the higher genus supermoduli spaces are very much involved:

[R. Donagi, E. Witten, \*Supermoduli Space Is Not Projected\*, arXiv:1304.7798](#)

These  $\mathcal{N} = 1$  and  $\mathcal{N} = 2$  super-Teichmüller spaces in the terminology of higher Teichmüller theory are related to *supergroups*

$$OSP(1|2), \quad OSP(2|2)$$

correspondingly.

In the late 80s the problem of construction of Penner's coordinates on  $ST(F)$  was introduced on [Yu.I. Manin's](#) Moscow seminar.

The  $\mathcal{N} = 1$  case was solved in:

[R. Penner](#), [A. Zeitlin](#), arXiv:1509.06302, to appear in [J. Diff. Geom.](#)

The  $\mathcal{N} = 2$  case was solved in:

[I. Ip](#), [R. Penner](#), [A. Zeitlin](#), [Adv. Math.](#) 336 (2018) 409-454, arXiv:1605.08094.

Full decoration removal for  $\mathcal{N} = 1$ :

[I. Ip](#), [R. Penner](#), [A. Zeitlin](#), arXiv:1709.06207, to appear in [Comm. Math. Phys.](#)

[Outline](#)[Introduction](#)[Cast of characters](#)[Coordinates on Super-Teichmüller space](#) [\$\mathcal{N} = 2\$  Super-Teichmüller theory](#)[Further work](#)[Open problems](#)

These  $\mathcal{N} = 1$  and  $\mathcal{N} = 2$  super-Teichmüller spaces in the terminology of higher Teichmüller theory are related to *supergroups*

$$OSP(1|2), \quad OSP(2|2)$$

correspondingly.

In the late 80s the problem of construction of Penner's coordinates on  $ST(F)$  was introduced on [Yu.I. Manin's](#) Moscow seminar.

The  $\mathcal{N} = 1$  case was solved in:

[R. Penner, A. Zeitlin](#), arXiv:1509.06302, to appear in *J. Diff. Geom.*

The  $\mathcal{N} = 2$  case was solved in:

[I. Ip, R. Penner, A. Zeitlin](#), *Adv. Math.* 336 (2018) 409-454, arXiv:1605.08094.

Full decoration removal for  $\mathcal{N} = 1$ :

[I. Ip, R. Penner, A. Zeitlin](#), arXiv:1709.06207, to appear in *Comm. Math. Phys.*

[Outline](#)[Introduction](#)[Cast of characters](#)[Coordinates on  
Super-Teichmüller  
space](#) [\$\mathcal{N} = 2\$   
Super-Teichmüller  
theory](#)[Further work](#)[Open problems](#)

These  $\mathcal{N} = 1$  and  $\mathcal{N} = 2$  super-Teichmüller spaces in the terminology of higher Teichmüller theory are related to *supergroups*

$$OSP(1|2), \quad OSP(2|2)$$

correspondingly.

In the late 80s the problem of construction of Penner's coordinates on  $ST(F)$  was introduced on [Yu.I. Manin's](#) Moscow seminar.

The  $\mathcal{N} = 1$  case was solved in:

[R. Penner](#), [A. Zeitlin](#), arXiv:1509.06302, to appear in [J. Diff. Geom.](#)

The  $\mathcal{N} = 2$  case was solved in:

[I. Ip](#), [R. Penner](#), [A. Zeitlin](#), [Adv. Math.](#) 336 (2018) 409-454, arXiv:1605.08094.

Full decoration removal for  $\mathcal{N} = 1$ :

[I. Ip](#), [R. Penner](#), [A. Zeitlin](#), arXiv:1709.06207, to appear in [Comm. Math. Phys.](#)

[Outline](#)[Introduction](#)[Cast of characters](#)[Coordinates on Super-Teichmüller space](#) [\$\mathcal{N} = 2\$  Super-Teichmüller theory](#)[Further work](#)[Open problems](#)

## i) Superspaces and supermanifolds

Let  $\Lambda(\mathbb{K}) = \Lambda^0(\mathbb{K}) \oplus \Lambda^1(\mathbb{K})$  be an exterior algebra over field  $\mathbb{K} = \mathbb{R}, \mathbb{C}$  with (in)finitely many generators  $1, e_1, e_2, \dots$ , so that

$$a = a^\# + \sum_i a_i e_i + \sum_{ij} a_{ij} e_i \wedge e_j + \dots, \quad \# : \Lambda(\mathbb{K}) \rightarrow \mathbb{K}$$

$a^\#$  is referred to as a *body* of a supernumber.

If  $a \in \Lambda^0(\mathbb{K})$ , it is called even (bosonic) number

If  $a \in \Lambda^1(\mathbb{K})$ , it is called odd (fermionic) number

Note, that odd numbers anticommute.

## i) Superspaces and supermanifolds

Let  $\Lambda(\mathbb{K}) = \Lambda^0(\mathbb{K}) \oplus \Lambda^1(\mathbb{K})$  be an exterior algebra over field  $\mathbb{K} = \mathbb{R}, \mathbb{C}$  with (in)finitely many generators  $1, e_1, e_2, \dots$ , so that

$$a = a^\# + \sum_i a_i e_i + \sum_{ij} a_{ij} e_i \wedge e_j + \dots, \quad \# : \Lambda(\mathbb{K}) \rightarrow \mathbb{K}$$

$a^\#$  is referred to as a *body* of a supernumber.

If  $a \in \Lambda^0(\mathbb{K})$ , it is called even (bosonic) number

If  $a \in \Lambda^1(\mathbb{K})$ , it is called odd (fermionic) number

Note, that odd numbers anticommute.

## i) Superspaces and supermanifolds

Let  $\Lambda(\mathbb{K}) = \Lambda^0(\mathbb{K}) \oplus \Lambda^1(\mathbb{K})$  be an exterior algebra over field  $\mathbb{K} = \mathbb{R}, \mathbb{C}$  with (in)finitely many generators  $1, e_1, e_2, \dots$ , so that

$$a = a^\# + \sum_i a_i e_i + \sum_{ij} a_{ij} e_i \wedge e_j + \dots, \quad \# : \Lambda(\mathbb{K}) \rightarrow \mathbb{K}$$

$a^\#$  is referred to as a *body* of a supernumber.

If  $a \in \Lambda^0(\mathbb{K})$ , it is called even (bosonic) number

If  $a \in \Lambda^1(\mathbb{K})$ , it is called odd (fermionic) number

Note, that odd numbers anticommute.

Superspace  $\mathbb{K}^{(n|m)}$  is:

$$\mathbb{K}^{(n|m)} = \{(z_1, z_2, \dots, z_n | \theta_1, \theta_2, \dots, \theta_m) : z_i \in \Lambda^0(\mathbb{K}), \theta_j \in \Lambda^1(\mathbb{K})\}$$

One can define  $(n|m)$  supermanifolds over  $\Lambda(\mathbb{K})$  based on superspaces  $\mathbb{K}^{(n|m)}$ , where  $\{z_i\}$  and  $\{\theta_j\}$  serve as *even and odd coordinates*.

Special spaces:

- Upper  $\mathcal{N} = N$  super-half-plane (we will need  $\mathcal{N} = 1, 2$ ):

$$H^+ = \{(z | \theta_1, \theta_2, \dots, \theta_N) \in \mathbb{C}^{(1|N)} \mid \operatorname{Im} z^\# > 0\}$$

- Positive superspace:

$$\mathbb{R}_+^{(n|m)} = \{(z_1, z_2, \dots, z_n | \theta_1, \theta_2, \dots, \theta_m) \in \mathbb{R}^{(n|m)} \mid z_i^\# > 0, i = 1, \dots, n\}$$



Superspace  $\mathbb{K}^{(n|m)}$  is:

$$\mathbb{K}^{(n|m)} = \{(z_1, z_2, \dots, z_n | \theta_1, \theta_2, \dots, \theta_m) : z_i \in \Lambda^0(\mathbb{K}), \theta_j \in \Lambda^1(\mathbb{K})\}$$

One can define  $(n|m)$  supermanifolds over  $\Lambda(\mathbb{K})$  based on superspaces  $\mathbb{K}^{(n|m)}$ , where  $\{z_i\}$  and  $\{\theta_j\}$  serve as *even and odd coordinates*.

Special spaces:

- Upper  $\mathcal{N} = N$  super-half-plane (we will need  $\mathcal{N} = 1, 2$ ):

$$H^+ = \{(z | \theta_1, \theta_2, \dots, \theta_N) \in \mathbb{C}^{(1|N)} \mid \text{Im } z^\# > 0\}$$

- Positive superspace:

$$\mathbb{R}_+^{(n|m)} = \{(z_1, z_2, \dots, z_n | \theta_1, \theta_2, \dots, \theta_m) \in \mathbb{R}^{(n|m)} \mid z_i^\# > 0, i = 1, \dots, n\}$$

Superspace  $\mathbb{K}^{(n|m)}$  is:

$$\mathbb{K}^{(n|m)} = \{(z_1, z_2, \dots, z_n | \theta_1, \theta_2, \dots, \theta_m) : z_i \in \Lambda^0(\mathbb{K}), \theta_j \in \Lambda^1(\mathbb{K})\}$$

One can define  $(n|m)$  supermanifolds over  $\Lambda(\mathbb{K})$  based on superspaces  $\mathbb{K}^{(n|m)}$ , where  $\{z_i\}$  and  $\{\theta_j\}$  serve as *even and odd coordinates*.

Special spaces:

- Upper  $\mathcal{N} = N$  super-half-plane (we will need  $\mathcal{N} = 1, 2$ ):

$$H^+ = \{(z | \theta_1, \theta_2, \dots, \theta_N) \in \mathbb{C}^{(1|N)} \mid \text{Im } z^\# > 0\}$$

- Positive superspace:

$$\mathbb{R}_+^{(n|m)} = \{(z_1, z_2, \dots, z_n | \theta_1, \theta_2, \dots, \theta_m) \in \mathbb{R}^{(n|m)} \mid z_i^\# > 0, i = 1, \dots, n\}$$

Superspace  $\mathbb{K}^{(n|m)}$  is:

$$\mathbb{K}^{(n|m)} = \{(z_1, z_2, \dots, z_n | \theta_1, \theta_2, \dots, \theta_m) : z_i \in \Lambda^0(\mathbb{K}), \theta_j \in \Lambda^1(\mathbb{K})\}$$

One can define  $(n|m)$  supermanifolds over  $\Lambda(\mathbb{K})$  based on superspaces  $\mathbb{K}^{(n|m)}$ , where  $\{z_i\}$  and  $\{\theta_j\}$  serve as *even and odd coordinates*.

Special spaces:

- Upper  $\mathcal{N} = N$  super-half-plane (we will need  $\mathcal{N} = 1, 2$ ):

$$H^+ = \{(z | \theta_1, \theta_2, \dots, \theta_N) \in \mathbb{C}^{(1|N)} \mid \text{Im } z^\# > 0\}$$

- Positive superspace:

$$\mathbb{R}_+^{(n|m)} = \{(z_1, z_2, \dots, z_n | \theta_1, \theta_2, \dots, \theta_m) \in \mathbb{R}^{(n|m)} \mid z_i^\# > 0, i = 1, \dots, n\}$$

ii) Supergroup  $OSp(1|2)$ 

Definition:

$(1|2) \times (1|2)$  supermatrices  $g$ , obeying the relation

$$g^{st} J g = J,$$

where

$$J = \begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix}$$

and the supertranspose  $g^{st}$  of  $g$  is given by

$$g = \begin{pmatrix} a & b & \alpha \\ c & d & \beta \\ \gamma & \delta & f \end{pmatrix} \quad \text{implies} \quad g^{st} = \begin{pmatrix} a & c & \gamma \\ b & d & \delta \\ -\alpha & -\beta & f \end{pmatrix}.$$

We want a connected component of identity, so we assume that Berezinian (super-analogue of determinant) = 1.

Outline

Introduction

Cast of characters

Coordinates on  
Super-Teichmüller  
space

$\mathcal{N} = 2$   
Super-Teichmüller  
theory

Further work

Open problems

ii) Supergroup  $OSp(1|2)$ 

Definition:

$(1|2) \times (1|2)$  supermatrices  $g$ , obeying the relation

$$g^{st} J g = J,$$

where

$$J = \begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix}$$

and the supertranspose  $g^{st}$  of  $g$  is given by

$$g = \begin{pmatrix} a & b & \alpha \\ c & d & \beta \\ \gamma & \delta & f \end{pmatrix} \quad \text{implies} \quad g^{st} = \begin{pmatrix} a & c & \gamma \\ b & d & \delta \\ -\alpha & -\beta & f \end{pmatrix}.$$

We want a connected component of identity, so we assume that Berezinian (super-analogue of determinant) = 1.

Outline

Introduction

Cast of characters

Coordinates on  
Super-Teichmüller  
space

$\mathcal{N} = 2$   
Super-Teichmüller  
theory

Further work

Open problems

Some remarks:

- Lie superalgebra  $osp(1|2)$ :

Three even  $h, X_{\pm}$  and two odd  $v_{\pm}$  generators, satisfying the following commutation relations:

$$[h, v_{\pm}] = \pm v_{\pm}, \quad [v_{\pm}, v_{\pm}] = \mp 2X_{\pm}, \quad [v_+, v_-] = h.$$

- Note, that the *body* of the supergroup  $OSP(1|2)$  is  $SL(2, \mathbb{R})$ , not  $PSL(2, \mathbb{R})$ !

Some remarks:

- Lie superalgebra  $osp(1|2)$ :

Three even  $h, X_{\pm}$  and two odd  $v_{\pm}$  generators, satisfying the following commutation relations:

$$[h, v_{\pm}] = \pm v_{\pm}, \quad [v_{\pm}, v_{\pm}] = \mp 2X_{\pm}, \quad [v_+, v_-] = h.$$

- Note, that the *body* of the supergroup  $OSP(1|2)$  is  $SL(2, \mathbb{R})$ , not  $PSL(2, \mathbb{R})$ !

$OSp(1|2)$  acts on  $\mathcal{N} = 1$  super half-plane  $H^+$ , with the absolute  $\partial H^+ = \mathbb{R}^{1|1}$  by superconformal fractional-linear transformations:

$$z \rightarrow \frac{az + b}{cz + d} + \eta \frac{\gamma z + \delta}{(cz + d)^2},$$
$$\eta \rightarrow \frac{\gamma z + \delta}{cz + d} + \eta \frac{1 + \frac{1}{2}\delta\gamma}{cz + d}.$$

Factor  $H^+/\Gamma$ , where  $\Gamma$  is a discrete subgroup of  $OSp(1|2)$ , such that its projection is a Fuchsian group, are called *super Riemann surfaces*.



$OSp(1|2)$  acts on  $\mathcal{N} = 1$  super half-plane  $H^+$ , with the absolute  $\partial H^+ = \mathbb{R}^{1|1}$  by superconformal fractional-linear transformations:

$$z \rightarrow \frac{az + b}{cz + d} + \eta \frac{\gamma z + \delta}{(cz + d)^2},$$
$$\eta \rightarrow \frac{\gamma z + \delta}{cz + d} + \eta \frac{1 + \frac{1}{2}\delta\gamma}{cz + d}.$$

Factor  $H^+/\Gamma$ , where  $\Gamma$  is a discrete subgroup of  $OSp(1|2)$ , such that its projection is a Fuchsian group, are called *super Riemann surfaces*.

Alternatively, *super Riemann surface* is a complex  $(1|1)$ -supermanifold  $S$  with everywhere non-integrable odd distribution  $\mathcal{D} \in TS$ , such that

$$0 \rightarrow \mathcal{D} \rightarrow TS \rightarrow \mathcal{D}^2 \rightarrow 0 \quad \text{is exact.}$$

There are more general fractional-linear transformations acting on  $H^+$ . They correspond to  $SL(1|2)$  supergroup, and factors  $H^+/\Gamma$  give  $(1|1)$ -supermanifolds which have relation to  $\mathcal{N} = 2$  super-Teichmüller theory.

Alternatively, *super Riemann surface* is a complex  $(1|1)$ -supermanifold  $S$  with everywhere non-integrable odd distribution  $\mathcal{D} \in TS$ , such that

$$0 \rightarrow \mathcal{D} \rightarrow TS \rightarrow \mathcal{D}^2 \rightarrow 0 \quad \text{is exact.}$$

There are more general fractional-linear transformations acting on  $H^+$ . They correspond to  $SL(1|2)$  supergroup, and factors  $H^+/\Gamma$  give  $(1|1)$ -supermanifolds which have relation to  $\mathcal{N} = 2$  super-Teichmüller theory.

### iii) ( $\mathcal{N} = 1$ ) Super-Teichmüller space

From now on let

$$ST(F) = \text{Hom}'(\pi_1(F), OSp(1|2))/OSp(1|2).$$

Super-Fuchsian representations comprising  $\text{Hom}'$  are defined to be those whose projections

$$\pi_1 \rightarrow OSp(1|2) \rightarrow SL(2, \mathbb{R}) \rightarrow PSL(2, \mathbb{R})$$

are Fuchsian groups, corresponding to  $F$ .

Trivial bundle  $S\tilde{T}(F) = \mathbb{R}_+^s \times ST(F)$  is called the decorated super-Teichmüller space.

Unlike (decorated) Teichmüller space,  $ST(F)$  ( $S\tilde{T}(F)$ ) has  $2^{2g+s-1}$  connected components labeled by spin structures on  $F$ .

iii) ( $\mathcal{N} = 1$ ) Super-Teichmüller space

From now on let

$$ST(F) = \text{Hom}'(\pi_1(F), \text{OSp}(1|2)) / \text{OSp}(1|2).$$

Super-Fuchsian representations comprising  $\text{Hom}'$  are defined to be those whose projections

$$\pi_1 \rightarrow \text{OSp}(1|2) \rightarrow \text{SL}(2, \mathbb{R}) \rightarrow \text{PSL}(2, \mathbb{R})$$

are Fuchsian groups, corresponding to  $F$ .

Trivial bundle  $S\tilde{T}(F) = \mathbb{R}_+^s \times ST(F)$  is called the decorated super-Teichmüller space.

Unlike (decorated) Teichmüller space,  $ST(F)$  ( $S\tilde{T}(F)$ ) has  $2^{2g+s-1}$  connected components labeled by spin structures on  $F$ .

iii) ( $\mathcal{N} = 1$ ) Super-Teichmüller space

From now on let

$$ST(F) = \text{Hom}'(\pi_1(F), \text{OSp}(1|2)) / \text{OSp}(1|2).$$

Super-Fuchsian representations comprising  $\text{Hom}'$  are defined to be those whose projections

$$\pi_1 \rightarrow \text{OSp}(1|2) \rightarrow \text{SL}(2, \mathbb{R}) \rightarrow \text{PSL}(2, \mathbb{R})$$

are Fuchsian groups, corresponding to  $F$ .

Trivial bundle  $S\tilde{T}(F) = \mathbb{R}_+^s \times ST(F)$  is called the decorated super-Teichmüller space.

Unlike (decorated) Teichmüller space,  $ST(F)$  ( $S\tilde{T}(F)$ ) has  $2^{2g+s-1}$  connected components labeled by spin structures on  $F$ .

iii) ( $\mathcal{N} = 1$ ) Super-Teichmüller space

From now on let

$$ST(F) = \text{Hom}'(\pi_1(F), \text{OSp}(1|2)) / \text{OSp}(1|2).$$

Super-Fuchsian representations comprising  $\text{Hom}'$  are defined to be those whose projections

$$\pi_1 \rightarrow \text{OSp}(1|2) \rightarrow \text{SL}(2, \mathbb{R}) \rightarrow \text{PSL}(2, \mathbb{R})$$

are Fuchsian groups, corresponding to  $F$ .

Trivial bundle  $S\tilde{T}(F) = \mathbb{R}_+^s \times ST(F)$  is called the decorated super-Teichmüller space.

Unlike (decorated) Teichmüller space,  $ST(F)$  ( $S\tilde{T}(F)$ ) has  $2^{2g+s-1}$  connected components labeled by spin structures on  $F$ .

#### iv) Ideal triangulations and trivalent fatgraphs

- Ideal triangulation of  $F$ : triangulation  $\Delta$  of  $F$  with punctures at the vertices, so that each arc connecting punctures is not homotopic to a point rel punctures.
- Trivalent fatgraph: trivalent graph  $\tau$  with cyclic orderings on half-edges about each vertex.

$\tau = \tau(\Delta)$ , if the following is true:

- 1) one fatgraph vertex per triangle
- 2) one edge of fatgraph intersects one shared edge of triangulation.



#### iv) Ideal triangulations and trivalent fatgraphs

- Ideal triangulation of  $F$ : triangulation  $\Delta$  of  $F$  with punctures at the vertices, so that each arc connecting punctures is not homotopic to a point rel punctures.
- Trivalent fatgraph: trivalent graph  $\tau$  with cyclic orderings on half-edges about each vertex.

$\tau = \tau(\Delta)$ , if the following is true:

- 1) one fatgraph vertex per triangle
- 2) one edge of fatgraph intersects one shared edge of triangulation.

#### iv) Ideal triangulations and trivalent fatgraphs

- Ideal triangulation of  $F$ : triangulation  $\Delta$  of  $F$  with punctures at the vertices, so that each arc connecting punctures is not homotopic to a point rel punctures.
- Trivalent fatgraph: trivalent graph  $\tau$  with cyclic orderings on half-edges about each vertex.

$\tau = \tau(\Delta)$ , if the following is true:



- 1) one fatgraph vertex per triangle
- 2) one edge of fatgraph intersects one shared edge of triangulation.

## iv) Ideal triangulations and trivalent fatgraphs

- Ideal triangulation of  $F$ : triangulation  $\Delta$  of  $F$  with punctures at the vertices, so that each arc connecting punctures is not homotopic to a point rel punctures.
- Trivalent fatgraph: trivalent graph  $\tau$  with cyclic orderings on half-edges about each vertex.

$\tau = \tau(\Delta)$ , if the following is true:

- 1) one fatgraph vertex per triangle
- 2) one edge of fatgraph intersects one shared edge of triangulation.

Fatgraph for  $F_1^1$ Fatgraph for  $F_0^3$

## v) Spin structures

Textbook definition:

Let  $M$  be an oriented  $n$ -dimensional Riemannian manifold,  $P_{SO}$  is an orthonormal frame bundle, associated with  $TM$ . A *spin structure* is a 2-fold covering map  $P \rightarrow P_{SO}$ , which restricts to  $Spin(n) \rightarrow SO(n)$  on each fiber.

This is not really useful for us, since we want to relate it to combinatorial geometric structures on  $F$ .

## v) Spin structures

Textbook definition:

Let  $M$  be an oriented  $n$ -dimensional Riemannian manifold,  $P_{SO}$  is an orthonormal frame bundle, associated with  $TM$ . A *spin structure* is a 2-fold covering map  $P \rightarrow P_{SO}$ , which restricts to  $Spin(n) \rightarrow SO(n)$  on each fiber.

This is not really useful for us, since we want to relate it to combinatorial geometric structures on  $F$ .

## v) Spin structures

Textbook definition:

Let  $M$  be an oriented  $n$ -dimensional Riemannian manifold,  $P_{SO}$  is an orthonormal frame bundle, associated with  $TM$ . A *spin structure* is a 2-fold covering map  $P \rightarrow P_{SO}$ , which restricts to  $Spin(n) \rightarrow SO(n)$  on each fiber.

This is not really useful for us, since we want to relate it to combinatorial geometric structures on  $F$ .

There are several ways to describe spin structures on  $F$ :

- [D. Johnson \(1980\)](#):

Quadratic forms  $q : H_1(F, \mathbb{Z}_2) \rightarrow \mathbb{Z}_2$ , which are quadratic with respect to the intersection pairing  $\cdot : H_1 \otimes H_1 \rightarrow \mathbb{Z}_2$ , i.e.

$$q(a + b) = q(a) + q(b) + a \cdot b \text{ if } a, b \in H_1.$$

- [S. Natanzon](#):

A spin structure on a uniformized surface  $F = \mathcal{U}/\Gamma$  is determined by a lift  $\tilde{\rho} : \pi_1 \rightarrow SL(2, \mathbb{R})$  of  $\rho : \pi_1 \rightarrow PSL_2(\mathbb{R})$ . Quadratic form  $q$  is computed using the following rules:  $\text{trace } \tilde{\rho}(\gamma) > 0$  if and only if  $q([\gamma]) \neq 0$ , where  $[\gamma] \in H_1$  is the image of  $\gamma \in \pi_1$  under the mod two Hurewicz map.

There are several ways to describe spin structures on  $F$ :

- [D. Johnson \(1980\)](#):

Quadratic forms  $q : H_1(F, \mathbb{Z}_2) \rightarrow \mathbb{Z}_2$ , which are quadratic with respect to the intersection pairing  $\cdot : H_1 \otimes H_1 \rightarrow \mathbb{Z}_2$ , i.e.

$$q(a + b) = q(a) + q(b) + a \cdot b \text{ if } a, b \in H_1.$$

- [S. Natanzon](#):

A spin structure on a uniformized surface  $F = \mathcal{U}/\Gamma$  is determined by a lift  $\tilde{\rho} : \pi_1 \rightarrow SL(2, \mathbb{R})$  of  $\rho : \pi_1 \rightarrow PSL_2(\mathbb{R})$ . Quadratic form  $q$  is computed using the following rules:  $\text{trace } \tilde{\rho}(\gamma) > 0$  if and only if  $q([\gamma]) \neq 0$ , where  $[\gamma] \in H_1$  is the image of  $\gamma \in \pi_1$  under the mod two Hurewicz map.



- [D. Cimasoni and N. Reshetikhin \(2007\)](#):

Combinatorial description of spin structures in terms of the so-called Kasteleyn orientations and dimer configurations on the one-skeleton of a suitable CW decomposition of  $F$ . They derive a formula for the quadratic form in terms of that combinatorial data.


- We gave a substantial simplification of the combinatorial formulation of spin structures on  $F$  (one of the main results of [R. Penner](#), [A. Zeitlin](#), [arXiv:1509.06302](#)):

Equivalence classes  $\mathcal{O}(\tau)$  of all orientations on a trivalent fatgraph spine  $\tau \subset F$ , where the equivalence relation is generated by reversing the orientation of each edge incident on some fixed vertex, with the added bonus of a computable evolution under flips:



- We gave a substantial simplification of the combinatorial formulation of spin structures on  $F$  (one of the main results of [R. Penner, A. Zeitlin, arXiv:1509.06302](#)):

Equivalence classes  $\mathcal{O}(\tau)$  of all orientations on a trivalent fatgraph spine  $\tau \subset F$ , where the equivalence relation is generated by reversing the orientation of each edge incident on some fixed vertex, with the added bonus of a computable evolution under flips:



Fix a surface  $F = F_g^s$  as above and

- ▶  $\tau \subset F$  is some trivalent fatgraph spine
- ▶  $\omega$  is an orientation on the edges of  $\tau$  whose class in  $\mathcal{O}(\tau)$  determines the component  $C$  of  $S\tilde{T}(F)$

Then there are global affine coordinates on  $C$ :

- ▶ one even coordinate called a  $\lambda$ -length for each edge
- ▶ one odd coordinate called a  $\mu$ -invariant for each vertex of  $\tau$ ,

the latter of which are taken modulo an overall change of sign.

Alternating the sign in one of the fermions corresponds to the reflection on the spin graph.

The above  $\lambda$ -lengths and  $\mu$ -invariants establish a real-analytic homeomorphism

$$C \rightarrow \mathbb{R}_+^{6g-6+3s} / \mathbb{Z}_2.$$

[Outline](#)[Introduction](#)[Cast of characters](#)[Coordinates on  
Super-Teichmüller  
space](#) [\$\mathcal{N} = 2\$   
Super-Teichmüller  
theory](#)[Further work](#)[Open problems](#)

# Coordinates on $S\tilde{T}(F)$

Fix a surface  $F = F_g^s$  as above and

- ▶  $\tau \subset F$  is some trivalent fatgraph spine
- ▶  $\omega$  is an orientation on the edges of  $\tau$  whose class in  $\mathcal{O}(\tau)$  determines the component  $C$  of  $S\tilde{T}(F)$

Then there are global affine coordinates on  $C$ :

- ▶ one even coordinate called a  $\lambda$ -length for each edge
- ▶ one odd coordinate called a  $\mu$ -invariant for each vertex of  $\tau$ ,

the latter of which are taken modulo an overall change of sign.

Alternating the sign in one of the fermions corresponds to the reflection on the spin graph.

The above  $\lambda$ -lengths and  $\mu$ -invariants establish a real-analytic homeomorphism

$$C \rightarrow \mathbb{R}_+^{6g-6+3s} / \mathbb{Z}_2.$$

Fix a surface  $F = F_g^s$  as above and

- ▶  $\tau \subset F$  is some trivalent fatgraph spine
- ▶  $\omega$  is an orientation on the edges of  $\tau$  whose class in  $\mathcal{O}(\tau)$  determines the component  $C$  of  $S\tilde{T}(F)$

Then there are global affine coordinates on  $C$ :

- ▶ one even coordinate called a  $\lambda$ -length for each edge
- ▶ one odd coordinate called a  $\mu$ -invariant for each vertex of  $\tau$ ,

the latter of which are taken modulo an overall change of sign.

Alternating the sign in one of the fermions corresponds to the reflection on the spin graph.

The above  $\lambda$ -lengths and  $\mu$ -invariants establish a real-analytic homeomorphism

$$C \rightarrow \mathbb{R}_+^{6g-6+3s} / \mathbb{Z}_2.$$

Outline

Introduction

Cast of characters

Coordinates on  
Super-Teichmüller  
space

$\mathcal{N} = 2$   
Super-Teichmüller  
theory

Further work

Open problems

Fix a surface  $F = F_g^s$  as above and

- ▶  $\tau \subset F$  is some trivalent fatgraph spine
- ▶  $\omega$  is an orientation on the edges of  $\tau$  whose class in  $\mathcal{O}(\tau)$  determines the component  $C$  of  $S\tilde{T}(F)$

Then there are global affine coordinates on  $C$ :

- ▶ one even coordinate called a  $\lambda$ -length for each edge
- ▶ one odd coordinate called a  $\mu$ -invariant for each vertex of  $\tau$ ,

the latter of which are taken modulo an overall change of sign.


Alternating the sign in one of the fermions corresponds to the reflection on the spin graph.

The above  $\lambda$ -lengths and  $\mu$ -invariants establish a real-analytic homeomorphism

$$C \rightarrow \mathbb{R}_+^{6g-6+3s|4g-4+2s} / \mathbb{Z}_2.$$

[Outline](#)[Introduction](#)[Cast of characters](#)[Coordinates on  
Super-Teichmüller  
space](#) [\$\mathcal{N} = 2\$   
Super-Teichmüller  
theory](#)[Further work](#)[Open problems](#)

When all  $a, b, c, d$  are different edges of the triangulations of  $F$ ,



Ptolemy transformations are as follows:

$$ef = (ac + bd) \left( 1 + \frac{\sigma\theta\sqrt{\chi}}{1 + \chi} \right),$$

$$\nu = \frac{\sigma + \theta\sqrt{\chi}}{\sqrt{1 + \chi}}, \quad \mu = \frac{\sigma\sqrt{\chi} - \theta}{\sqrt{1 + \chi}}.$$

$\chi = \frac{ac}{bd}$  denotes the cross-ratio, and the evolution of spin graph follows from the construction associated to the spin graph evolution rule.

[Outline](#)
[Introduction](#)
[Cast of characters](#)
[Coordinates on  
Super-Teichmüller  
space](#)
 [\$\mathcal{N} = 2\$   
Super-Teichmüller  
theory](#)
[Further work](#)
[Open problems](#)



- These coordinates are natural in the sense that if  $\varphi \in MC(F)$  has induced action  $\tilde{\varphi}$  on  $\tilde{\Gamma} \in ST(F)$ , then  $\tilde{\varphi}(\tilde{\Gamma})$  is determined by the orientation and coordinates on edges and vertices of  $\varphi(\tau)$  induced by  $\varphi$  from the orientation  $\omega$ , the  $\lambda$ -lengths and  $\mu$ -invariants on  $\tau$ .

- There is an even 2-form on  $ST(F)$  which is invariant under super Ptolemy transformations, namely,

$$\omega = \sum_v d \log a \wedge d \log b + d \log b \wedge d \log c + d \log c \wedge d \log a - (d\theta)^2,$$

where the sum is over all vertices  $v$  of  $\tau$  where the consecutive half edges incident on  $v$  in clockwise order have induced  $\lambda$ -lengths  $a, b, c$  and  $\theta$  is the  $\mu$ -invariant of  $v$ .

- Coordinates on  $ST(F)$ :

Take instead of  $\lambda$ -lengths shear coordinates  $z_e = \log \left( \frac{ac}{bd} \right)$  for every edge  $e$ , which are subject to linear relation: the sum of all  $z_e$  adjacent to a given vertex = 0.

- These coordinates are natural in the sense that if  $\varphi \in MC(F)$  has induced action  $\tilde{\varphi}$  on  $\tilde{\Gamma} \in ST(F)$ , then  $\tilde{\varphi}(\tilde{\Gamma})$  is determined by the orientation and coordinates on edges and vertices of  $\varphi(\tau)$  induced by  $\varphi$  from the orientation  $\omega$ , the  $\lambda$ -lengths and  $\mu$ -invariants on  $\tau$ .

- There is an even 2-form on  $ST(F)$  which is invariant under super Ptolemy transformations, namely,

$$\omega = \sum_v d \log a \wedge d \log b + d \log b \wedge d \log c + d \log c \wedge d \log a - (d\theta)^2,$$

where the sum is over all vertices  $v$  of  $\tau$  where the consecutive half edges incident on  $v$  in clockwise order have induced  $\lambda$ -lengths  $a, b, c$  and  $\theta$  is the  $\mu$ -invariant of  $v$ .

- Coordinates on  $ST(F)$ :

Take instead of  $\lambda$ -lengths shear coordinates  $z_e = \log \left( \frac{ac}{bd} \right)$  for every edge  $e$ , which are subject to linear relation: the sum of all  $z_e$  adjacent to a given vertex = 0.

- These coordinates are natural in the sense that if  $\varphi \in MC(F)$  has induced action  $\tilde{\varphi}$  on  $\tilde{\Gamma} \in ST(F)$ , then  $\tilde{\varphi}(\tilde{\Gamma})$  is determined by the orientation and coordinates on edges and vertices of  $\varphi(\tau)$  induced by  $\varphi$  from the orientation  $\omega$ , the  $\lambda$ -lengths and  $\mu$ -invariants on  $\tau$ .

- There is an even 2-form on  $ST(F)$  which is invariant under super Ptolemy transformations, namely,

$$\omega = \sum_v d \log a \wedge d \log b + d \log b \wedge d \log c + d \log c \wedge d \log a - (d\theta)^2,$$

where the sum is over all vertices  $v$  of  $\tau$  where the consecutive half edges incident on  $v$  in clockwise order have induced  $\lambda$ -lengths  $a, b, c$  and  $\theta$  is the  $\mu$ -invariant of  $v$ .

- Coordinates on  $ST(F)$ :

Take instead of  $\lambda$ -lengths shear coordinates  $z_e = \log \left( \frac{ac}{bd} \right)$  for every edge  $e$ , which are subject to linear relation: the sum of all  $z_e$  adjacent to a given vertex = 0.

# Sketch of construction via hyperbolic supergeometry

XIXth century perspective on hyperbolic (super)geometry:

$OSp(1|2)$  acts on super-Minkowski space  $\mathbb{R}^{2,1|2}$  (in the bosonic case  
 $PSL(2, \mathbb{R})$  acts on  $\mathbb{R}^{2,1}$ ).

If  $A = (x_1, x_2, y, \phi, \theta)$  and  $A' = (x'_1, x'_2, y', \phi', \theta')$  in  $\mathbb{R}^{2,1|2}$ , the pairing is:

$$\langle A, A' \rangle = \frac{1}{2}(x_1 x'_2 + x'_1 x_2) - yy' + \phi\theta' + \phi'\theta.$$

Two surfaces of special importance for us are

- ▶ Superhyperboloid  $\mathbb{H}$  consisting of points  $A \in \mathbb{R}^{2,1|2}$  satisfying the condition  $\langle A, A \rangle = 1$
- ▶ Positive super light cone  $L^+$  consisting of points  $B \in \mathbb{R}^{2,1|2}$  satisfying  $\langle B, B \rangle = 0$ ,

where  $x_1^\#, x_2^\# \geq 0$ .

There is an equivariant projection from  $\mathbb{H}$  on the  $\mathcal{N} = 1$  super upper half-plane  $H^+$ .

XIXth century perspective on hyperbolic (super)geometry:

$OSp(1|2)$  acts on super-Minkowski space  $\mathbb{R}^{2,1|2}$  (in the bosonic case  
 $PSL(2, \mathbb{R})$  acts on  $\mathbb{R}^{2,1}$ ).

If  $A = (x_1, x_2, y, \phi, \theta)$  and  $A' = (x'_1, x'_2, y', \phi', \theta')$  in  $\mathbb{R}^{2,1|2}$ , the pairing is:

$$\langle A, A' \rangle = \frac{1}{2}(x_1 x'_2 + x'_1 x_2) - yy' + \phi\theta' + \phi'\theta.$$

Two surfaces of special importance for us are

- ▶ Superhyperboloid  $\mathbb{H}$  consisting of points  $A \in \mathbb{R}^{2,1|2}$  satisfying the condition  $\langle A, A \rangle = 1$
- ▶ Positive super light cone  $L^+$  consisting of points  $B \in \mathbb{R}^{2,1|2}$  satisfying  $\langle B, B \rangle = 0$ ,

where  $x_1^\#, x_2^\# \geq 0$ .

There is an equivariant projection from  $\mathbb{H}$  on the  $\mathcal{N} = 1$  super upper half-plane  $H^+$ .

[Outline](#)[Introduction](#)[Cast of characters](#)[Coordinates on  
Super-Teichmüller  
space](#) [\$\mathcal{N} = 2\$   
Super-Teichmüller  
theory](#)[Further work](#)[Open problems](#)

XIXth century perspective on hyperbolic (super)geometry:

$OSp(1|2)$  acts on super-Minkowski space  $\mathbb{R}^{2,1|2}$  (in the bosonic case  
 $PSL(2, \mathbb{R})$  acts on  $\mathbb{R}^{2,1}$ ).

If  $A = (x_1, x_2, y, \phi, \theta)$  and  $A' = (x'_1, x'_2, y', \phi', \theta')$  in  $\mathbb{R}^{2,1|2}$ , the pairing is:

$$\langle A, A' \rangle = \frac{1}{2}(x_1 x'_2 + x'_1 x_2) - yy' + \phi\theta' + \phi'\theta.$$

Two surfaces of special importance for us are

- ▶ Superhyperboloid  $\mathbb{H}$  consisting of points  $A \in \mathbb{R}^{2,1|2}$  satisfying the condition  $\langle A, A \rangle = 1$
- ▶ Positive super light cone  $L^+$  consisting of points  $B \in \mathbb{R}^{2,1|2}$  satisfying  $\langle B, B \rangle = 0$ ,

where  $x_1^\#, x_2^\# \geq 0$ .

There is an equivariant projection from  $\mathbb{H}$  on the  $\mathcal{N} = 1$  super upper half-plane  $H^+$ .

[Outline](#)[Introduction](#)[Cast of characters](#)[Coordinates on  
Super-Teichmüller  
space](#) [\$\mathcal{N} = 2\$   
Super-Teichmüller  
theory](#)[Further work](#)[Open problems](#)

XIXth century perspective on hyperbolic (super)geometry:

$OSp(1|2)$  acts on super-Minkowski space  $\mathbb{R}^{2,1|2}$  (in the bosonic case  
 $PSL(2, \mathbb{R})$  acts on  $\mathbb{R}^{2,1}$ ).

If  $A = (x_1, x_2, y, \phi, \theta)$  and  $A' = (x'_1, x'_2, y', \phi', \theta')$  in  $\mathbb{R}^{2,1|2}$ , the pairing is:

$$\langle A, A' \rangle = \frac{1}{2}(x_1 x'_2 + x'_1 x_2) - yy' + \phi\theta' + \phi'\theta.$$

Two surfaces of special importance for us are

- ▶ Superhyperboloid  $\mathbb{H}$  consisting of points  $A \in \mathbb{R}^{2,1|2}$  satisfying the condition  $\langle A, A \rangle = 1$
- ▶ Positive super light cone  $L^+$  consisting of points  $B \in \mathbb{R}^{2,1|2}$  satisfying  $\langle B, B \rangle = 0$ ,

where  $x_1^\#, x_2^\# \geq 0$ .

There is an equivariant projection from  $\mathbb{H}$  on the  $\mathcal{N} = 1$  super upper half-plane  $H^+$ .

[Outline](#)[Introduction](#)[Cast of characters](#)[Coordinates on  
Super-Teichmüller  
space](#) [\$\mathcal{N} = 2\$   
Super-Teichmüller  
theory](#)[Further work](#)[Open problems](#)

XIXth century perspective on hyperbolic (super)geometry:

$OSp(1|2)$  acts on super-Minkowski space  $\mathbb{R}^{2,1|2}$  (in the bosonic case  
 $PSL(2, \mathbb{R})$  acts on  $\mathbb{R}^{2,1}$ ).

If  $A = (x_1, x_2, y, \phi, \theta)$  and  $A' = (x'_1, x'_2, y', \phi', \theta')$  in  $\mathbb{R}^{2,1|2}$ , the pairing is:

$$\langle A, A' \rangle = \frac{1}{2}(x_1 x'_2 + x'_1 x_2) - yy' + \phi\theta' + \phi'\theta.$$

Two surfaces of special importance for us are

- ▶ Superhyperboloid  $\mathbb{H}$  consisting of points  $A \in \mathbb{R}^{2,1|2}$  satisfying the condition  $\langle A, A \rangle = 1$
- ▶ Positive super light cone  $L^+$  consisting of points  $B \in \mathbb{R}^{2,1|2}$  satisfying  $\langle B, B \rangle = 0$ ,

where  $x_1^\#, x_2^\# \geq 0$ .

There is an equivariant projection from  $\mathbb{H}$  on the  $\mathcal{N} = 1$  super upper half-plane  $H^+$ .

[Outline](#)[Introduction](#)[Cast of characters](#)[Coordinates on  
Super-Teichmüller  
space](#) [\$\mathcal{N} = 2\$   
Super-Teichmüller  
theory](#)[Further work](#)[Open problems](#)



$OSp(1|2)$  does not act transitively on  $L^+$ :

The space of orbits is labelled by odd variable up to a sign.

We pick an orbit of the vector  $(1, 0, 0, 0, 0)$  and denote it  $L_0^+$ .

There is an equivariant projection from  $L_0^+$  to  $\mathbb{R}^{1|1} = \partial H^+$ .

Goal: Construction of the  $\pi_1$ -equivariant lift for all the data from the universal cover  $\tilde{F}$ , associated to its triangulation to  $L_0^+$ .

Such equivariant lift gives the representation of  $\pi_1$  in  $OSp(1|2)$ .

Outline

Introduction

Cast of characters

Coordinates on  
Super-Teichmüller  
space

$\mathcal{N} = 2$   
Super-Teichmüller  
theory

Further work

Open problems

$OSp(1|2)$  does not act transitively on  $L^+$ :

The space of orbits is labelled by odd variable up to a sign.

We pick an orbit of the vector  $(1, 0, 0, 0, 0)$  and denote it  $L_0^+$ .

There is an equivariant projection from  $L_0^+$  to  $\mathbb{R}^{1|1} = \partial H^+$ .

Goal: Construction of the  $\pi_1$ -equivariant lift for all the data from the universal cover  $\tilde{F}$ , associated to its triangulation to  $L_0^+$ .

Such equivariant lift gives the representation of  $\pi_1$  in  $OSp(1|2)$ .

$OSp(1|2)$  does not act transitively on  $L^+$ :

The space of orbits is labelled by odd variable up to a sign.

We pick an orbit of the vector  $(1, 0, 0, 0, 0)$  and denote it  $L_0^+$ .

There is an equivariant projection from  $L_0^+$  to  $\mathbb{R}^{1|1} = \partial H^+$ .

Goal: Construction of the  $\pi_1$ -equivariant lift for all the data from the universal cover  $\tilde{F}$ , associated to its triangulation to  $L_0^+$ .

Such equivariant lift gives the representation of  $\pi_1$  in  $OSp(1|2)$ .

$OSp(1|2)$  does not act transitively on  $L^+$ :

The space of orbits is labelled by odd variable up to a sign.

We pick an orbit of the vector  $(1, 0, 0, 0, 0)$  and denote it  $L_0^+$ .

There is an equivariant projection from  $L_0^+$  to  $\mathbb{R}^{1|1} = \partial H^+$ .

Goal: Construction of the  $\pi_1$ -equivariant lift for all the data from the universal cover  $\tilde{F}$ , associated to its triangulation to  $L_0^+$ .

Such equivariant lift gives the representation of  $\pi_1$  in  $OSp(1|2)$ .

- There is a unique  $OSp(1|2)$ -invariant of two linearly independent vectors  $A, B \in L_0^+$ , and it is given by the pairing  $\langle A, B \rangle$ , the square root of which we will call  $\lambda$ -length.

Let  $\zeta^b \zeta^e \zeta^a$  be a positive triple in  $L_0^+$ . Then there is  $g \in OSp(1|2)$ , which is unique up to composition with the fermionic reflection, and unique even  $r, s, t$ , which have positive bodies, and odd  $\theta$  so that

$$g \cdot \zeta^e = t(1, 1, 1, \theta, \theta), \quad g \cdot \zeta^b = r(0, 1, 0, 0, 0), \quad g \cdot \zeta^a = s(1, 0, 0, 0, 0).$$

- The moduli space of  $OSp(1|2)$ -orbits of positive triples in the light cone is given by  $(a, b, e, \theta) \in \mathbb{R}_+^{3|1} / \mathbb{Z}_2$ , where  $\mathbb{Z}_2$  acts by fermionic reflection.

On the superline  $\mathbb{R}^{1|1}$  the parameter  $\theta$  is known as *Manin invariant*.

[Outline](#)[Introduction](#)[Cast of characters](#)[Coordinates on  
Super-Teichmüller  
space](#) [\$\mathcal{N} = 2\$   
Super-Teichmüller  
theory](#)[Further work](#)[Open problems](#)

- There is a unique  $OSp(1|2)$ -invariant of two linearly independent vectors  $A, B \in L_0^+$ , and it is given by the pairing  $\langle A, B \rangle$ , the square root of which we will call  $\lambda$ -length.

Let  $\zeta^b \zeta^e \zeta^a$  be a positive triple in  $L_0^+$ . Then there is  $g \in OSp(1|2)$ , which is unique up to composition with the fermionic reflection, and unique even  $r, s, t$ , which have positive bodies, and odd  $\theta$  so that

$$g \cdot \zeta^e = t(1, 1, 1, \theta, \theta), \quad g \cdot \zeta^b = r(0, 1, 0, 0, 0), \quad g \cdot \zeta^a = s(1, 0, 0, 0, 0).$$

- The moduli space of  $OSp(1|2)$ -orbits of positive triples in the light cone is given by  $(a, b, e, \theta) \in \mathbb{R}_+^{3|1} / \mathbb{Z}_2$ , where  $\mathbb{Z}_2$  acts by fermionic reflection.

On the superline  $\mathbb{R}^{1|1}$  the parameter  $\theta$  is known as *Manin invariant*.

[Outline](#)[Introduction](#)[Cast of characters](#)[Coordinates on  
Super-Teichmüller  
space](#) [\$\mathcal{N} = 2\$   
Super-Teichmüller  
theory](#)[Further work](#)[Open problems](#)

- There is a unique  $OSp(1|2)$ -invariant of two linearly independent vectors  $A, B \in L_0^+$ , and it is given by the pairing  $\langle A, B \rangle$ , the square root of which we will call  $\lambda$ -length.

Let  $\zeta^b \zeta^e \zeta^a$  be a positive triple in  $L_0^+$ . Then there is  $g \in OSp(1|2)$ , which is unique up to composition with the fermionic reflection, and unique even  $r, s, t$ , which have positive bodies, and odd  $\theta$  so that

$$g \cdot \zeta^e = t(1, 1, 1, \theta, \theta), \quad g \cdot \zeta^b = r(0, 1, 0, 0, 0), \quad g \cdot \zeta^a = s(1, 0, 0, 0, 0).$$

- The moduli space of  $OSp(1|2)$ -orbits of positive triples in the light cone is given by  $(a, b, e, \theta) \in \mathbb{R}_+^{3|1} / \mathbb{Z}_2$ , where  $\mathbb{Z}_2$  acts by fermionic reflection.

On the superline  $\mathbb{R}^{1|1}$  the parameter  $\theta$  is known as *Manin invariant*.

[Outline](#)[Introduction](#)[Cast of characters](#)[Coordinates on  
Super-Teichmüller  
space](#) [\$\mathcal{N} = 2\$   
Super-Teichmüller  
theory](#)[Further work](#)[Open problems](#)

- There is a unique  $OSp(1|2)$ -invariant of two linearly independent vectors  $A, B \in L_0^+$ , and it is given by the pairing  $\langle A, B \rangle$ , the square root of which we will call  $\lambda$ -length.

Let  $\zeta^b \zeta^e \zeta^a$  be a positive triple in  $L_0^+$ . Then there is  $g \in OSp(1|2)$ , which is unique up to composition with the fermionic reflection, and unique even  $r, s, t$ , which have positive bodies, and odd  $\theta$  so that

$$g \cdot \zeta^e = t(1, 1, 1, \theta, \theta), \quad g \cdot \zeta^b = r(0, 1, 0, 0, 0), \quad g \cdot \zeta^a = s(1, 0, 0, 0, 0).$$

- The moduli space of  $OSp(1|2)$ -orbits of positive triples in the light cone is given by  $(a, b, e, \theta) \in \mathbb{R}_+^{3|1} / \mathbb{Z}_2$ , where  $\mathbb{Z}_2$  acts by fermionic reflection.

On the superline  $\mathbb{R}^{1|1}$  the parameter  $\theta$  is known as *Manin invariant*.


[Outline](#)[Introduction](#)[Cast of characters](#)[Coordinates on  
Super-Teichmüller  
space](#) [\$\mathcal{N} = 2\$   
Super-Teichmüller  
theory](#)[Further work](#)[Open problems](#)



# Orbits of 4 points in $L_0^+$ : basic calculation

Suppose points  $A, B, C$  are put in the standard position.

The 4th point  $D$ , so that two new  $\lambda$ - lengths are  $c, d$ .



Fixing the sign of  $\theta$ , we fix the sign of Manin invariant  $\sigma$  in terms of coordinates of  $D$ .

Important observation: if we turn the picture upside down, then

$$(\theta, \sigma) \rightarrow (\sigma, -\theta)$$

Outline

Introduction

Cast of characters

Coordinates on  
Super-Teichmüller  
space

$\mathcal{N} = 2$   
Super-Teichmüller  
theory


Further work

Open problems

# Orbits of 4 points in $L_0^+$ : basic calculation

Suppose points  $A, B, C$  are put in the standard position.

The 4th point  $D$ , so that two new  $\lambda$ - lengths are  $c, d$ .



Fixing the sign of  $\theta$ , we fix the sign of Manin invariant  $\sigma$  in terms of coordinates of  $D$ .

Important observation: if we turn the picture upside down, then

$$(\theta, \sigma) \rightarrow (\sigma, -\theta)$$

Outline

Introduction

Cast of characters

Coordinates on  
Super-Teichmüller  
space

$\mathcal{N} = 2$   
Super-Teichmüller  
theory


Further work

Open problems

# Orbits of 4 points in $L_0^+$ : basic calculation

Suppose points  $A, B, C$  are put in the standard position.

The 4th point  $D$ , so that two new  $\lambda$ - lengths are  $c, d$ .



Fixing the sign of  $\theta$ , we fix the sign of Manin invariant  $\sigma$  in terms of coordinates of  $D$ .

Important observation: if we turn the picture upside down, then

$$(\theta, \sigma) \rightarrow (\sigma, -\theta)$$

# The lift of ideal triangulation to super-Minkowski space

Denote:

- ▶  $\Delta$  is ideal triangulation of  $F$ ,  $\tilde{\Delta}$  is ideal triangulation of the universal cover  $\tilde{F}$
- ▶  $\Delta_\infty$  ( $\tilde{\Delta}_\infty$ )-collection of ideal points of  $F$  ( $\tilde{F}$ ).

Consider  $\Delta$  together with:

- the orientation on the fatgraph  $\tau(\Delta)$ ,
- coordinate system  $\tilde{C}(F, \Delta)$ , i.e.
  - positive even coordinate for every edge
  - odd coordinate for every triangle

We call coordinate vectors  $\vec{c}, \vec{c}'$  equivalent if they are identical up to overall reflection of sign of odd coordinates.

Let  $C(F, \Delta) \equiv \tilde{C}(F, \Delta) / \sim$ . This implies that

$$C(F, \Delta) \simeq \mathbb{R}_+^{6g+3s-6|4g+2s-4} / \mathbb{Z}_2$$

Outline

Introduction

Cast of characters

Coordinates on  
Super-Teichmüller  
space

$\mathcal{N} = 2$   
Super-Teichmüller  
theory

Further work

Open problems

# The lift of ideal triangulation to super-Minkowski space

Denote:

- ▶  $\Delta$  is ideal triangulation of  $F$ ,  $\tilde{\Delta}$  is ideal triangulation of the universal cover  $\tilde{F}$
- ▶  $\Delta_\infty$  ( $\tilde{\Delta}_\infty$ )-collection of ideal points of  $F$  ( $\tilde{F}$ ).

Consider  $\Delta$  together with:

- the orientation on the fatgraph  $\tau(\Delta)$ ,

coordinate system  $\tilde{C}(F, \Delta)$ , i.e.

- positive even coordinate for every edge
- odd coordinate for every triangle

We call coordinate vectors  $\vec{c}$ ,  $\vec{c}'$  equivalent if they are identical up to overall reflection of sign of odd coordinates.

Let  $C(F, \Delta) \equiv \tilde{C}(F, \Delta) / \sim$ . This implies that

$$C(F, \Delta) \simeq \mathbb{R}_+^{6g+3s-6|4g+2s-4} / \mathbb{Z}_2$$

Outline

Introduction

Cast of characters

Coordinates on  
Super-Teichmüller  
space

$\mathcal{N} = 2$   
Super-Teichmüller  
theory

Further work

Open problems

# The lift of ideal triangulation to super-Minkowski space

Denote:

- ▶  $\Delta$  is ideal triangulation of  $F$ ,  $\tilde{\Delta}$  is ideal triangulation of the universal cover  $\tilde{F}$
- ▶  $\Delta_\infty$  ( $\tilde{\Delta}_\infty$ )-collection of ideal points of  $F$  ( $\tilde{F}$ ).

Consider  $\Delta$  together with:

- the orientation on the fatgraph  $\tau(\Delta)$ ,
- coordinate system  $\tilde{C}(F, \Delta)$ , i.e.
- positive even coordinate for every edge
- odd coordinate for every triangle

We call coordinate vectors  $\vec{c}, \vec{c}'$  equivalent if they are identical up to overall reflection of sign of odd coordinates.

Let  $C(F, \Delta) \equiv \tilde{C}(F, \Delta) / \sim$ . This implies that

$$C(F, \Delta) \simeq \mathbb{R}_+^{6g+3s-6} |4g+2s-4| / \mathbb{Z}_2$$

Outline

Introduction

Cast of characters

Coordinates on  
Super-Teichmüller  
space

$\mathcal{N} = 2$   
Super-Teichmüller  
theory

Further work

Open problems

# The lift of ideal triangulation to super-Minkowski space

Denote:

- ▶  $\Delta$  is ideal triangulation of  $F$ ,  $\tilde{\Delta}$  is ideal triangulation of the universal cover  $\tilde{F}$
- ▶  $\Delta_\infty$  ( $\tilde{\Delta}_\infty$ )-collection of ideal points of  $F$  ( $\tilde{F}$ ).

Consider  $\Delta$  together with:

- the orientation on the fatgraph  $\tau(\Delta)$ ,
- coordinate system  $\tilde{C}(F, \Delta)$ , i.e.
- positive even coordinate for every edge
- odd coordinate for every triangle

We call coordinate vectors  $\vec{c}$ ,  $\vec{c}'$  equivalent if they are identical up to overall reflection of sign of odd coordinates.

Let  $C(F, \Delta) \equiv \tilde{C}(F, \Delta) / \sim$ . This implies that

$$C(F, \Delta) \simeq \mathbb{R}_+^{6g+3s-6} |4g+2s-4| / \mathbb{Z}_2$$

Outline

Introduction

Cast of characters

Coordinates on  
Super-Teichmüller  
space

$\mathcal{N} = 2$   
Super-Teichmüller  
theory


Further work

Open problems

Then there exists a lift for each  $\vec{c} \in \ell : \check{\Delta}_\infty \rightarrow L_0^+$ , with the property:

for every quadrilateral  $ABCD$ , if the arrow is pointing from  $\sigma$  to  $\theta$  then the lift is given by the picture from the previous slide up to post-composition with the element of  $OSP(1|2)$ .

The construction of  $\ell$  can be done in a recursive way:



Such lift is unique up to post-composition with  $OSP(1|2)$  group element and it is  $\pi_1$ -equivariant. This allows us to construct representation of  $\pi_1$  in  $OSP(1|2)$ , based on the provided data.

Outline

Introduction

Cast of characters

Coordinates on  
Super-Teichmüller  
space

$\mathcal{N} = 2$   
Super-Teichmüller  
theory

Further work


Open problems



Then there exists a lift for each  $\vec{c} \in \ell : \tilde{\Delta}_\infty \rightarrow L_0^+$ , with the property:

for every quadrilateral  $ABCD$ , if the arrow is pointing from  $\sigma$  to  $\theta$  then the lift is given by the picture from the previous slide up to post-composition with the element of  $OSp(1|2)$ .

The construction of  $\ell$  can be done in a recursive way:



Such lift is unique up to post-composition with  $OSp(1|2)$  group element and it is  $\pi_1$ -equivariant. This allows us to construct representation of  $\pi_1$  in  $OSP(1|2)$ , based on the provided data.

Outline

Introduction

Cast of characters

Coordinates on  
Super-Teichmüller  
space

$\mathcal{N} = 2$   
Super-Teichmüller  
theory


Further work

Open problems

Then there exists a lift for each  $\vec{c} \in \ell : \tilde{\Delta}_\infty \rightarrow L_0^+$ , with the property:

for every quadrilateral  $ABCD$ , if the arrow is pointing from  $\sigma$  to  $\theta$  then the lift is given by the picture from the previous slide up to post-composition with the element of  $OSP(1|2)$ .

The construction of  $\ell$  can be done in a recursive way:




Such lift is unique up to post-composition with  $OSP(1|2)$  group element and it is  $\pi_1$ -equivariant. This allows us to construct representation of  $\pi_1$  in  $OSP(1|2)$ , based on the provided data.

Then there exists a lift for each  $\vec{c} \in \ell : \tilde{\Delta}_\infty \rightarrow L_0^+$ , with the property:

for every quadrilateral  $ABCD$ , if the arrow is pointing from  $\sigma$  to  $\theta$  then the lift is given by the picture from the previous slide up to post-composition with the element of  $OSP(1|2)$ .

The construction of  $\ell$  can be done in a recursive way:



Such lift is unique up to post-composition with  $OSP(1|2)$  group element and it is  $\pi_1$ -equivariant. This allows us to construct representation of  $\pi_1$  in  $OSP(1|2)$ , based on the provided data.

# Theorem

Fix  $F, \Delta, \tau(\Delta)$  as before. Let  $\omega$  be an orientation, corresponding to a specified spin structure  $s$  of  $F$ . Given a coordinate vector  $\vec{c} \in \tilde{C}(F, \Delta)$ , there exists a map called the lift,

$$\ell_\omega : \tilde{\Delta}_\infty \rightarrow L_0^+$$

which is uniquely determined up to post-composition by  $OSp(1|2)$  under admissibility conditions discussed above, and only depends on the equivalent classes  $C(F, \Delta)$  of the coordinates.

There is a representation  $\hat{\rho} : \pi_1 := \pi_1(F) \rightarrow OSp(1|2)$ , uniquely determined up to conjugacy by an element of  $OSp(1|2)$  such that

- (1)  $\ell$  is  $\pi_1$ -equivariant, i.e.  $\hat{\rho}(\gamma)(\ell(a)) = \ell(\gamma(a))$  for each  $\gamma \in \pi_1$  and  $a \in \tilde{\Delta}_\infty$ ;
- (2)  $\hat{\rho}$  is a super-Fuchsian representation, i.e. the natural projection

$$\rho : \pi_1 \xrightarrow{\hat{\rho}} OSp(1|2) \rightarrow SL(2, \mathbb{R}) \rightarrow PSL(2, \mathbb{R})$$

is a Fuchsian representation for  $F$ ;

- (3) the space of all lifts  $\tilde{\rho} : \pi_1 \xrightarrow{\hat{\rho}} OSp(1|2) \rightarrow SL(2, \mathbb{R})$  is in one-to-one correspondence with the spin structures  $s$  on  $F$ .

# Theorem

Fix  $F, \Delta, \tau(\Delta)$  as before. Let  $\omega$  be an orientation, corresponding to a specified spin structure  $s$  of  $F$ . Given a coordinate vector  $\vec{c} \in \tilde{C}(F, \Delta)$ , there exists a map called the lift,

$$\ell_\omega : \tilde{\Delta}_\infty \rightarrow L_0^+$$

which is uniquely determined up to post-composition by  $OSp(1|2)$  under admissibility conditions discussed above, and only depends on the equivalent classes  $C(F, \Delta)$  of the coordinates.

There is a representation  $\hat{\rho} : \pi_1 := \pi_1(F) \rightarrow OSp(1|2)$ , uniquely determined up to conjugacy by an element of  $OSp(1|2)$  such that

- (1)  $\ell$  is  $\pi_1$ -equivariant, i.e.  $\hat{\rho}(\gamma)(\ell(a)) = \ell(\gamma(a))$  for each  $\gamma \in \pi_1$  and  $a \in \tilde{\Delta}_\infty$ ;
- (2)  $\hat{\rho}$  is a super-Fuchsian representation, i.e. the natural projection

$$\rho : \pi_1 \xrightarrow{\hat{\rho}} OSp(1|2) \rightarrow SL(2, \mathbb{R}) \rightarrow PSL(2, \mathbb{R})$$

is a Fuchsian representation for  $F$ ;

- (3) the space of all lifts  $\tilde{\rho} : \pi_1 \xrightarrow{\hat{\rho}} OSp(1|2) \rightarrow SL(2, \mathbb{R})$  is in one-to-one correspondence with the spin structures  $s$  on  $F$ .

# Theorem

Fix  $F, \Delta, \tau(\Delta)$  as before. Let  $\omega$  be an orientation, corresponding to a specified spin structure  $s$  of  $F$ . Given a coordinate vector  $\vec{c} \in \tilde{C}(F, \Delta)$ , there exists a map called the lift,

$$\ell_\omega : \tilde{\Delta}_\infty \rightarrow L_0^+$$

which is uniquely determined up to post-composition by  $OSp(1|2)$  under admissibility conditions discussed above, and only depends on the equivalent classes  $C(F, \Delta)$  of the coordinates.

There is a representation  $\hat{\rho} : \pi_1 := \pi_1(F) \rightarrow OSp(1|2)$ , uniquely determined up to conjugacy by an element of  $OSp(1|2)$  such that

(1)  $\ell$  is  $\pi_1$ -equivariant, i.e.  $\hat{\rho}(\gamma)(\ell(a)) = \ell(\gamma(a))$  for each  $\gamma \in \pi_1$  and  $a \in \tilde{\Delta}_\infty$ ;

(2)  $\hat{\rho}$  is a super-Fuchsian representation, i.e. the natural projection

$$\rho : \pi_1 \xrightarrow{\hat{\rho}} OSp(1|2) \rightarrow SL(2, \mathbb{R}) \rightarrow PSL(2, \mathbb{R})$$

is a Fuchsian representation for  $F$ ;

(3) the space of all lifts  $\tilde{\rho} : \pi_1 \xrightarrow{\hat{\rho}} OSp(1|2) \rightarrow SL(2, \mathbb{R})$  is in one-to-one correspondence with the spin structures  $s$  on  $F$ .

Fix  $F, \Delta, \tau(\Delta)$  as before. Let  $\omega$  be an orientation, corresponding to a specified spin structure  $s$  of  $F$ . Given a coordinate vector  $\vec{c} \in \tilde{C}(F, \Delta)$ , there exists a map called the lift,

$$\ell_\omega : \tilde{\Delta}_\infty \rightarrow L_0^+$$

which is uniquely determined up to post-composition by  $OSp(1|2)$  under admissibility conditions discussed above, and only depends on the equivalent classes  $C(F, \Delta)$  of the coordinates.

There is a representation  $\hat{\rho} : \pi_1 := \pi_1(F) \rightarrow OSp(1|2)$ , uniquely determined up to conjugacy by an element of  $OSp(1|2)$  such that

- (1)  $\ell$  is  $\pi_1$ -equivariant, i.e.  $\hat{\rho}(\gamma)(\ell(a)) = \ell(\gamma(a))$  for each  $\gamma \in \pi_1$  and  $a \in \tilde{\Delta}_\infty$ ;
- (2)  $\hat{\rho}$  is a super-Fuchsian representation, i.e. the natural projection

$$\rho : \pi_1 \xrightarrow{\hat{\rho}} OSp(1|2) \rightarrow SL(2, \mathbb{R}) \rightarrow PSL(2, \mathbb{R})$$

is a Fuchsian representation for  $F$ ;

- (3) the space of all lifts  $\tilde{\rho} : \pi_1 \xrightarrow{\hat{\rho}} OSp(1|2) \rightarrow SL(2, \mathbb{R})$  is in one-to-one correspondence with the spin structures  $s$  on  $F$ .

Fix  $F, \Delta, \tau(\Delta)$  as before. Let  $\omega$  be an orientation, corresponding to a specified spin structure  $s$  of  $F$ . Given a coordinate vector  $\vec{c} \in \tilde{C}(F, \Delta)$ , there exists a map called the lift,

$$\ell_\omega : \tilde{\Delta}_\infty \rightarrow L_0^+$$

which is uniquely determined up to post-composition by  $OSp(1|2)$  under admissibility conditions discussed above, and only depends on the equivalent classes  $C(F, \Delta)$  of the coordinates.

There is a representation  $\hat{\rho} : \pi_1 := \pi_1(F) \rightarrow OSp(1|2)$ , uniquely determined up to conjugacy by an element of  $OSp(1|2)$  such that

- (1)  $\ell$  is  $\pi_1$ -equivariant, i.e.  $\hat{\rho}(\gamma)(\ell(a)) = \ell(\gamma(a))$  for each  $\gamma \in \pi_1$  and  $a \in \tilde{\Delta}_\infty$ ;
- (2)  $\hat{\rho}$  is a super-Fuchsian representation, i.e. the natural projection


$$\rho : \pi_1 \xrightarrow{\hat{\rho}} OSp(1|2) \rightarrow SL(2, \mathbb{R}) \rightarrow PSL(2, \mathbb{R})$$

is a Fuchsian representation for  $F$ ;

- (3) the space of all lifts  $\tilde{\rho} : \pi_1 \xrightarrow{\hat{\rho}} OSp(1|2) \rightarrow SL(2, \mathbb{R})$  is in one-to-one correspondence with the spin structures  $s$  on  $F$ .



## The super-Ptolemy transformations



$$ef = (ac + bd) \left( 1 + \frac{\sigma\theta\sqrt{\chi}}{1 + \chi} \right),$$

$$\nu = \frac{\sigma + \theta\sqrt{\chi}}{\sqrt{1 + \chi}}, \quad \mu = \frac{\sigma\sqrt{\chi} - \theta}{\sqrt{1 + \chi}}$$

are the consequence of light cone geometry.

Outline

Introduction

Cast of characters

Coordinates on  
Super-Teichmüller  
space $\mathcal{N} = 2$   
Super-Teichmüller  
theory

Further work

Open problems

The space of all such lifts  $\ell_\omega$  coincides with the decorated super-Teichmüller space  $S\tilde{T}(F) = \mathbb{R}_+^s \times ST(F)$ .

In order to remove the decoration, one can pass to shear coordinates  $z_e = \log\left(\frac{ac}{bd}\right)$ .

It is easy to check that the 2-form

$$\omega = \sum_{\Delta} d \log a \wedge d \log b + d \log b \wedge d \log c + d \log c \wedge d \log a - (d\theta)^2$$

is invariant under the flip transformations. This is a generalization of the formula for Weil-Petersson 2-form.

The space of all such lifts  $\ell_\omega$  coincides with the decorated super-Teichmüller space  $S\tilde{T}(F) = \mathbb{R}_+^s \times ST(F)$ .

In order to remove the decoration, one can pass to shear coordinates  $z_e = \log \left( \frac{ac}{bd} \right)$ .

It is easy to check that the 2-form

$$\omega = \sum_{\Delta} d \log a \wedge d \log b + d \log b \wedge d \log c + d \log c \wedge d \log a - (d\theta)^2$$

is invariant under the flip transformations. This is a generalization of the formula for Weil-Petersson 2-form.

The space of all such lifts  $\ell_\omega$  coincides with the decorated super-Teichmüller space  $S\tilde{T}(F) = \mathbb{R}_+^s \times ST(F)$ .

In order to remove the decoration, one can pass to shear coordinates  $z_e = \log\left(\frac{ac}{bd}\right)$ .

It is easy to check that the 2-form

$$\omega = \sum_{\Delta} d \log a \wedge d \log b + d \log b \wedge d \log c + d \log c \wedge d \log a - (d\theta)^2$$

is invariant under the flip transformations. This is a generalization of the formula for Weil-Petersson 2-form.

The space of all such lifts  $\ell_\omega$  coincides with the decorated super-Teichmüller space  $S\tilde{T}(F) = \mathbb{R}_+^s \times ST(F)$ .

In order to remove the decoration, one can pass to shear coordinates  $z_e = \log\left(\frac{ac}{bd}\right)$ .

It is easy to check that the 2-form

$$\omega = \sum_{\Delta} d \log a \wedge d \log b + d \log b \wedge d \log c + d \log c \wedge d \log a - (d\theta)^2$$

is invariant under the flip transformations. This is a generalization of the formula for Weil-Petersson 2-form.

Further reduction of the decoration:  $S\tilde{T}(F) = \mathbb{R}_+^{6g+3s-6|4g+2s-4}/\mathbb{Z}_2$  is actually an  $\mathbb{R}_+^{(s|n_R)}$ -decoration over physically relevant Teichmüller space.

Here  $n_R$  is the number of Ramond punctures, which means that the small contour  $\gamma$  surrounding the puncture is such that  $q[\gamma] = 1$ , i.e.  $\text{tr}(\tilde{\rho}(\gamma)) > 0$ .

On the level of hyperbolic geometry, the appropriate constraint is that the monodromy group element has to be true parabolic, i.e. to be conjugated to the parabolic element of  $SL(2, \mathbb{R})$  subgroup.

We formulated it in terms of invariant constraints on shear coordinates in:

I. Ip, R. Penner, A. Zeitlin, arXiv:1709.06207, to appear in Comm. Math. Phys.

Outline

Introduction

Cast of characters

Coordinates on  
Super-Teichmüller  
space

$\mathcal{N} = 2$   
Super-Teichmüller  
theory

Further work

Open problems

Further reduction of the decoration:  $S\tilde{T}(F) = \mathbb{R}_+^{6g+3s-6|4g+2s-4} / \mathbb{Z}_2$  is actually an  $\mathbb{R}_+^{(s|n_R)}$ -decoration over physically relevant Teichmüller space.

Here  $n_R$  is the number of Ramond punctures, which means that the small contour  $\gamma$  surrounding the puncture is such that  $q[\gamma] = 1$ , i.e.  $\text{tr}(\tilde{\rho}(\gamma)) > 0$ .

On the level of hyperbolic geometry, the appropriate constraint is that the monodromy group element has to be true parabolic, i.e. to be conjugated to the parabolic element of  $SL(2, \mathbb{R})$  subgroup.

We formulated it in terms of invariant constraints on shear coordinates in:

I. Ip, R. Penner, A. Zeitlin, arXiv:1709.06207, to appear in Comm. Math. Phys.

[Outline](#)[Introduction](#)[Cast of characters](#)[Coordinates on Super-Teichmüller space](#) [\$\mathcal{N} = 2\$  Super-Teichmüller theory](#)[Further work](#)[Open problems](#)

Further reduction of the decoration:  $S\tilde{T}(F) = \mathbb{R}_+^{6g+3s-6|4g+2s-4} / \mathbb{Z}_2$  is actually an  $\mathbb{R}_+^{(s|n_R)}$ -decoration over physically relevant Teichmüller space.

Here  $n_R$  is the number of Ramond punctures, which means that the small contour  $\gamma$  surrounding the puncture is such that  $q[\gamma] = 1$ , i.e.  $tr(\tilde{\rho}(\gamma)) > 0$ .

On the level of hyperbolic geometry, the appropriate constraint is that the monodromy group element has to be true parabolic, i.e. to be conjugated to the parabolic element of  $SL(2, \mathbb{R})$  subgroup.

We formulated it in terms of invariant constraints on shear coordinates in:

I. Ip, R. Penner, A. Zeitlin, arXiv:1709.06207, to appear in Comm. Math. Phys.

[Outline](#)[Introduction](#)[Cast of characters](#)[Coordinates on Super-Teichmüller space](#) [\$\mathcal{N} = 2\$  Super-Teichmüller theory](#)[Further work](#)[Open problems](#)



Further reduction of the decoration:  $S\tilde{T}(F) = \mathbb{R}_+^{6g+3s-6|4g+2s-4} / \mathbb{Z}_2$  is actually an  $\mathbb{R}_+^{(s|n_R)}$ -decoration over physically relevant Teichmüller space.

Here  $n_R$  is the number of Ramond punctures, which means that the small contour  $\gamma$  surrounding the puncture is such that  $q[\gamma] = 1$ , i.e.  $\text{tr}(\tilde{\rho}(\gamma)) > 0$ .

On the level of hyperbolic geometry, the appropriate constraint is that the monodromy group element has to be true parabolic, i.e. to be conjugated to the parabolic element of  $SL(2, \mathbb{R})$  subgroup.

We formulated it in terms of invariant constraints on shear coordinates in:

I. Ip, R. Penner, A. Zeitlin, arXiv:1709.06207, to appear in Comm. Math. Phys.

Outline

Introduction

Cast of characters

Coordinates on  
Super-Teichmüller  
space

$\mathcal{N} = 2$   
Super-Teichmüller  
theory

Further work

Open problems

Further reduction of the decoration:  $S\tilde{T}(F) = \mathbb{R}_+^{6g+3s-6|4g+2s-4} / \mathbb{Z}_2$  is actually an  $\mathbb{R}_+^{(s|n_R)}$ -decoration over physically relevant Teichmüller space.

Here  $n_R$  is the number of Ramond punctures, which means that the small contour  $\gamma$  surrounding the puncture is such that  $q[\gamma] = 1$ , i.e.  $\text{tr}(\tilde{\rho}(\gamma)) > 0$ .

On the level of hyperbolic geometry, the appropriate constraint is that the monodromy group element has to be true parabolic, i.e. to be conjugated to the parabolic element of  $SL(2, \mathbb{R})$  subgroup.

We formulated it in terms of invariant constraints on shear coordinates in:

I. Ip, R. Penner, A. Zeitlin, arXiv:1709.06207, to appear in Comm. Math. Phys.

[Outline](#)[Introduction](#)[Cast of characters](#)[Coordinates on Super-Teichmüller space](#) [\$\mathcal{N} = 2\$  Super-Teichmüller theory](#)[Further work](#)[Open problems](#)

# $\mathcal{N} = 2$ super-Teichmüller theory: prerequisites

$\mathcal{N} = 2$  super-Teichmüller space is related to  $OSP(2|2)$  supergroup of rank 2.

It is more useful to work with its  $3 \times 3$  incarnation, which is isomorphic to  $\Psi \ltimes SL(1|2)_0$ , where  $\Psi$  is a certain automorphism of the Lie algebra  $\mathfrak{sl}(1|2) \simeq \mathfrak{osp}(2|2)$ .

$SL(1|2)_0$  is a supergroup, consisting of supermatrices

$$g = \begin{pmatrix} a & b & \alpha \\ c & d & \beta \\ \gamma & \delta & f \end{pmatrix}$$

such that  $f > 0$  and their Berezinian = 1.

This group acts on the space  $\mathbb{C}^{1|2}$  as superconformal fractional-linear transformations.

As before,  $\mathcal{N} = 2$  super-Fuchsian groups are the ones whose projections

$$\pi_1 \rightarrow OSP(2|2) \rightarrow GL^+(2, \mathbb{R}) \rightarrow SL(2, \mathbb{R}) \rightarrow PSL(2, \mathbb{R})$$

are Fuchsian.

# $\mathcal{N} = 2$ super-Teichmüller theory: prerequisites

$\mathcal{N} = 2$  super-Teichmüller space is related to  $OSP(2|2)$  supergroup of rank 2.

It is more useful to work with its  $3 \times 3$  incarnation, which is isomorphic to  $\Psi \ltimes SL(1|2)_0$ , where  $\Psi$  is a certain automorphism of the Lie algebra  $\mathfrak{sl}(1|2) \simeq \mathfrak{osp}(2|2)$ .

$SL(1|2)_0$  is a supergroup, consisting of supermatrices

$$g = \begin{pmatrix} a & b & \alpha \\ c & d & \beta \\ \gamma & \delta & f \end{pmatrix}$$

such that  $f > 0$  and their Berezinian = 1.

This group acts on the space  $\mathbb{C}^{1|2}$  as superconformal fractional-linear transformations.

As before,  $\mathcal{N} = 2$  super-Fuchsian groups are the ones whose projections

$$\pi_1 \rightarrow OSP(2|2) \rightarrow GL^+(2, \mathbb{R}) \rightarrow SL(2, \mathbb{R}) \rightarrow PSL(2, \mathbb{R})$$

are Fuchsian.

$\mathcal{N} = 2$  super-Teichmüller theory: prerequisites

$\mathcal{N} = 2$  super-Teichmüller space is related to  $OSP(2|2)$  supergroup of rank 2.

It is more useful to work with its  $3 \times 3$  incarnation, which is isomorphic to  $\Psi \ltimes SL(1|2)_0$ , where  $\Psi$  is a certain automorphism of the Lie algebra  $\mathfrak{sl}(1|2) \simeq \mathfrak{osp}(2|2)$ .

$SL(1|2)_0$  is a supergroup, consisting of supermatrices

$$g = \begin{pmatrix} a & b & \alpha \\ c & d & \beta \\ \gamma & \delta & f \end{pmatrix}$$

such that  $f > 0$  and their Berezinian = 1.

This group acts on the space  $\mathbb{C}^{1|2}$  as superconformal fractional-linear transformations.

As before,  $\mathcal{N} = 2$  super-Fuchsian groups are the ones whose projections

$$\pi_1 \rightarrow OSP(2|2) \rightarrow GL^+(2, \mathbb{R}) \rightarrow SL(2, \mathbb{R}) \rightarrow PSL(2, \mathbb{R})$$

are Fuchsian.

Outline

Introduction

Cast of characters

Coordinates on  
Super-Teichmüller  
space

$\mathcal{N} = 2$   
Super-Teichmüller  
theory

Further work

Open problems

$\mathcal{N} = 2$  super-Teichmüller theory: prerequisites

$\mathcal{N} = 2$  super-Teichmüller space is related to  $OSP(2|2)$  supergroup of rank 2.

It is more useful to work with its  $3 \times 3$  incarnation, which is isomorphic to  $\Psi \ltimes SL(1|2)_0$ , where  $\Psi$  is a certain automorphism of the Lie algebra  $\mathfrak{sl}(1|2) \simeq \mathfrak{osp}(2|2)$ .

$SL(1|2)_0$  is a supergroup, consisting of supermatrices

$$g = \begin{pmatrix} a & b & \alpha \\ c & d & \beta \\ \gamma & \delta & f \end{pmatrix}$$

such that  $f > 0$  and their Berezinian = 1.

This group acts on the space  $\mathbb{C}^{1|2}$  as superconformal fractional-linear transformations.

As before,  $\mathcal{N} = 2$  super-Fuchsian groups are the ones whose projections

$$\pi_1 \rightarrow OSP(2|2) \rightarrow GL^+(2, \mathbb{R}) \rightarrow SL(2, \mathbb{R}) \rightarrow PSL(2, \mathbb{R})$$

are Fuchsian.

Outline

Introduction

Cast of characters

Coordinates on  
Super-Teichmüller  
space

$\mathcal{N} = 2$   
Super-Teichmüller  
theory

Further work

Open problems

$\mathcal{N} = 2$  super-Teichmüller theory: prerequisites

$\mathcal{N} = 2$  super-Teichmüller space is related to  $OSP(2|2)$  supergroup of rank 2.

It is more useful to work with its  $3 \times 3$  incarnation, which is isomorphic to  $\Psi \ltimes SL(1|2)_0$ , where  $\Psi$  is a certain automorphism of the Lie algebra  $\mathfrak{sl}(1|2) \simeq \mathfrak{osp}(2|2)$ .

$SL(1|2)_0$  is a supergroup, consisting of supermatrices

$$g = \begin{pmatrix} a & b & \alpha \\ c & d & \beta \\ \gamma & \delta & f \end{pmatrix}$$

such that  $f > 0$  and their Berezinian = 1.

This group acts on the space  $\mathbb{C}^{1|2}$  as superconformal fractional-linear transformations.

As before,  $\mathcal{N} = 2$  super-Fuchsian groups are the ones whose projections

$$\pi_1 \rightarrow OSP(2|2) \rightarrow GL^+(2, \mathbb{R}) \rightarrow SL(2, \mathbb{R}) \rightarrow PSL(2, \mathbb{R})$$

are Fuchsian.

Outline

Introduction

Cast of characters

Coordinates on  
Super-Teichmüller  
space

$\mathcal{N} = 2$   
Super-Teichmüller  
theory

Further work

Open problems

Note, that the pure bosonic part of  $SL(1|2)_0$  is  $GL^+(2, \mathbb{R})$ .

Therefore, the construction of coordinates requires a new notion:  
 $\mathbb{R}_+$ -graph connection.

A  $G$ -graph connection on  $\tau$  is the assignment  $h_e \in G$  to each oriented edge  $e$  of  $\tau$  so that  $h_{\bar{e}} = h_e^{-1}$  if  $\bar{e}$  is the opposite orientation to  $e$ .

Two assignments  $\{h_e\}, \{h'_e\}$  are equivalent iff there are  $t_v \in G$  for each vertex  $v$  of  $\tau$  such that  $h'_e = t_v h_e t_w^{-1}$  for each oriented edge  $e \in \tau$  with initial point  $v$  and terminal point  $w$ .

The moduli space of flat  $G$ -connections on  $F$  is isomorphic to the space of equivalent  $G$ -graph connections on  $\tau$ .

By the way, spin structures can be identified with equivalence classes of  $\mathbb{Z}_2$ -graph connections.

[Outline](#)[Introduction](#)[Cast of characters](#)[Coordinates on  
Super-Teichmüller  
space](#) [\$\mathcal{N} = 2\$   
Super-Teichmüller  
theory](#)[Further work](#)[Open problems](#)



Note, that the pure bosonic part of  $SL(1|2)_0$  is  $GL^+(2, \mathbb{R})$ .

Therefore, the construction of coordinates requires a new notion:

$\mathbb{R}_+$ -graph connection.

A  $G$ -graph connection on  $\tau$  is the assignment  $h_e \in G$  to each oriented edge  $e$  of  $\tau$  so that  $h_{\bar{e}} = h_e^{-1}$  if  $\bar{e}$  is the opposite orientation to  $e$ .

Two assignments  $\{h_e\}, \{h'_e\}$  are equivalent iff there are  $t_v \in G$  for each vertex  $v$  of  $\tau$  such that  $h'_e = t_v h_e t_w^{-1}$  for each oriented edge  $e \in \tau$  with initial point  $v$  and terminal point  $w$ .

The moduli space of flat  $G$ -connections on  $F$  is isomorphic to the space of equivalent  $G$ -graph connections on  $\tau$ .

By the way, spin structures can be identified with equivalence classes of  $\mathbb{Z}_2$ -graph connections.

Note, that the pure bosonic part of  $SL(1|2)_0$  is  $GL^+(2, \mathbb{R})$ .

Therefore, the construction of coordinates requires a new notion:

$\mathbb{R}_+$ -graph connection.

A  $G$ -graph connection on  $\tau$  is the assignment  $h_e \in G$  to each oriented edge  $e$  of  $\tau$  so that  $h_{\bar{e}} = h_e^{-1}$  if  $\bar{e}$  is the opposite orientation to  $e$ .

Two assignments  $\{h_e\}, \{h'_e\}$  are equivalent iff there are  $t_v \in G$  for each vertex  $v$  of  $\tau$  such that  $h'_e = t_v h_e t_w^{-1}$  for each oriented edge  $e \in \tau$  with initial point  $v$  and terminal point  $w$ .

The moduli space of flat  $G$ -connections on  $F$  is isomorphic to the space of equivalent  $G$ -graph connections on  $\tau$ .

By the way, spin structures can be identified with equivalence classes of  $\mathbb{Z}_2$ -graph connections.

Note, that the pure bosonic part of  $SL(1|2)_0$  is  $GL^+(2, \mathbb{R})$ .

Therefore, the construction of coordinates requires a new notion:

$\mathbb{R}_+$ -graph connection.

A  $G$ -graph connection on  $\tau$  is the assignment  $h_e \in G$  to each oriented edge  $e$  of  $\tau$  so that  $h_{\bar{e}} = h_e^{-1}$  if  $\bar{e}$  is the opposite orientation to  $e$ .

Two assignments  $\{h_e\}, \{h'_e\}$  are equivalent iff there are  $t_v \in G$  for each vertex  $v$  of  $\tau$  such that  $h'_e = t_v h_e t_w^{-1}$  for each oriented edge  $e \in \tau$  with initial point  $v$  and terminal point  $w$ .



The moduli space of flat  $G$ -connections on  $F$  is isomorphic to the space of equivalent  $G$ -graph connections on  $\tau$ .

By the way, spin structures can be identified with equivalence classes of  $\mathbb{Z}_2$ -graph connections.

## Data on triangulation/fatgraphs:

- ▶ One positive parameter per edge of fatgraph/triangulation
- ▶ Two odd parameters per triangle
- ▶ Two spin structures: generated by reflection of signs and the permutation of odd parameters
- ▶  $\mathbb{R}_+$ -graph connection

# Generic Ptolemy transformations are:



and the transformation formulas are as follows:

$$ef = (ac + bd) \left( 1 + \frac{h_e^{-1}\sigma_1\theta_2}{2(\sqrt{\chi} + \sqrt{\chi^{-1}})} + \frac{h_e\sigma_2\theta_1}{2(\sqrt{\chi} + \sqrt{\chi^{-1}})} \right),$$

$$\mu_1 = \frac{h_e\theta_1 + \sqrt{\chi}\sigma_1}{\mathcal{D}}, \quad \mu_2 = \frac{h_e^{-1}\theta_2 + \sqrt{\chi}\sigma_2}{\mathcal{D}},$$

$$\nu_1 = \frac{\sigma_1 - \sqrt{\chi}h_e\theta_1}{\mathcal{D}}, \quad \nu_2 = \frac{\sigma_2 - \sqrt{\chi}h_e^{-1}\theta_2}{\mathcal{D}},$$

$$h'_a = \frac{h_a}{h_e c_\theta}, \quad h'_b = \frac{h_b c_\theta}{h_e}, \quad h'_c = h_c \frac{c_\theta}{c_\mu}, \quad h'_d = h_d \frac{c_\nu}{c_\theta}, \quad h_f = \frac{c_\sigma}{c_\theta^2},$$

where

$$\mathcal{D} := \sqrt{1 + \chi + \frac{\sqrt{\chi}}{2}(h_e^{-1}\sigma_1\theta_2 + h_e\sigma_2\theta_1)},$$

$$c_\theta := 1 + \frac{\theta_1\theta_2}{6}.$$

Outline

Introduction

Cast of characters

Coordinates on  
Super-Teichmüller  
space

$\mathcal{N} = 2$   
Super-Teichmüller  
theory

Further work

Open problems

There is a parallel construction, based on Jenkins-Strebel differentials.

How to glue a Riemann surface based on a fatgraph with the metric data?

Jenkins-Strebel differential and the underlying fatgraph →  
special covering of Riemann surfaces with double overlaps,  
corresponding to the edges.

M. Kontsevich'92; M. Mulase, M. Penkava'98

In a joint work with A. Schwarz, we

- ▶ Explicitly construct deformations for the class of  $(1|1)$ -supermanifolds "of middle degree" with punctures as Čech cocycles
- ▶ Get in contact with the analogue of Penner's convex hull construction
- ▶ Construct  $N=1$  SRS using the dualities of  $(1|1)$ -supermanifolds/ $N = 2$  SRS

Outline

Introduction

Cast of characters

Coordinates on  
Super-Teichmüller  
space

$\mathcal{N} = 2$   
Super-Teichmüller  
theory

Further work

Open problems

There is a parallel construction, based on Jenkins-Strebel differentials.

How to glue a Riemann surface based on a fatgraph with the metric data?

Jenkins-Strebel differential and the underlying fatgraph  $\rightarrow$   
special covering of Riemann surfaces with double overlaps,  
corresponding to the edges.

M. Kontsevich'92; M. Mulase, M. Penkava'98

In a joint work with A. Schwarz, we

- ▶ Explicitly construct deformations for the class of  $(1|1)$ -supermanifolds "of middle degree" with punctures as Čech cocycles
- ▶ Get in contact with the analogue of Penner's convex hull construction
- ▶ Construct  $N=1$  SRS using the dualities of  $(1|1)$ -supermanifolds/ $N = 2$  SRS

Outline

Introduction

Cast of characters

Coordinates on  
Super-Teichmüller  
space

$\mathcal{N} = 2$   
Super-Teichmüller  
theory

Further work

Open problems



There is a parallel construction, based on Jenkins-Strebel differentials.

How to glue a Riemann surface based on a fatgraph with the metric data?

Jenkins-Strebel differential and the underlying fatgraph  $\rightarrow$  special covering of Riemann surfaces with double overlaps, corresponding to the edges.

M. Kontsevich'92; M. Mulase, M. Penkava'98

In a joint work with A. Schwarz, we

- ▶ Explicitly construct deformations for the class of  $(1|1)$ -supermanifolds "of middle degree" with punctures as Čech cocycles
- ▶ Get in contact with the analogue of Penner's convex hull construction
- ▶ Construct  $N=1$  SRS using the dualities of  $(1|1)$ -supermanifolds/ $N = 2$  SRS

Outline

Introduction

Cast of characters

Coordinates on Super-Teichmüller space

$\mathcal{N} = 2$  Super-Teichmüller theory

Further work

Open problems

# McShane-type identities, path to volumes?

The simplest McShane identity (G. McShane'92):

$$\frac{1}{2} = \sum_{\gamma} \frac{1}{1 + e^{l_{\gamma}}}$$

on a cusped torus, where sum is over all simple geodesics  $\gamma$  and  $l_{\gamma}$  is the length.

M. Mirzakhani used such types of identities to deal with the volumes of the moduli spaces.

Y. Huang recently shown how to deal with McShane identities using Penner's lambda length coordinates.

Together with Y. Huang, R. Penner, we have shown that the following generalization of McShane identity holds:

$$\frac{1}{2} = \sum_{\gamma} \left( \frac{1}{1 + e^{l_{\gamma}}} + \frac{e^{\frac{3}{2}l_{\gamma}} T}{e^{2l_{\gamma}} - 1} \right)$$

where  $l_{\gamma}$  is the superanalogue of geodesic length and  $T$  is a product of  $\mu$ -coordinates.

# McShane-type identities, path to volumes?

The simplest McShane identity (G. McShane'92):

$$\frac{1}{2} = \sum_{\gamma} \frac{1}{1 + e^{l_{\gamma}}}$$

on a cusped torus, where sum is over all simple geodesics  $\gamma$  and  $l_{\gamma}$  is the length.

M. Mirzakhani used such types of identities to deal with the volumes of the moduli spaces.

Y. Huang recently shown how to deal with McShane identities using Penner's lambda length coordinates.

Together with Y. Huang, R. Penner, we have shown that the following generalization of McShane identity holds:

$$\frac{1}{2} = \sum_{\gamma} \left( \frac{1}{1 + e^{l_{\gamma}}} + \frac{e^{\frac{3}{2}l_{\gamma}} T}{e^{2l_{\gamma}} - 1} \right)$$

where  $l_{\gamma}$  is the superanalogue of geodesic length and  $T$  is a product of  $\mu$ -coordinates.

- 1) Cluster superalgebras
- 2) Weil-Petersson-form in  $\mathcal{N} = 2$  case
- 3) Quantization of super-Teichmüller spaces
- 4) Analogues of Weil-Petersson volumes
- 5) Relation to Strebel theory
- 6) Quasi-abelianization to  $GL(1|1)$ /spectral network approach in the style of Gaiotto-Moore-Neitzke

Outline

Introduction

Cast of characters

Coordinates on  
Super-Teichmüller  
space

$\mathcal{N} = 2$   
Super-Teichmüller  
theory

Further work

**Open problems**

Outline

Introduction

Cast of characters

Coordinates on  
Super-Teichmüller  
space

$\mathcal{N} = 2$   
Super-Teichmüller  
theory

Further work

**Open problems**

Thank you!