

# Liouville-type theorems and decay estimates for solutions to higher order elliptic equations

Guozhen Lu, Peiyong Wang and Jiuyi Zhu

ABSTRACT. Liouville-type theorems are powerful tools in partial differential equations. Boundedness assumption of solutions are often imposed in deriving such Liouville-type theorems. In this paper, we establish some Liouville-type theorems without the boundedness assumption of nonnegative solutions to certain classes of elliptic equations and systems. Using a rescaling technique and doubling lemma developed recently in [PQS], we improve several Liouville-type theorems in higher order elliptic equations, some semi-linear equations and elliptic systems. More specifically, we remove the boundedness assumption of the solutions which is required in the proofs of the corresponding Liouville-type theorems in the recent literature. Moreover, we also investigate the singularity and decay estimates of higher order elliptic equations.

## 1. Introduction

The aim of this paper is to establish some Liouville-type theorems without the boundedness assumption on nonnegative solutions to certain classes of elliptic equations. Liouville-type theorems are powerful tools to prove a priori bounds for nonnegative solutions in a bounded domain. Using the rescaling method (also called the "blow-up" method) in the elegant paper [GS2], an equation in a bounded domain will blow up to become another equation in the whole Euclidean space or a half space. With the aid of the corresponding Liouville-type theorem in the Euclidean space  $\mathbb{R}^N$  and half space  $\mathbb{R}_+^N$  and a contradiction argument, the a priori bounds could be deduced. Moreover, the existence of nonnegative solutions to elliptic equations is established by the topological degree method using a priori estimates (see e.g. [DLN]).

In [PQS], the authors develop a general method for the derivation of universal, pointwise a priori estimates of local solutions from the Liouville-type theorem. Their results show that the universal bounds theorems for local solutions and the Liouville-type theorem are essentially equivalent. By further exploring their rescaling method, we also obtain that the boundedness assumption of solutions in proving the Liouville-type theorem is not essential in many cases.

---

1991 *Mathematics Subject Classification.* 35B53, 35J40, 35J47, 35B45,

*Key words and phrases.* Higher order elliptic equations, Polyharmonic operators on half spaces, Dirichlet problem, Navier boundary condition, Liouville-type theorem, Decay estimates for solutions, Doubling property, Without boundedness assumptions.

Research is partly supported by a US NSF grant #DMS0901761.

Corresponding Author: Jiuyi Zhu at jiuyi.zhu@wayne.edu.

We denote the Sobolev critical exponent by

$$p_S := \begin{cases} \frac{N+2m}{N-2m} & \text{if } N > 2m, \\ \infty & \text{if } N \leq 2m, \end{cases}$$

where  $m \in \mathbb{N}$  and  $N$  is the dimension of Euclidean space  $\mathbb{R}^N$  or half space  $\mathbb{R}_+^N$ . The Liouville-type theorem for the subcritical higher order elliptic equations in Euclidean space was established in [WX] as follows,

**Theorem A:** Let  $1 < q < p_S$ . If  $u$  is a classical nonnegative solution of

$$(1.1) \quad (-\Delta)^m u = u^q \quad \text{in } \mathbb{R}^N,$$

then  $u \equiv 0$ .

For the higher order elliptic equation in half space, generally speaking, two types of boundary conditions, i.e. the Dirichlet boundary problem and the Navier boundary problem, are considered. The Liouville theorem for the Dirichlet boundary type of higher order equation in half space has been obtained in [RW].

**Theorem B:** Let  $1 < q < p_S$ . If  $u$  is a classical solution of

$$(1.2) \quad \begin{cases} (-\Delta)^m u = u^q & \text{in } \mathbb{R}_+^N, \\ u \geq 0 & \text{in } \mathbb{R}_+^N, \\ u = \frac{\partial u}{\partial x_N} = \dots = \frac{\partial^{m-1} u}{\partial x_N^{m-1}} = 0 & \text{on } \partial \mathbb{R}_+^N, \end{cases}$$

then  $u \equiv 0$ .

The Liouville-type theorem for the Navier boundary problem of higher order elliptic equation in half space has been studied in [BM], [GL] and [S]. The authors consider higher order elliptic equation

$$(1.3) \quad \begin{cases} (-\Delta)^m u = u^p & \text{in } \mathbb{R}_+^N, \\ u \geq 0 & \text{in } \mathbb{R}_+^N, \\ u = \Delta u = \dots = \Delta^{m-1} u = 0 & \text{on } \partial \mathbb{R}_+^N. \end{cases}$$

Under the following boundedness assumption on the solution:

$$(1.4) \quad \sup u < \infty, \quad \sup |\Delta^i u| < \infty, \quad i = 1, 2, \dots, m-1.$$

They established that the solutions in (1.3) are trivial if  $2m+1 < N$  and  $1 < p < \frac{N-1+2m}{N-1-2m}$  or if  $2m+1 \geq N$  and  $1 < p < \infty$ . The proof of their results is based on an idea of [D]. The idea is the following: if there exists a solution of (1.3) and one is able to show that any solution is increasing in the  $x_N$  direction, then passing the limit as  $x_N \rightarrow \infty$ , one could get a solution of the same equation in  $\mathbb{R}^{N-1}$ , which in turn allows the use of the Liouville-type theorem in the whole space. (Note that the critical exponent in (1.3) is  $\frac{N-1+2m}{N-1-2m}$ ). We would like to mention that the Liouville-type theorem for (1.3) in [D] is for the case of  $m = 1$  with the assumption that the solution is bounded. Later on this result was improved to hold for unbounded solution in [BCN], among many other results.

We also mention that Liouville-type theorems for differential inequalities and systems involving polyharmonic operators and related to the Hardy-Littlewood-Sobolev inequalities have also been extensively studied. We refer the reader to the paper by Caristi, D'Ambrosio and Mitidieri [CDM] and references therein (see also [M1]).

A natural question is then whether the boundedness assumption for the Liouville-type theorem is necessary for higher order elliptic equation or not. In particular, we are

interested in the Liouville-type theorems for solutions to polyharmonic equations. We show that it is indeed unnecessary for such equations and establish that

**THEOREM 1.** *If  $u$  is a classical solution of (1.3) with  $2m + 1 \leq N$  and  $1 < p < \frac{N+2m}{N-2m}$  or with  $2m + 1 > N$  and  $1 < p < \infty$ , then  $u \equiv 0$ .*

In addition, we consider the a priori estimates of possible singularities for local solutions of higher order elliptic equation

$$(1.5) \quad (-\Delta)^m u = f(u)$$

when  $f$  satisfies certain conditions. More precisely, we will obtain

**THEOREM 2.** *Let  $1 < p < p_S$  and  $\Omega \neq \mathbb{R}^N$  be a domain in  $\mathbb{R}^N$ . Assume that the function  $f : [0, \infty) \rightarrow \mathbb{R}$  is continuous,*

$$(1.6) \quad \lim_{u \rightarrow \infty} u^{-p} f(u) = \rho \in (0, \infty).$$

*Then there exists  $C(N, f, m) > 0$  independent of  $\Omega$  and  $u$  such that for any positive solution  $u$  of (1.5) in  $\Omega$ , there holds*

$$(1.7) \quad \sum_{|\nu|=0}^{2m-1} |D^\nu u(x)| \leq C(N, f, m)(1 + \text{dist}^{-\frac{2m}{p-1}}(x, \partial\Omega)), \quad x \in \Omega,$$

where  $\nu = \{\nu_1, \dots, \nu_N\}$ ,  $D^\nu u = D_{x_1}^{\nu_1} u \cdots D_{x_N}^{\nu_N} u$  and  $\text{dist}(x, \partial\Omega)$  is the distance of  $x$  from  $\partial\Omega$ .

*In particular, if  $\Omega = B_R \setminus \{0\}$  for some  $R > 0$ , then*

$$\sum_{|\nu|=0}^{2m-1} |D^\nu u(x)| \leq C(N, f, m)(1 + |x|^{-\frac{2m}{p-1}}), \quad 0 < |x| \leq \frac{R}{2},$$

where  $B_R$  is a ball centered at 0 with radius  $R$ .

In the special case of  $f(u) = u^p$ . More precise results can be given in bounded or exterior domains.

**THEOREM 3.** *Let  $1 < p < p_S$  and  $\Omega \neq \mathbb{R}^N$  be a domain in  $\mathbb{R}^N$ . Then there exists  $C(N, p, m) > 0$  independent of  $\Omega$  and  $u$  such that any nonnegative solution  $u$  in (1.5) satisfies*

$$(1.8) \quad \sum_{|\nu|=0}^{2m-1} |D^\nu u(x)| \leq C(N, p, m) \text{dist}^{-\frac{2m}{p-1}}(x, \partial\Omega), \quad x \in \Omega.$$

*In particular if  $\Omega$  is an exterior domain, i.e. the set  $\{x \in \mathbb{R}^N; |x| > R\} \subset \Omega$  for some  $R > 0$ , then*

$$\sum_{|\nu|=0}^{2m-1} |D^\nu u(x)| \leq C(N, p, m) |x|^{-\frac{2m}{p-1}}, \quad |x| \geq 2R.$$

The Liouville-type theorems for semilinear elliptic equations in Euclidean space  $\mathbb{R}^N$  and half space  $\mathbb{R}_+^N$  are well-known in literature. Among many results in [GS1], the Liouville-type theorem for the subcritical elliptic equation is shown. Namely, there is no nontrivial  $C^2$  solution of the problem

$$(1.9) \quad \begin{cases} -\Delta u = u^p & \text{in } \mathbb{R}^N, \\ u \geq 0 & \text{in } \mathbb{R}^N, \end{cases}$$

if  $N \geq 3$  and  $1 < p < \frac{N+2}{N-2}$ . See also [CL] for a simpler proof with the Kelvin transform and moving plane method of Gidas-Ni-Nirenberg [GNN]. In [DG], the authors consider the following mixed (Dirichlet-Neumann) boundary conditions in a half space, i.e. the problem

$$(1.10) \quad \begin{cases} -\Delta u = u^p & \text{in } \mathbb{R}_+^N, \\ u \geq 0 & \text{in } \mathbb{R}_+^N, \\ u = 0 & \text{on } \Gamma_0 := \{x \in \partial\mathbb{R}_+^N | x_N = 0, x_1 > 0\}, \\ \frac{\partial u}{\partial x_N} = 0 & \text{on } \Gamma_1 := \{x \in \partial\mathbb{R}_+^N | x_N = 0, x_1 < 0\}, \end{cases}$$

where  $N \geq 3$  and  $1 < p < \frac{N+2}{N-2}$ . Using the idea of the Kelvin transform combined with the moving plane method, they prove the nonexistence of nontrivial solutions in (1.10) under the assumption that the solution is bounded. In next theorem, we are able to remove this boundedness assumption of the solution. In fact, our theorem is stated as:

**THEOREM 4.** *If  $u \in C^2(\mathbb{R}_+^N) \cap C(\overline{\mathbb{R}_+^N})$  is a solution of (1.10) with  $N \geq 3$  and  $1 < p < \frac{N+2}{N-2}$ , then  $u \equiv 0$ .*

For the semilinear elliptic system, an important model is the Lane-Emden system:

$$(1.11) \quad \begin{cases} -\Delta u_1 = u_2^p & \text{in } \mathbb{R}^N, \\ -\Delta u_2 = u_1^q & \text{in } \mathbb{R}^N. \end{cases}$$

It is conjectured that if

$$\frac{1}{p+1} + \frac{1}{q+1} > \frac{n-2}{n},$$

then there are no nontrivial classical solutions of (1.11) in  $\mathbb{R}^N$ . The conjecture has been proved to be true for radial solutions in all dimensions in [M]. The cases of  $N = 3, 4$  for the conjecture in general have been also solved recently in [PQS] and [So] respectively. The interested reader can refer to the above papers and references therein for detailed descriptions (see also the works [BMa], [SZ], etc.). In [DS], the authors study the elliptic system

$$(1.12) \quad \begin{cases} \Delta u_1 + u_1^{\alpha_1} + u_2^{\frac{\alpha_2-1}{\alpha_1-1}} = 0 & \text{in } \mathbb{R}^N, \\ \Delta u_2 + u_2^{\alpha_2} + u_1^{\frac{\alpha_1-1}{\alpha_2-1}} = 0 & \text{in } \mathbb{R}^N, \end{cases}$$

and prove that (1.12) does not have bounded positive classical solutions in  $\mathbb{R}^N$  if

$$(1.13) \quad 1 < \alpha_1, \alpha_2 < \frac{N+2}{N-2}.$$

We also show that the boundedness assumption of the solution for the elliptic system (1.12) is not necessary. Indeed, we can establish that

**THEOREM 5.** *The classical positive solutions  $u_1$  and  $u_2$  in (1.12) are trivial under the assumption of (1.13).*

We end our introduction by mentioning that nonexistence results of nonnegative solutions have been recently established by the first and third authors [LZ] for certain classes of integral equations which are closely related to polyharmonic equations in half spaces by a completely different method from that used in the current paper, namely, the method of moving planes in integral form developed in [CLO].

The outline of the paper is as follows. Section 2 is devoted to obtaining the Liouville-type theorem, then singularity and decay estimates for higher order elliptic equations. In Section 3, the Liouville-type theorem with mixed boundary condition for semilinear elliptic equation is shown. The Liouville-type theorem for the elliptic systems is considered in Section 4. Throughout the paper,  $C$  and  $c$  denote generic positive constants, which are independent of  $u$  and may vary from line to line.

## 2. Higher order elliptic equations

We state the following technical lemma used frequently in this paper.

**LEMMA 1.** *(Doubling lemma) Let  $(X, d)$  be a complete metric space and  $\emptyset \neq D \subset \Sigma \subset X$ , with  $\Sigma$  closed. Define  $M : D \rightarrow (0, \infty)$  to be bounded on compact subsets of  $D$  and fix a positive number  $k$ . If  $y \in D$  is such that*

$$M(y) \text{dist}(y, \Gamma) > 2k,$$

where  $\Gamma = \Sigma \setminus D$ , then there exists  $x \in D$  such that

$$M(x) \text{dist}(x, \Gamma) > 2k, \quad M(x) \geq M(y)$$

and

$$M(z) \leq 2M(x), \quad \forall z \in D \cap \bar{B}(x, kM^{-1}(x)).$$

**REMARK 1.** *If  $\Gamma = \emptyset$ , then  $\text{dist}(x, \Gamma) := \infty$ .*

The proof of the above lemma is given in [PQS]. Based on the doubling property, we can start the rescaling process to prove local estimates of solutions of superlinear problems. The idea is by a contradiction argument. One can argue that the local estimates of solutions are violated. By an appropriate rescaling, the blow up sequence of solutions converges to a bounded solution of a limiting equation in  $\mathbb{R}^N$ . Then from the non-existence of solutions in the limiting equation, we could infer that the local estimates exist. In this spirit, we conclude the proof of Theorem 1. We first state the lemma about the local a priori estimates for higher order elliptic equation (see [ADN] or [RW]). Let  $u$  satisfy the following equation

$$(2.1) \quad (-\Delta)^m u = g(x) \quad \text{in } \Omega.$$

Then we have

**LEMMA 2.** *Let  $\Omega$  be a ball  $\{x \in \mathbb{R}^N : |x| < R\}$ . Suppose  $u \in W^{2m,q}$  satisfies (2.1). Then there exists a constant  $C > 0$  depending only on  $\Omega, N, m$  and  $R$  such that for any  $\sigma \in (0, 1)$*

$$(2.2) \quad \|u\|_{W^{2m,q}(\Omega \cap B_{\sigma R})} \leq \frac{C}{(1-\sigma)^{2m}} (\|g\|_{L^q(\Omega)} + \|u\|_{L^q(\Omega)}).$$

The following lemma is the Liouville-type theorem for higher order elliptic equations with boundedness assumption in [S].

LEMMA 3. *The higher order elliptic equation (1.3) does not have positive solutions under assumption of (1.4) provided  $p < \frac{N-1+2m}{N-1-2m}$  and  $2m+1 < N$  (or  $N \geq 2m+1$  and  $p < \infty$ ).*

PROOF OF THEOREM 1: Suppose that a solution  $u$  to the equation (1.3) is unbounded in the sense that (1.4) is violated. Namely, the following boundedness is not true

$$\sup u < \infty, \quad \sup |\Delta^i u| < \infty, \quad i = 1, 2, \dots, m-1.$$

Then, there exists a sequence of  $(y_k) \in \mathbb{R}^N$  such that

$$\sum_{i=0}^{m-1} |\Delta^i u(y_k)| \rightarrow \infty$$

as  $k \rightarrow \infty$ . Set

$$M(y) := \left( \sum_{i=0}^{m-1} |\Delta^i u(y)| \right)^{\frac{p-1}{2m}} : \mathbb{R}_+^N \rightarrow \mathbb{R}.$$

Then  $M(y_k) \rightarrow \infty$  as  $k \rightarrow \infty$ . By taking  $D = \Sigma = X = \overline{\mathbb{R}_+^N}$  in the doubling lemma and Remark 1 ( See also, e.g. [RW]), there exists another sequence of  $(x_k)$  such that

$$M(x_k) \geq M(y_k)$$

and

$$M(z) \leq 2M(x_k), \quad \forall z \in B_{k/M(x_k)}(x_k) \cap \overline{\mathbb{R}_+^N}.$$

Define

$$d_k := x_{k,N} M(x_k)$$

and

$$H_k := \{\xi \in \mathbb{R}^N | \xi_N > -d_k\}.$$

We introduce a new function

$$v_k(\xi) := \frac{u(x_k + \frac{\xi}{M(x_k)})}{M^{\frac{2m}{p-1}}(x_k)}.$$

Then,  $v_k(\xi)$  is the nonnegative solution of

$$(2.3) \quad \begin{cases} (-\Delta)^m v_k = v_k^p & \text{in } H_k, \\ v_k = \Delta v_k = \dots = \Delta^{m-1} v_k = 0 & \text{on } \partial H_k \end{cases}$$

satisfying

$$(2.4) \quad \sum_{i=0}^{m-1} |\Delta^i v_k(0)|^{\frac{p-1}{2m+2(p-1)i}} = 1$$

and

$$(2.5) \quad \sum_{i=0}^{m-1} |\Delta^i v_k(\xi)|^{\frac{p-1}{2m+2(p-1)i}} \leq 2, \quad \forall \xi \in H_k \cap B_k(0).$$

Two cases may occur as  $k \rightarrow \infty$ , either case (1)

$$x_{k,N} M(x_k) \rightarrow \infty$$

for a subsequence still denoted as before, or case (2)

$$x_{k,N}M(x_k) \rightarrow d$$

for a subsequence still denoted as before, here  $d \geq 0$ . If case (1) occurs, i.e.  $H_k \cap B_k(0) \rightarrow \mathbb{R}^N$  as  $k \rightarrow \infty$ , then for any smooth compact  $D$  in  $\mathbb{R}^N$ , there exists  $k_0$  large enough such that  $D \subset (H_k \cap B_k(0))$  as  $k \geq k_0$ . By the classical  $W^{2m,q}$  estimates for higher order elliptic equation in Lemma 2 and (2.5), we have  $\|v_k\|_{W^{2m,q}(\bar{D})} \leq C$  for any  $1 < q < \infty$ . Therefore, we can extract a convergent subsequence  $v_k \rightarrow v$  in  $D$ , where  $v \in C^{2m-1,\tau}$  for some  $\tau > 0$ . Furthermore, using a diagonal line argument,  $v_k \rightarrow v$  in  $C_{loc}^{2m-1,\tau}(\mathbb{R}^N)$  and  $v$  solves

$$(-\Delta)^m v = v^p \quad \text{in } \mathbb{R}^N.$$

From Theorem A,  $u$  is trivial. However, (2.4) implies that

$$\sum_{i=0}^{m-1} |\Delta^i v(0)|^{\frac{p-1}{2m+2(p-1)i}} = 1,$$

which indicates that  $u$  is nontrivial. Obviously a contradiction is arrived.

If case (2) occurs, we make a further translation. Set

$$\tilde{v}_k(\xi) := v_k(\xi - d_k e_N) \quad \text{for } \xi \in \overline{\mathbb{R}_+^N}.$$

Then  $\tilde{v}_k$  satisfies

$$(2.6) \quad \begin{cases} (-\Delta)^m \tilde{v}_k = \tilde{v}_k^p & \text{in } \mathbb{R}_+^N, \\ \tilde{v}_k = \Delta \tilde{v}_k = \dots = \Delta^{m-1} \tilde{v}_k = 0 & \text{on } \partial \mathbb{R}_+^N. \end{cases}$$

While

$$(2.7) \quad \sum_{i=0}^{m-1} |\Delta^i \tilde{v}_k(d_k e_N)|^{\frac{p-1}{2m+2(p-1)i}} = 1$$

and

$$(2.8) \quad \sum_{i=0}^{m-1} |\Delta^i \tilde{v}_k(\xi)|^{\frac{p-1}{2m+2(p-1)i}} \leq 2, \quad \forall \xi \in \mathbb{R}_+^N \cap B_k(d_k e_N).$$

Observe that (2.6) could also be reduced to a system of elliptic equations. Let

$$(\tilde{v}_k, \dots, (-\Delta)^{m-1} \tilde{v}_k) := (w_k^{(0)}, \dots, w_k^{(m-1)}).$$

Then,  $(w_k^{(0)}, \dots, w_k^{(m-1)})$  solves

$$(2.9) \quad \begin{cases} -\Delta w_k^{(m-1)} = (w_k^{(0)})^p & \text{in } \mathbb{R}_+^N, \\ -\Delta w_k^{(i)} = w_k^{(i+1)} & \text{in } \mathbb{R}_+^N, \quad i = 0, \dots, m-2, \\ w_k^{(0)} = \dots = w_k^{(m-1)} = 0 & \text{on } \partial \mathbb{R}_+^N. \end{cases}$$

Furthermore,

$$\sum_{i=0}^{m-1} |w_k^{(i)}(d_k e_N)|^{\frac{p-1}{2m+2(p-1)i}} = 1$$

and

$$\sum_{i=0}^{m-1} |w_k^{(i)}(\xi)|^{\frac{p-1}{2m+2(p-1)i}} \leq 2, \quad \forall \xi \in \mathbb{R}_+^N \cap B_k(d_k e_N).$$

For any smooth compact  $D$  in  $\overline{\mathbb{R}_+^N}$ , there exists  $k_0$  large enough such that  $D \subset (\overline{\mathbb{R}_+^N} \cap B_k(d_k e_N))$  for any  $k > k_0$ . By the classical elliptic estimates,

$$\sum_{i=0}^{m-1} \|w_k^{(i)}(\xi)\|_{C^{2,\tau}(D)} \leq C$$

for some  $\tau > 0$ . Thanks to the Arzelá-Ascoli Theorem, there exists a function  $(v^{(0)}, \dots, v^{(m-1)})$  such that  $(w_k^{(0)}, \dots, w_k^{(m-1)})$  converges to  $(v^{(0)}, \dots, v^{(m-1)})$  in  $C^2(\bar{D})$ . Through a diagonal line argument,  $(w_k^{(0)}, \dots, w_k^{(m-1)})$  converges in  $C_{loc}^2(\overline{\mathbb{R}_+^N})$  to  $(v^{(0)}, \dots, v^{(m-1)})$ , which solves

$$(2.10) \quad \begin{cases} -\Delta v^{(m-1)} = (v^{(0)})^p & \text{in } \mathbb{R}_+^N, \\ -\Delta v^{(i)} = v^{(i+1)} & \text{in } \mathbb{R}_+^N, \quad i = 0, \dots, m-2, \\ v^{(0)} = \dots = v^{(m-1)} = 0 & \text{on } \partial\mathbb{R}_+^N, \end{cases}$$

and satisfies

$$(2.11) \quad \sum_{i=0}^{m-1} |v^{(i)}(de_N)|^{\frac{p-1}{2m+2(p-1)i}} = 1$$

and

$$(2.12) \quad \sum_{i=0}^{m-1} |v^{(i)}(\xi)|^{\frac{p-1}{2m+2(p-1)i}} \leq 2, \quad \forall \xi \in \mathbb{R}_+^N.$$

From (2.10),  $v^{(0)}$  actually solves

$$(2.13) \quad \begin{cases} (-\Delta)^m v^{(0)} = (v^{(0)})^p & \text{in } \mathbb{R}_+^N, \\ v^{(0)} = \Delta v^{(0)} = \dots = \Delta^{m-1} v^{(0)} = 0 & \text{on } \partial\mathbb{R}_+^N. \end{cases}$$

Hence  $v^{(0)} \equiv 0$  by lemma 3. Note that  $\frac{N-1+2m}{N-1-2m} > \frac{N+2m}{N-2m}$ . Then  $v^{(1)} \equiv \dots \equiv v^{(m-1)} \equiv 0$ , which contradicts (2.11). Therefore, the proof is completed.  $\square$

Next, we consider the singularity and decay estimates of the higher order elliptic equation. The main idea is the same as the case of the semilinear elliptic equation in [PQS].

**PROOF OF THEOREM 2:** Suppose (1.7) is not true. Then there exist sequences of  $\Omega_k, u_k, y_k \in \Omega_k$  such that  $u_k$  solves (1.5) on  $\Omega_k$  and the function

$$M_k := \sum_{|\nu|=0}^{2m-1} |D^\nu u_k|^{\frac{p-1}{2m+(p-1)|\nu|}}$$

satisfies

$$M_k(y_k) \geq 2k(1 + \text{dist}^{-1}(y_k, \partial\Omega)) \geq 2k \text{dist}^{-1}(y_k, \partial\Omega).$$

Adapting from the doubling lemma, there exists  $x_k \in \Omega_k$  such that

$$M_k(x_k) \geq M_k(y_k),$$

$$M_k(x_k) > 2k \text{dist}^{-1}(x_k, \partial\Omega_k)$$

and

$$M_k(z) \leq 2M_k(x_k), \quad \text{if } |z - x_k| \leq kM_k^{-1}(x_k).$$



We introduce a new function

$$v_k(\xi) := \frac{u_k(x_k + \frac{\xi}{M_k(x_k)})}{M_k^{\frac{2m}{p-1}}(x_k)}, \quad \forall |\xi| \leq k.$$

Then, the function  $v_k$  solves

$$(2.14) \quad (-\Delta)^m v_k(\xi) = f_k(v_k(\xi)) := M_k^{\frac{-2mp}{p-1}}(x_k) f(M_k^{\frac{2m}{p-1}}(x_k) v_k).$$

Also,

$$(2.15) \quad \sum_{|\nu|=0}^{2m-1} |D^\nu v_k|^{\frac{p-1}{2m+(p-1)|\nu|}}(0) = 1$$

and

$$(2.16) \quad \sum_{|\nu|=0}^{2m-1} |D^\nu v_k|^{\frac{p-1}{2m+(p-1)|\nu|}}(\xi) \leq 2, \quad \forall |\xi| \leq k.$$

By (1.6) and the continuity of  $f$ , we have

$$-c \leq f(s) \leq C(s^p + 1), \quad \text{for } s \geq 0.$$

Furthermore, it follows that

$$-c M_k^{\frac{-2mp}{p-1}}(x_k) \leq f_k(v_k(\xi)) \leq C, \quad \forall |\xi| \leq k.$$

For any smooth compact  $D$  in  $\mathbb{R}^N$ , there exists  $k_0$  large enough such that  $D \subset B_k(0)$  if  $k \geq k_0$ . Using the classical estimate of the higher order elliptic equation in (2.2) and (2.16),  $v_k(\xi) \in W_{loc}^{2m,q}(\mathbb{R}^N)$  for any  $1 < q < \infty$  and

$$\|v_k\|_{W^{2m,q}(\bar{D})} \leq C.$$

By the Sobolev imbedding and a diagonal line argument,  $v_k(\xi)$  converges in  $C_{loc}^{2m-1,\tau}(\mathbb{R}^N)$  to a function  $v \in C_{loc}^{2m-1,\tau}(\mathbb{R}^N)$  for some  $\tau > 0$ . Since  $v_k(\xi)$  is positive for  $\xi \in \mathbb{R}^N$ , then

$$f_k(v_k) \rightarrow \rho v^p \quad \text{as } k \rightarrow \infty.$$

Consequently,  $v$  is a nonnegative solution of

$$(-\Delta)^m v(\xi) = \rho v^p(\xi), \quad \xi \in \mathbb{R}^N.$$

By an appropriate rescaling,  $v$  is trivial from Theorem A, which contradicts (2.15). Hence, the conclusion holds.  $\square$

Next, we turn to the special case of  $f(u) = u^p$ .

**PROOF OF THEOREM 3:** Suppose (1.8) fails, Then, there exist sequences of  $\Omega_k$ ,  $u_k$ ,  $y_k \in \Omega_k$  such that  $u_k$  solves (1.5) in  $\Omega_k$  and the function

$$M_k := \sum_{|\nu|=0}^{2m-1} |D^\nu u_k|^{\frac{p-1}{2m+(p-1)|\nu|}}$$

satisfies

$$M_k(y_k) > 2k \text{dist}^{-1}(y_k, \partial\Omega_k).$$

By the doubling lemma, it follows that there exists  $x_k \in \Omega_k$  such that

$$\begin{aligned} M_k(x_k) &\geq M_k(y_k), \\ M_k(x_k) &> 2k \operatorname{dist}^{-1}(x_k, \partial\Omega_k) \end{aligned}$$

and

$$M_k(z) \leq 2M_k(x_k), \quad \text{if } |z - x_k| \leq kM_k^{-1}(x_k).$$

As before, we introduce a new function

$$v_k(\xi) := \frac{u_k(x_k + \frac{\xi}{M_k(x_k)})}{M_k^{\frac{2m}{p-1}}(x_k)}, \quad \forall |\xi| \leq k.$$

Then, the function  $v_k$  satisfies

$$(2.17) \quad (-\Delta)^m v_k(\xi) = v_k^p(\xi), \quad \forall |\xi| \leq k$$

with

$$(2.18) \quad \sum_{|\nu|=0}^{2m-1} |D^\nu v_k|^{\frac{p-1}{2m+(p-1)|\nu|}}(0) = 1$$

and

$$\sum_{|\nu|=0}^{2m-1} |D^\nu v_k|^{\frac{p-1}{2m+(p-1)|\nu|}}(\xi) \leq 2, \quad \forall |\xi| \leq k.$$

For any smooth compact  $D$  in  $\mathbb{R}^N$ , there exists  $k_0$  large enough such that  $D \subset B_k(0)$  if  $k \geq k_0$ . By the classical  $W^{2m,q}$  estimates for higher order elliptic equation in Lemma 2, we have  $\|v_k\|_{W^{2m,q}(\bar{D})} \leq C$  for any  $1 < q < \infty$ . Furthermore, using the Sobolev imbedding Theorem and a diagonal line argument,  $v_k \rightarrow v$  in  $C_{loc}^{2m-1,\tau}(\mathbb{R}^N)$  and  $v$  solves

$$(-\Delta)^m v = v^p.$$

With the aid of Theorem A,  $u$  is trivial. Nevertheless, from (2.18), it is impossible to be trivial. Hence, a contradiction is arrived. Then the theorem is verified.  $\square$

### 3. Elliptic equation with mixed boundary condition

In this section, we consider the Liouville-type theorem with mixed boundary condition.

LEMMA 4. *Let  $u \in W_{loc}^{1,2}(\mathbb{R}_+^N) \cap C^0(\overline{\mathbb{R}_+^N})$  be a weak solution of (1.10). Then the only bounded solution is  $u \equiv 0$ .*

The lemma above is the Liouville-type theorem with boundedness assumptions of solutions in [DG].

PROOF OF THEOREM 4: Suppose that there exists a unbounded solution  $u$ . Then, there exists a sequence  $(y_k) \in \mathbb{R}_+^N$  such that  $u(y_k) \rightarrow \infty$  as  $k \rightarrow \infty$ . Let  $M := u^{\frac{p-1}{2}} : \mathbb{R}_+^N \rightarrow \mathbb{R}$ , then  $M(y_k) \rightarrow \infty$  as  $k \rightarrow \infty$  since  $p > 1$ . Following from the doubling lemma by taking  $D = \Sigma = X = \overline{\mathbb{R}_+^N}$  and Remark 1, there exists another sequence of  $(x_k)$  such that

$$M(x_k) \geq M(y_k)$$

and

$$M(z) \leq 2M(x_k), \quad \forall z \in B_{k/M(x_k)}(x_k) \cap \overline{\mathbb{R}_+^N}.$$

Define

$$d_k := x_{k,N}M(x_k)$$

and

$$H_k := \{\xi \in \mathbb{R}^N \mid \xi_N > -d_k\}.$$

As in Theorem 1, we introduce a new function

$$v_k(\xi) = \frac{u(x_k + \frac{\xi}{M(x_k)})}{M^{\frac{2}{p-1}}(x_k)}.$$

Then,  $v_k(\xi)$  is the nonnegative solution of

$$(3.1) \quad \begin{cases} -\Delta v_k = v_k^p & \text{in } H_k, \\ v_k \geq 0 & \text{in } H_k, \\ v_k = 0 & \text{on } \Gamma_0^k := \{\xi \in H_k \mid \xi_N = -d_k, \xi_1 > 0\}, \\ \frac{\partial v_k}{\partial \xi_N} = 0 & \text{on } \Gamma_1^k := \{\xi \in H_k \mid \xi_N = -d_k, \xi_1 < 0\} \end{cases}$$

satisfying

$$(3.2) \quad v_k(0)^{\frac{p-1}{2}} = 1$$

and

$$(3.3) \quad v_k^{\frac{p-1}{2}}(\xi) \leq 2, \quad \forall \xi \in H_k \cap B_k(0).$$

As before, two cases may occur as  $k \rightarrow \infty$ , either case (1)

$$x_{k,N}M(x_k) \rightarrow \infty$$

for a subsequence still denoted as before, or case (2)

$$x_{k,N}M(x_k) \rightarrow d$$

for a subsequence still denoted as before, here  $d \geq 0$ . If case (1) occurs, i.e.  $H_k \cap B_k(0) \rightarrow \mathbb{R}^N$  as  $k \rightarrow \infty$ , then for any smooth compact  $D$  in  $\mathbb{R}^N$ , there exists  $k_0$  large enough such that  $D \subset (H_k \cap B_k(0))$  as  $k \geq k_0$ . By the classical elliptic equation estimates and (3.3), we have  $\|v_k\|_{W^{2,q}(\bar{D})} \leq C$  for any  $1 < q < \infty$ . Therefore, from the Sobolev embedding theorem, we can extract a convergent subsequence  $v_k$  that converges to  $v$  in  $D$ , where  $v \in C^{1,\tau}(\bar{D})$ . Furthermore, using a diagonalization argument,  $v_k \rightarrow v$  in  $C_{loc}^{1,\tau}(\mathbb{R}^N)$  and  $v$  solves

$$-\Delta v = v^p \quad \text{in } \mathbb{R}^N.$$

By the classical Liouville theorem of semilinear elliptic equation in Euclidean space,  $v$  is trivial, which contradicts (3.2).

If case (2) occurs, we make a further translation. Define

$$\tilde{v}_k(\xi) := v_k(\xi - d_k e_N) \quad \text{for } \xi \in \overline{\mathbb{R}_+^N}.$$

Then  $\tilde{v}_k$  satisfies

$$(3.4) \quad \begin{cases} -\Delta \tilde{v}_k = \tilde{v}_k^p & \text{in } \mathbb{R}_+^N, \\ \tilde{v}_k \geq 0 & \text{in } \mathbb{R}_+^N, \\ \tilde{v}_k = 0 & \text{on } \tilde{\Gamma}_0^k := \{\xi \in \mathbb{R}_+^N \mid \xi_N = 0, \xi_1 > 0\}, \\ \frac{\partial \tilde{v}_k}{\partial \xi_N} = 0 & \text{on } \tilde{\Gamma}_1^k := \{\xi \in \mathbb{R}_+^N \mid \xi_N = 0, \xi_1 < 0\}. \end{cases}$$

While,

$$(3.5) \quad \tilde{v}_k^{\frac{p-1}{2}}(d_k e_N) = 1$$

and

$$(3.6) \quad \tilde{v}_k^{\frac{p-1}{2}}(\xi) \leq 2, \quad \forall \xi \in \mathbb{R}_+^N \cap B_k(d_k e_N).$$

For any smooth compact neighborhood  $D$  of the origin in  $\overline{\mathbb{R}_+^N}$ , we have  $\{\tilde{v}_k\}$  is uniformly bounded in  $C^{2,\tau}(K) \cap C^\tau(\bar{D})$  for some  $0 < \tau < \frac{1}{2}$ , where  $K$  is a compact set of  $\overline{\mathbb{R}_+^N}$  with  $\text{dist}(K, \tilde{\Gamma}) > 0$  (The a priori estimates could be done in section 6 in [CP]) and where  $\tilde{\Gamma} := \{\xi \in \mathbb{R}_+^N | \xi_N = 0, \xi_1 = 0\}$ . By the Arzelá-Ascoli Theorem,  $\tilde{v}_k$  converges to  $\tilde{v}$  in  $C^2(K) \cap C(\bar{D})$ . In addition, by a diagonalization argument,  $\tilde{v}_k$  converges locally in  $\overline{\mathbb{R}_+^N}$  to  $\tilde{v} \in C^2(\mathbb{R}_+^N) \cap C(\overline{\mathbb{R}_+^N})$  such that  $\tilde{v}$  satisfies

$$(3.7) \quad \begin{cases} -\Delta \tilde{v} = \tilde{v}^p & \text{in } \mathbb{R}_+^N, \\ \tilde{v}(\xi) \leq 2^{\frac{2}{p-1}} & \text{in } \mathbb{R}_+^N, \\ \tilde{v} = 0 & \text{on } \Gamma_0 = \{\xi \in \mathbb{R}_+^N | \xi_N = 0, \xi_1 > 0\}, \\ \frac{\partial \tilde{v}}{\partial \xi_N} = 0 & \text{on } \Gamma_1 = \{\xi \in \mathbb{R}_+^N | \xi_N = 0, \xi_1 < 0\}. \end{cases}$$

Deduced from Lemma 4, we have  $\tilde{v} \equiv 0$ . However, from (3.5),  $\tilde{v}(d_k e_N) = 1$ . Clearly, a contradiction is achieved. Therefore, the proof Theorem 4 is complete.  $\square$

#### 4. Elliptic systems

In the last section, we consider the elliptic systems. We could show that boundedness is removable in the Liouville theorem of (1.12). The following lemma is established in [DS].

LEMMA 5. *The system (1.12) does not have a bounded positive solution in  $\mathbb{R}^N$ , provided*

$$1 < \alpha_1, \alpha_2 < \frac{N+2}{N-2}.$$

PROOF OF THEOREM 5. Suppose by contradiction that there exists a sequence of  $(y_k)$  such that  $u_1(y_k) \rightarrow \infty$  or  $u_2(y_k) \rightarrow \infty$  as  $k \rightarrow \infty$ . Let

$$M(y) := u_1^{\frac{\alpha_1-1}{2}}(y) + u_2^{\frac{\alpha_2-1}{2}}(y).$$

Then,  $M(y_k) \rightarrow \infty$  as  $k \rightarrow \infty$ . From the doubling lemma by letting  $D = \Sigma = X = \overline{\mathbb{R}^N}$  and Remark 1, there exists a sequence of  $(x_k)$  such that

$$M(x_k) \geq M(y_k)$$

and

$$M(z) \leq 2M(x_k), \quad \text{if } |z - x_k| \leq kM^{-1}(x_k).$$

Let  $\alpha = \frac{2}{\alpha_1-1}$  and  $\beta = \frac{2}{\alpha_2-1}$ . We rescale  $(u_1, u_2)$  by

$$\tilde{v}_{1,k}(\xi) := \frac{u_1(x_k + \frac{\xi}{M(x_k)})}{M^\alpha(x_k)}$$

and

$$\tilde{v}_{2,k}(\xi) := \frac{u_2(x_k + \frac{\xi}{M(x_k)})}{M^\beta(x_k)}.$$

Then,  $(\tilde{v}_{1,k}, \tilde{v}_{2,k})$  satisfies

$$(4.1) \quad \begin{cases} \Delta \tilde{v}_{1,k} + \tilde{v}_{1,k}^{\alpha_1} + \tilde{v}_{2,k}^{\alpha_1 \frac{\alpha_2-1}{\alpha_1-1}} = 0, \\ \Delta \tilde{v}_{2,k} + \tilde{v}_{2,k}^{\alpha_2} + \tilde{v}_{1,k}^{\alpha_2 \frac{\alpha_1-1}{\alpha_2-1}} = 0. \end{cases}$$

Moreover

$$\tilde{v}_{1,k}^{\frac{1}{\alpha}}(0) + \tilde{v}_{2,k}^{\frac{1}{\beta}}(0) = 1$$

and

$$\tilde{v}_{1,k}^{\frac{1}{\alpha}}(\xi) + \tilde{v}_{2,k}^{\frac{1}{\beta}}(\xi) \leq 2, \quad \forall |\xi| \leq k.$$

For any smooth compact  $D$  in  $\mathbb{R}^N$ , there exists  $k_0$  large enough such that  $D \subset B_k(0)$  if  $k \geq k_0$ . Using the classical  $W^{2,q}$  estimates for elliptic equation, we have

$$\sum_{i=1}^2 \|\tilde{v}_{i,k}\|_{W^{2,q}(D)} \leq C$$

for  $1 < q < \infty$ . By the standard Sobolev imbedding theorem, there exists a sequence of  $(\tilde{v}_{1,k}, \tilde{v}_{2,k})$  that converges to  $(\tilde{v}_1, \tilde{v}_2)$  in  $C^{1,\tau}(D)$  for some  $\tau > 0$ . Employing a diagonal line argument, we readily deduce that  $(\tilde{v}_{1,k}, \tilde{v}_{2,k})$  converges in  $C_{loc}^{1,\tau}(\mathbb{R}^N)$  to a solution  $(\tilde{v}_1, \tilde{v}_2)$  in  $\mathbb{R}^N$  which satisfies

$$(4.2) \quad \begin{cases} \Delta \tilde{v}_1 + \tilde{v}_1^{\alpha_1} + \tilde{v}_2^{\alpha_1 \frac{\alpha_2-1}{\alpha_1-1}} = 0, \\ \Delta \tilde{v}_2 + \tilde{v}_2^{\alpha_2} + \tilde{v}_1^{\alpha_2 \frac{\alpha_1-1}{\alpha_2-1}} = 0. \end{cases}$$

Furthermore,

$$\tilde{v}_1^{\frac{1}{\alpha}}(0) + \tilde{v}_2^{\frac{1}{\beta}}(0) = 1.$$

So  $(\tilde{v}_1, \tilde{v}_2)$  is nontrivial. This contradicts Lemma 5. Therefore, we then complete the proof of the theorem.  $\square$

**Acknowledgement:** Part of the work was done when the first author was visiting Beijing Normal University in China.

## References

- [ADN] S. Agmon, A. Douglis and L. Nirenberg, Estimates near the boundary for solutions of elliptic partial differential equation satisfying general boundary conditions, I. *Comm. Pure Appl. Math.* 12(1959), 623-729.
- [BCN] H. Berestycki, L. Caffarelli and L. Nirenberg, Further qualitative properties for elliptic equation in unbounded domains. Dedicated to Ennio De Giorgi. *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* 25(1997), 69-94.
- [BM] I. Birindelli and E. Mitidieri, Liouville theorems for elliptic inequalities and applications, *Proc. Roy. Soc. Edinburgh* 128A(1998), 1217-1247.
- [BMa] J. Busca and R. Manasevich, A Liouville-type theorem for Lane-Emden systems, *Indiana Univ. Math. J.* 51 (2002), no. 1, 37-51.
- [CDM] G. Caristi, L. D'Ambrosio and E. Mitidieri, Representation formulae for solutions to some classes of higher order systems and related Liouville theorems. *Milan J. Math.* 76 (2008), 27-67.
- [CL] W. Chen and C. Li, Classification of solutions of some nonlinear elliptic equations, *Duke Math. J.* 63(1991), 615-622.

- [CLO] W. Chen, C. Li and B. Ou, Classification of solutions for an integral equation, *Comm. Pure Appl. Math.* 59 (2006), no. 3, 330–343.
- [CP] E. Colorado and I. Peral, Semilinear elliptic problems with mixed Dirichlet-Neumann boundary conditions, *Journal of Functional Analysis* 199(2003), 468-507.
- [D] E. Dancer, Some notes on the method of moving planes, *Bull. Austral. Math. Soc.* 46(1992), 425-434.
- [DG] L. Damascelli and F. Gladiali, Some nonexistence results for positive solutions of elliptic equations in unbounded domains, *Rev. Mat. Iberoamericana* 20(2004), 67-86.
- [DLN] D. G. deFigueiredo, P. L. Lions and R. D. Nussbaum, A priori estimate and existence of positive solution to semilinear elliptic equations, *J. Math. Pures at Appl.* 61(1982), 41-63.
- [DS] D. de Figueiredo and B. Sirakov, Liouville type theorem, monotonicity results and a priori bounds for positive solution of elliptic systems, *Math. Ann.* 33(2005), no. 2, 231-260.
- [GL] Y. Guo and J. Liu, Liouville-type theorems for polyharmonic equations in  $\mathbb{R}^N$  and in  $\mathbb{R}_+^N$ , *Proceedings of the Royal Society of Edinburgh* 138A (2008), 339-359.
- [LGZ] J. Liu, Y. Guo and Y. Zhang, Liouville-type theorems for polyharmonic systems in  $R^N$ , *J. Differential Equations*, 225 (2006), 685-709.
- [GNN] B. Gidas; W. Ni and L. Nirenberg, Symmetry and related properties via the maximum principle. *Comm. Math. Phys.* 68 (1979), no. 3, 209-243.
- [GS1] B. Gidas and J. Spruck, Global and local behavior of positive solutions of nonlinear elliptic equations, *Commun. Pure Appl. Math.* 34(1981), 525-598.
- [GS2] B. Gidas and J. Spruck, A priori bounds for positive solutions for nonlinear elliptic equations, *Commun. Partial Diff. Equations* 6(1981), 883-901.
- [LZ] G. Lu and J. Zhu, The axial symmetry and regularity of solutions to an integral equation in a half space, *Pacific Journal of Mathematics* 253 (2011), No. 2, 455-473.
- [M] E. Mitidieri, Nonexistence of positive solutions of semilinear elliptic systems in  $\mathbb{R}^N$ , *Differ. Integral Equations* 9 (1996), 465-479.
- [M1] E. Mitidieri, A Rellich type identity and applications. *Comm. Partial Differential Equations* 18 (1993), no. 1-2, 125-151.
- [PQS] P. Poláčik, P. Quittner and P. Souplet, Singularity and decay estimates in superlinear problems via Liouville-type theorems, *Duke Math. J.* 139 (2007), 555-579.
- [RW] W. Reichel and T. Weth, existence of solutions to nonlinear, subcritical higher order elliptic Dirichlet problems, *J. Differential Equations* 248(2010), 1866-1878.
- [SZ] J. Serrin, H. Zou, Non-existence of positive solutions of Lane-Emden systems. *Differential Integral Equations* 9 (1996), no. 4, 635-653.
- [S] B. Sirakov, existence results and a priori bounds for higher order elliptic equations and systems. *J. Math. Pures Appl.* 89 (2008), 114-133.
- [So] P. Souplet, The proof of the Lane-Emden conjecture in four space dimensions, *Adv. Math.* 221(2009), no. 5, 1409-1427.
- [WX] J. Wei and X. Xu. Classification of solutions of higher order conformally invariant equations, *Math. Ann.* 313(1999), 207-228.

GUOZHEN LU, SCHOOL OF MATHEMATICAL SCIENCES, BEIJING NORMAL UNIVERSITY, BEIJING, CHINA 100875, AND, DEPARTMENT OF MATHEMATICS, WAYNE STATE UNIVERSITY, DETROIT, MI 48202, USA, EMAILS: GZLU@MATH.WAYNE.EDU

PEIYONG WANG AND JIUYI ZHU, DEPARTMENT OF MATHEMATICS, WAYNE STATE UNIVERSITY, DETROIT, MI 48202, USA, EMAILS: PYWANG@MATH.WAYNE.EDU AND JIUYI.ZHU@WAYNE.EDU